



# National Inventory Document 2025

**SLOVAK REPUBLIC**

Submission under the Regulation (EU) 2018/19

**Bratislava  
March 2025**

## PREFACE

<b>TITLE OF REPORT</b>	<b>NATIONAL GREENHOUSE GAS INVENTORY DOCUMENT 1990 – 2023 UNDER THE REGULATION (EU) 2018/1999</b>
<b>CONTACTS:</b>	Compiled by Janka Szemesová Coordinator of the National Inventory System of the Slovak Republic
	Approved by Jozef Škultéty Head of the Climate Change Department of the Ministry of Environment of the Slovak Republic National Focal Point to the UNFCCC
<b>ORGANISATION:</b>	Slovak Hydrometeorological Institute
	Ministry of Environment of the Slovak Republic
<b>ADDRESS:</b>	Jeséniova 17, Bratislava
	Nám. Ľudovíta Štúra 1, Bratislava
<b>EMAILS:</b>	<a href="mailto:janka.szemesova@shmu.sk">janka.szemesova@shmu.sk</a>
	<a href="mailto:jozef.skultety@enviro.gov.sk">jozef.skultety@enviro.gov.sk</a>
<b>VERSION:</b>	2. version
<b>DATE OF SUBMISSION:</b>	March, 2025

In 2025, the Slovak Republic is submitting report to the European Commission under the Article 26 (3) of the Regulation (EU) 2018/1999 and the Chapter III and Articles 8-24 of the Commission Implementing Regulation (EU) 2020/1208. The whole package of the National Inventory Document 2025 of the Slovak Republic comprises:

1. SVK NID 2025 – Sectoral chapters of Slovakia's National Greenhouse Gas Emission Inventory Document prepared using the MPG under ETF;
2. SVK\_GHG\_1990-2023 – SVK-CRT-2025-V0.2 (2025) generated using the ETF software according to the decision 18/CMA.1 (Annex 1), accompanied by the json file;
3. Tabular format specified in Articles 10-15, 17, 19-24 to the 2020/1208 Regulation, including LULUCF accounting tables.

The Slovakia Inventory Document as well as CRT tables can be downloaded from the following address: <http://oeab.shmu.sk>. GHG emissions are also published in publication Životné prostredie v SR (Chapter 1.3 Air, page 19) prepared by the Statistical Office of the Slovak Republic.

This version of the annual GHG emissions inventory is the second submission of the national inventory in 2025 of the Slovak Republic to the European Union under the Energy Governance.

Submission is uploaded via the EIONET Central Data Repository tool of the EEA.

### Key points:

The decrease in greenhouse gas emissions in Slovakia exceeded 50% in 2023 compared to the base year 1990. The emissions from road transport decreased year-on-year by 0.6%, marking the first such decline since 1993 without being affected by any global situation or crisis. In addition, it is significant that the decreasing trend in emissions from landfilling was confirmed, for the second year in a row.

The statistical data collected for year 2023 by individual sectors and categories confirmed the decreasing trend of emissions in the energy industry and manufacturing, in services, and in households. Industrial sectors reduced greenhouse gas emissions by 3% year-on-year, mainly in the mineral and chemical industries. In contrast, nitrous oxide emissions increased in agriculture by 10% year-on-year, mainly due to the intensification of agricultural soil fertilization. Emissions from waste management decreased by 4% year-on-year. Removals from forests and soil increased slightly, which is, however, only a very temporary trend. In the long term, a decrease in removals in Slovakia is expected due to the aging forest.

Emissions from road transportation in Slovakia wasn't successfully reduced for the long time. Despite some measures that were implemented in the past, they appeared to be ineffective in reversing the trends of increasing diesel oil and gasoline consumption. In 2023, diesel oil consumption decreased for the first time since the early 1990s and emissions decreased slightly. It will be interesting to see whether this trend is confirmed in the following years, as the first estimate for 2024 will be available this summer. Overall, greenhouse gas emissions in 2023 reached 36 114.74 Gg CO<sub>2</sub> equivalents without LULUCF removals and including indirect emissions from solvent use.

Major changes and corrections included in this SVK NID 2025 are connected with the implementation of the IPCC 2019 Refinement and are focused on following issues:

- General: Harmonization of indirect emissions of the NO<sub>x</sub>, CO, NMVOC, SO<sub>2</sub> and NH<sub>3</sub> in line with the CLRTAP and NECD submissions reported in February 15, 2025 in all sectors ([Chapter ES.5](#)).
- Energy: Road transport was recalculated due to the COPERT model update. In addition, based on the new statistical input in households, biomass consumption in the 1.A.4.b was recalculated.
- Fugitive Emissions: This subcategory was recalculated based on the changes in EF and corrections, mainly caused by implementation of the IPCC 2019 Refinement.
- IPPU: No major recalculations needed in this sector except of correction of the EF and addition new vehicle categories due to the COPERT model update in 2.D.3.
- Agriculture: Several recalculations were occurred, as addition emissions from rabbits.
- LULUCF: No major recalculations needed in this sector except of corrections of calculation errors.
- Waste: Several recalculations in all categories connected with the implementation of the IPCC 2019 Refinement and the changes in activity data and methodologies.

More information on recalculations made in the GHG inventory preparation can be found in the sectoral chapters of this Report and the [Chapter 10](#).

---

<b>PREFACE.....</b>	<b>2</b>
<b>EXECUTIVE SUMMARY .....</b>	<b>5</b>
ES.1. Background Information on Greenhouse Gas Inventories and Climate Change ....	5
ES.2. Summary of National Emission and Removal Trends .....	6
ES.3. Overview of Source and Sink Category .....	8
ES.4. Indirect Emissions and Precursors of Greenhouse Gases .....	15
<b>CHAPTER 1. INTRODUCTION .....</b>	<b>17</b>
1.1. Background Information on Greenhouse Gas Inventories and Climate Change ..	17
1.2. Description of the National Inventory Arrangements .....	22
<b>CHAPTER 2. TRENDS IN GREENHOUSE GAS EMISSIONS .....</b>	<b>38</b>
2.1. Description and Interpretation of Emission Trends for Aggregated GHG Emissions .....	38
2.2. Description and Interpretation of Emission Trends by Gas .....	40
2.3. Description and Interpretation of Emission Trends by Category.....	41
<b>CHAPTER 3. ENERGY.....</b>	<b>49</b>
<b>CHAPTER 4. IPPU.....</b>	<b>127</b>
<b>CHAPTER 5. AGRICULTURE.....</b>	<b>231</b>
<b>CHAPTER 6. LULUCF.....</b>	<b>308</b>
<b>CHAPTER 7. WASTE.....</b>	<b>383</b>
<b>CHAPTER 8. OTHER.....</b>	<b>427</b>
<b>CHAPTER 9. INDIRECT CO<sub>2</sub> AND NITROUS OXIDE EMISSIONS.....</b>	<b>427</b>
<b>CHAPTER 10. RECALCULATIONS AND IMPROVEMENTS.....</b>	<b>427</b>
<b>CHAPTER 11. INFORMATION ON CHANGES IN NATIONAL SYSTEM.....</b>	<b>434</b>
<b>CHAPTER 12. INFORMATION IN ACCORDANCE WITH ART. 3, PARA 14.....</b>	<b>434</b>

---

# EXECUTIVE SUMMARY

## ES.1. Background Information on Greenhouse Gas Inventories and Climate Change

Climate change is a key environmental, economic and social challenge globally and in Europe. On the one hand, most economic activities are contributing to climate change by emitting greenhouse gases or affecting carbon sinks (e.g. through land use change); on the other hand, all ecosystems, many economic activities as well as human health and well-being are sensitive to climate change.

Because the impact of the climate change differs in various regions of the world, its socio-economic and environmental impact always requires an active solution. Necessary political measures have to stem from detailed analysis of the current greenhouse gas (GHG) emissions in every sector, emission projections and impact assessment of adopted or planned policy measures. Such detailed analyses are good starting points for any policy reflected in national communication of a party prepared according to rules of the United Nations Framework Convention on Climate Change (UNFCCC).

Climate change, caused by increasing anthropogenic emissions of greenhouse gases, represents one of the most serious environmental threats for humankind. Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) are the most important anthropogenic greenhouse gases with increasing concentration in atmosphere. The GHGs inventory includes also halogenated hydrocarbons (PFCs, HFCs) and SF<sub>6</sub>, which are not controlled by the Montreal Protocol.

According to the WMO it is officially confirmed that 2024 is the warmest year on record, by a huge margin. The annual average global temperature approached 1.5° Celsius above pre-industrial levels – symbolic because the Paris Agreement on climate change aims to limit the long-term temperature increase (averaged over decades rather than an individual year like 2024) to no more than 1.5° Celsius above pre-industrial levels. Six leading international datasets used for monitoring global temperatures and consolidated by WMO show that the annual average global temperature was 1.45 ± 0.12 °C above pre-industrial levels (1850-1900) in 2024. Global temperatures in every month between June and December set new monthly records. July and August were the two hottest months on record. Long-term monitoring of global temperatures is just one indicator of climate and how it is changing. Other key indicators include atmospheric greenhouse gas concentrations, ocean heat and acidification, sea level, sea ice extent and glacier mass balance.

WMO's provisional State of the Global Climate in 2024 report, published on 30 November, showed that records were broken across the board. WMO will issue its final State of the Global Climate 2024 report in March 2025. This will include details on socio-economic impacts on food security, displacement and health.

The Paris Agreement seeks to hold the increase in the global average temperature to well below 2°C above pre-industrial levels while pursuing efforts to limit the temperature increase to 1.5°C above pre-industrial levels. The Intergovernmental Panel on Climate Change says that climate-related risks for natural and human systems are higher for global warming of 1.5°C than at present, but lower than at 2°C. A study by WMO and the UK's Met Office last year predicted that there is a 66% likelihood that the annual average near-surface global temperature between 2023 and 2027 will be more than 1.5°C above pre-industrial levels for at least one year. This does not mean that we will permanently exceed the 1.5°C level specified in the Paris Agreement which refers to long-term warming over many years. The chance

---

of temporarily exceeding 1.5°C has risen steadily since 2015, when it was close to zero. For the years between 2017 and 2021, there was a 10% chance of exceedance.<sup>1</sup>

The European Climate Law writes into law the goal set out in the European Green Deal for Europe's economy and society to become climate-neutral by 2050. The law also sets the intermediate target of reducing net greenhouse gas emissions by at least 55% by 2030, compared to 1990 levels. Climate neutrality means achieving net zero greenhouse gas emissions for EU countries as a whole, mainly by cutting emissions, investing in green technologies and protecting the natural environment. The law aims to ensure that all EU policies contribute to this goal and that all sectors of the economy and society play their part. The Climate Law includes measures to keep track of progress and adjust our actions accordingly, based on existing systems such as the governance process for Member States' integrated national energy and climate plans, regular reports by the European Environment Agency, and the latest scientific evidence on climate change and its impacts. Slovakia is a part of these actions and agreed the climate neutrality until 2050 among the first countries in the EU (end of 2019).

During the year 2020, many countries were going through the worst economic contraction since the 1930s due to COVID-19 pandemic. Some economists believe it will be essentially V-shaped: first a steep fall, then a steep return to normal. In May 2020, the EU Commission proposed stimulus packages called "sustainable recovery" mostly address to investments into the buildings, transport, power and industry sectors. Aim of this plan is not only reduce emissions, but also create new jobs, make innovations and build circular economy.

On 14 July 2021, the European Commission adopted a series of legislative proposals setting out how it intends to achieve climate neutrality in the EU by 2050, including the intermediate target. The package proposes to revise several pieces of EU climate legislation, including the ETS directive, Effort Sharing Regulation, transport and land use legislation, setting out in real terms the ways in which the Commission intends to reach EU climate targets under the European Green Deal.

From 2021, the fourth EU ETS trading period has gone operational. Main change is the increase of linear reduction factor from 1.74% per annum to 2.2% per annum, which should bring at least 43% reduction within the EU ETS sectors by 2030. To achieve the ambitious reductions, several low carbon-funding mechanisms were introduced, in particular Innovation Fund (to support demonstration of innovative renewable energy and low-carbon innovation in industry, as well as carbon capture, use and storage) and a Modernisation Fund (to contribute to modernising the energy systems of 10 EU Member States with lower GDP).

## ES.2. Summary of National Emission and Removal Trends

The GHG emissions presented in the National Inventory Document 2025 were updated and recalculated using the last updated methods based on the IPCC 2019 Refinement to the IPCC 2006 Guidelines for National Greenhouse Gas Inventories, national conditions and data published by the Statistical Office of the Slovak Republic.

In 2024, UNFCCC review did not take place and all previous recommendation were implemented. ESR review took place in 2024 resulted without open observation and with no recommendation.

Total GHG emissions were 36 114.74 Gg of CO<sub>2</sub> eq. in 2023 (without LULUCF and with indirect emissions). This represents a reduction by 50.9% against the base year 1990. In comparison with 2022, the emissions decreased by 2%. The decrease in total emissions of 2023 compared to 2022 was due to decrease in the Energy and IPPU sectors.

---


<sup>1</sup> <https://wmo.int/media/news/wmo-confirms-2023-smashes-global-temperature-record>

The 2025 submission includes indirect CO<sub>2</sub> emissions in the solvents category (IPPU). This means, that indirect emissions were 41.08 Gg of CO<sub>2</sub> eq. in 2023. Indirect CO<sub>2</sub> emissions were estimated and reported for the time series 1990 – 2023.

The major changes in the 2025 national inventory of GHG emissions are caused by recalculations in the Fugitive Emissions, Agriculture and Waste sectors for the particular years or whole time series.

The emissions with LULUCF and with indirect emissions decreased in 2023 compared to 2022 by 4.4%. During period 1991 – 2023, the total greenhouse gas emissions in the Slovak Republic did not exceed the level of 1990. [Tables ES.2](#) and [ES.3](#) show the aggregated GHG emissions expressed in CO<sub>2</sub> equivalents and according to the gases in the period 1990 – 2023. [Figure ES.1](#) shows trend in the gases without LULUCF. The emissions of F-gases are the only emissions from consumption HFCs, PFCs and SF<sub>6</sub> in industry with the increasing trend since 1990 (despite decrease of PFCs gases from aluminium production).

**Figure ES.1: GHG emission trends compared with the base year (%) in the Slovak Republic**



*GHG emissions in % to base years without LULUCF and indirect emissions; emissions are determined as of March 2025.*

Slovakia decreased its emissions by around 21% between 2010 and 2023. The latest available GHG emission projections have demonstrated emissions decrease as an evidence of the successful implementation of the policies and measures and their effect on the improvement in energy intensity and industrial production efficiency. These emission projections are in line with the [First Biennial Transparency Report](#) (submitted in December 2024). New drivers and parameters reflecting the actual pandemic situation were projected.

Reduction of emissions in Slovakia in past years was a conjunction of different impacts starting from impressive industrial and technological restructuring connected with the fuel switching of fossil fuels from coal and oil to the natural gas (air pollution legislation since 1991 was the main driving force), economy restructuring towards the less energy intensive production (mostly in recent years) and also by temporary changes in production intensity (driven by global and EU markets). Transport (mostly the road transport), with continuously increasing emissions in past, started to decline its trend in the recent years. The continuous pressure is being made in formulating the effective strategy and policy to achieve further reduction of emissions in this sector, too. For example, combination of regulatory and economic instruments (toll pay for freight vehicles based on their environmental characteristics in a combination with fuel and emission standards for new cars). The car tax system and the level of fuel taxation, which is close to the EU average, contribute to limit the increase of greenhouse gas emissions in the transport.

In Slovakia, the structural changes in the manufacturing industry towards less energy intensive industries, such as machinery and automotive industry, can explain why after 2009 the energy consumption did not pick up the same pace as prior to that year when led to a significant decrease in

---

primary energy intensity (the GDP grew twice as fast as primary energy consumption). Therefore, the trend observed particularly in primary energy consumption is mainly due to other factors although some energy efficiency improvements did take place particularly during the period after the year 2012. The policy package still needs various improvements across the sectors including the sectoral mitigation targets particularly in transport, buildings, agriculture and waste.

Although this optimistic trend recognised in previous years, it is visible since last 3 years, that the improvement of several indicators such as GHG per capita or GHG/GDP started slowed down and reached minimum level. GHG emissions level reached minimum in 2014 (since 1990), then the trend started to increase slowly until peaked in 2018. After post pandemic recovery (2021), GHG emissions were the lowest since the base in 2022 and 2023 ([Chapter 2](#)).

Covid-19 pandemic situation connected with emissions drop in 2020 in conjunction with the industrial changes in iron and steel production, transformation of electricity and heat production sectors and changes in fuels combustion caused by increasing prices, led to the dramatically high decrease of the total emissions in 2020. However, despite this optimistic development, the emission trend in 2021 increased back to the pre-pandemic level. Further reduction of emissions in 2023 was caused by the energy prices policy and due to economic reasons, several important industrial plants reduced or closed the operation. More information are in energy and IPPU sectoral chapters.

### ES.3. Overview of Source and Sink Category

The emissions without LULUCF in 2023 and with indirect emissions are lower than in 2022 due to the essential decrease in Energy and IPPU sectors, mostly in energy and manufacturing industry, mineral production, chemical industry and metal industry.

The Energy sector (including transport) with the share of 69% was the main contributor to total GHG emissions in 2023. Transport with 21% share on total emissions contributes significantly to the GHG budget. In 2023, the transport in total emissions has decreased by more than 0.55% in comparison with previous year 2022. In addition to fuel combustion in stationary sources of pollution, also the pollution from small sources of residential heating systems and fugitive methane emissions from transport, processing and distribution of oil and natural gas contribute significantly to the total GHG emissions.

The Industrial Processes and Product Use sector was the second important sector in 2023 with its 20% share in total GHG emissions, producing mainly technological emissions from processing mineral products, chemical production and steel and iron production. The reduction of emissions from technological processes is very costly and there exist specific technical limits, therefore the emissions have not been changed since the reference year as significantly as for other categories. Mostly the production volume in industrial processes influences their level. The most growing emissions within the IPPU sector are HFCs and SF<sub>6</sub> emissions as result of industrial demand and use of these substances in construction, insulation of building, electro-technical and/or automobile industry.


In 2023, the share of the Agriculture sector on total GHG emissions was 6% and the trend in emissions is slightly decreasing since 1999. The most significant reduction of emissions from agriculture was achieved at the beginning of nineties as result of reduction in breeding livestock number together with restricted use of fertilizers.

The Waste sector contributed by 5% to total GHG emissions in 2023. Using of more exact methodology for the evaluation of methane emissions from solid waste disposal on sites and included also older layer into calculation resulted in continual increase of emissions by more than 100% compared to the base year 1990. Similar trend is expected to remain in future years, although the increase should not be so substantial as before. Volume of emissions from landfills depends, largely, on applied methodology to evaluate landfills and on the scale of implementation energy recovery of landfill gases by landfill operators.



The shares of individual sectors in total GHG emissions have not been changed significantly compared to the base year 1990. Nevertheless, increase in transport emissions in trend since 1990 and decreased share of stationary sources of pollution in the Energy sector are noticeable. Combustion of fossil fuels, which account for about 75% of the total CO<sub>2</sub> emissions in the Slovak Republic (without LULUCF), represent the most important anthropogenic source of CO<sub>2</sub> emissions (*Figure ES.2, Table ES.3*).

*Figure ES.2: GHG emissions share by the sectors (%) in the Slovak Republic in 2023*



*Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.*

**Table ES.1: Summary of the GHG emissions according to the gases and the sectors in 2023 and 2022**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2023					
	Gg of CO <sub>2</sub> equivalents					
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	SF <sub>6</sub>
1. Energy	23 972.41	782.47	172.49	NO	NO	NO
2. Industrial Processes	6 718.99	16.06	104.95	437.89	0.01	14.70
3. Agriculture	72.47	1 138.58	970.80	NO	NO	NO
4. LULUCF	-7 817.81	15.14	27.05	NO	NO	NO
5. Waste	2.38	1 472.24	197.24	NO	NO	NO
<i>Memo Items - International Transport</i>	169.66	0.09	1.22	NO	NO	NO
<b>Total (excluding LULUCF)</b>	30 766.25	3 409.35	1 445.47	437.89	0.01	14.70
<b>Total (including LULUCF)</b>	22 948.43	3 424.49	1 472.52	437.89	0.01	14.70

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2022					
	Gg of CO <sub>2</sub> equivalents					
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	SF <sub>6</sub>
1. Energy	24 609.40	791.31	187.43	NO	NO	NO
2. Industrial Processes	6 915.91	15.02	111.05	480.86	5.91	15.38
3. Agriculture	60.84	1 139.85	765.28	NO	NO	NO
4. LULUCF	-7 317.22	45.85	44.20	NO	NO	NO
5. Waste	3.31	1 537.32	202.05	NO	NO	NO
<i>Memo Items - International Transport</i>	148.02	0.08	1.07	NO	NO	NO
<b>Total (excluding LULUCF)</b>	31 589.46	3 483.50	1 265.82	480.86	5.91	15.38
<b>Total (including LULUCF)</b>	24 272.24	3 529.35	1 310.02	480.86	5.91	15.38

**Table ES.2: Summary of the GHG emissions according to the gases in 1990 – 2021**

GREENHOUSE GAS EMISSIONS	Base year 1990	1991	1992	1993	1994	1995	1996	1997
	<i>Gg of CO<sub>2</sub> equivalents</i>							
CO <sub>2</sub> emissions excluding net CO <sub>2</sub> from LULUCF	61 528.54	53 333.93	48 930.98	46 394.33	43 803.52	44 194.17	44 075.52	44 147.08
CO <sub>2</sub> emissions including net CO <sub>2</sub> from LULUCF	52 505.31	43 499.03	38 463.07	36 120.97	34 124.52	35 073.46	35 046.20	35 340.74
CH <sub>4</sub> emissions excluding CH <sub>4</sub> from LULUCF	8 274.46	7 854.76	6 747.16	6 160.48	5 802.18	5 800.89	5 633.63	5 353.11
CH <sub>4</sub> emissions including CH <sub>4</sub> from LULUCF	8 286.91	7 865.06	6 761.48	6 187.28	5 809.41	5 809.54	5 645.67	5 362.47
N <sub>2</sub> O emissions excluding N <sub>2</sub> O from LULUCF	3 408.58	2 692.28	2 239.11	1 881.33	2 284.33	2 427.11	2 600.79	2 577.68
N <sub>2</sub> O emissions including N <sub>2</sub> O from LULUCF	3 526.84	2 802.30	2 348.48	1 994.52	2 382.58	2 512.13	2 681.83	2 649.92
HFCs	NO	NO	NO	NO	0.20	12.38	26.31	38.33
PFCs	213.92	210.43	195.83	122.51	104.11	90.15	36.89	36.48
SF <sub>6</sub>	0.06	0.04	0.04	0.09	18.16	10.47	11.51	11.83
<b>Total (excluding LULUCF)</b>	<b>73 425.57</b>	<b>64 091.45</b>	<b>58 113.12</b>	<b>54 558.74</b>	<b>52 012.51</b>	<b>52 535.17</b>	<b>52 384.64</b>	<b>52 164.51</b>
<b>Total (including LULUCF)</b>	<b>64 533.04</b>	<b>54 376.85</b>	<b>47 768.91</b>	<b>44 425.38</b>	<b>42 438.99</b>	<b>43 508.13</b>	<b>43 448.40</b>	<b>43 439.76</b>
<i>Total (excluding LULUCF, including indirect emissions)</i>	<i>73 513.34</i>	<i>64 177.97</i>	<i>58 198.55</i>	<i>54 642.99</i>	<i>52 095.68</i>	<i>52 617.26</i>	<i>52 465.61</i>	<i>52 244.35</i>
<i>Total (including LULUCF, including indirect emissions)</i>	<i>64 620.81</i>	<i>54 463.38</i>	<i>47 854.33</i>	<i>44 509.63</i>	<i>42 522.16</i>	<i>43 590.22</i>	<i>43 529.37</i>	<i>43 519.59</i>

GREENHOUSE GAS EMISSIONS	1998	1999	2000	2001	2002	2003	2004	2005
	<i>Gg of CO<sub>2</sub> equivalents</i>							
CO <sub>2</sub> emissions excluding net CO <sub>2</sub> from LULUCF	43 877.53	43 088.35	41 187.04	43 272.72	42 033.70	42 359.23	42 856.56	42 857.27
CO <sub>2</sub> emissions including net CO <sub>2</sub> from LULUCF	34 010.51	33 986.98	32 171.98	34 966.33	33 238.82	34 109.14	34 644.52	38 517.34
CH <sub>4</sub> emissions excluding CH <sub>4</sub> from LULUCF	5 156.12	5 084.02	4 849.45	4 737.91	4 680.74	4 626.59	4 582.65	4 582.93
CH <sub>4</sub> emissions including CH <sub>4</sub> from LULUCF	5 165.31	5 144.36	4 879.79	4 752.00	4 704.30	4 672.53	4 598.58	4 612.38
N <sub>2</sub> O emissions excluding N <sub>2</sub> O from LULUCF	2 300.56	1 929.07	2 129.10	2 392.60	2 334.25	2 351.62	2 520.67	2 507.81
N <sub>2</sub> O emissions including N <sub>2</sub> O from LULUCF	2 366.73	2 018.75	2 192.90	2 442.44	2 384.67	2 410.01	2 563.20	2 554.02
HFCs	50.73	71.82	98.20	130.29	167.96	201.17	240.28	277.09
PFCs	28.34	19.03	17.83	18.84	19.87	26.55	27.00	27.48
SF <sub>6</sub>	13.04	13.03	13.44	13.74	15.23	15.52	15.91	16.89
<b>Total (excluding LULUCF)</b>	<b>51 426.30</b>	<b>50 205.32</b>	<b>48 295.06</b>	<b>50 566.10</b>	<b>49 251.74</b>	<b>49 580.67</b>	<b>50 243.07</b>	<b>50 269.48</b>

GREENHOUSE GAS EMISSIONS	1998	1999	2000	2001	2002	2003	2004	2005
	<i>Gg of CO<sub>2</sub> equivalents</i>							
<b>Total (including LULUCF)</b>	<b>41 634.65</b>	<b>41 253.97</b>	<b>39 374.15</b>	<b>42 323.64</b>	<b>40 530.84</b>	<b>41 434.91</b>	<b>42 089.49</b>	<b>46 005.21</b>
<i>Total (excluding LULUCF, including indirect emissions)</i>	<i>51 505.00</i>	<i>50 282.12</i>	<i>48 360.51</i>	<i>50 631.62</i>	<i>49 323.51</i>	<i>49 648.66</i>	<i>50 318.75</i>	<i>50 336.41</i>
<b>Total (including LULUCF, including indirect emissions)</b>	<b>41 713.36</b>	<b>41 330.77</b>	<b>39 439.59</b>	<b>42 389.16</b>	<b>40 602.60</b>	<b>41 502.90</b>	<b>42 165.17</b>	<b>46 072.14</b>

GREENHOUSE GAS EMISSIONS	2006	2007	2008	2009	2010	2011	2012	2013
	<i>Gg of CO<sub>2</sub> equivalents</i>							
<b>CO<sub>2</sub> emissions excluding net CO<sub>2</sub> from LULUCF</b>	42 639.35	41 031.85	41 425.30	37 681.13	38 464.35	38 045.11	35 961.41	35 621.71
<b>CO<sub>2</sub> emissions including net CO<sub>2</sub> from LULUCF</b>	35 224.70	34 047.68	35 554.50	32 026.41	33 704.86	32 916.83	29 694.28	28 636.37
<b>CH<sub>4</sub> emissions excluding CH<sub>4</sub> from LULUCF</b>	4 590.35	4 388.57	4 319.33	4 119.24	4 124.95	4 059.04	3 912.04	3 896.32
<b>CH<sub>4</sub> emissions including CH<sub>4</sub> from LULUCF</b>	4 608.96	4 419.20	4 338.54	4 147.59	4 147.38	4 085.98	3 963.59	3 913.40
<b>N<sub>2</sub>O emissions excluding N<sub>2</sub>O from LULUCF</b>	2 583.31	2 503.41	2 497.31	2 171.67	2 220.02	1 588.93	1 441.59	1 465.42
<b>N<sub>2</sub>O emissions including N<sub>2</sub>O from LULUCF</b>	2 621.29	2 545.69	2 530.94	2 209.27	2 253.42	1 624.63	1 490.64	1 496.95
<b>HFCs</b>	323.94	368.16	431.50	492.20	569.22	576.43	602.07	620.99
<b>PFCs</b>	40.96	31.39	41.43	24.50	28.27	24.63	28.62	17.02
<b>SF<sub>6</sub></b>	17.22	17.93	19.43	20.11	20.23	21.44	21.90	22.99
<b>Total (excluding LULUCF)</b>	<b>50 195.13</b>	<b>48 341.32</b>	<b>48 734.30</b>	<b>44 508.84</b>	<b>45 427.05</b>	<b>44 315.58</b>	<b>41 967.63</b>	<b>41 644.45</b>
<b>Total (including LULUCF)</b>	<b>42 837.07</b>	<b>41 430.06</b>	<b>42 916.34</b>	<b>38 920.07</b>	<b>40 723.38</b>	<b>39 249.94</b>	<b>35 801.10</b>	<b>34 707.72</b>
<i>Total (excluding LULUCF, including indirect emissions)</i>	<i>50 266.70</i>	<i>48 398.28</i>	<i>48 797.00</i>	<i>44 567.68</i>	<i>45 476.25</i>	<i>44 373.20</i>	<i>42 014.11</i>	<i>41 690.86</i>
<b>Total (including LULUCF, including indirect emissions)</b>	<b>42 908.64</b>	<b>41 487.01</b>	<b>42 979.04</b>	<b>38 978.91</b>	<b>40 772.59</b>	<b>39 307.55</b>	<b>35 847.59</b>	<b>34 754.13</b>

GREENHOUSE GAS EMISSIONS	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Gg of CO<sub>2</sub> equivalents</i>							
<b>CO<sub>2</sub> emissions excluding net CO<sub>2</sub> from LULUCF</b>	33 708.17	34 527.85	34 977.55	36 183.94	36 184.07	33 853.19	31 176.16	35 251.32
<b>CO<sub>2</sub> emissions including net CO<sub>2</sub> from LULUCF</b>	28 891.15	29 224.94	29 603.59	30 914.67	31 888.48	28 788.22	23 932.38	27 987.54
<b>CH<sub>4</sub> emissions excluding CH<sub>4</sub> from LULUCF</b>	3 708.40	3 771.18	3 758.99	3 758.36	3 685.12	3 618.21	3 599.22	3 561.56
<b>CH<sub>4</sub> emissions including CH<sub>4</sub> from LULUCF</b>	3 733.74	3 799.64	3 782.50	3 784.48	3 710.90	3 648.43	3 626.55	3 581.35
<b>N<sub>2</sub>O emissions excluding N<sub>2</sub>O from LULUCF</b>	1 563.37	1 364.10	1 548.39	1 422.49	1 381.11	1 431.13	1 458.90	1 359.98
<b>N<sub>2</sub>O emissions including N<sub>2</sub>O from LULUCF</b>	1 599.88	1 403.90	1 585.59	1 460.81	1 418.86	1 470.70	1 495.55	1 391.15

GREENHOUSE GAS EMISSIONS	2014	2015	2016	2017	2018	2019	2020	2021
	Gg of CO <sub>2</sub> equivalents							
HFCs	626.14	704.84	647.95	710.19	675.62	688.69	646.65	672.37
PFCs	18.27	16.53	15.17	16.75	16.14	14.28	13.22	14.23
SF <sub>6</sub>	14.60	14.75	6.00	7.30	9.68	9.14	17.73	17.44
<b>Total (excluding LULUCF)</b>	<b>39 638.96</b>	<b>40 399.26</b>	<b>40 954.05</b>	<b>42 099.03</b>	<b>41 951.74</b>	<b>39 614.63</b>	<b>36 911.88</b>	<b>40 876.90</b>
<b>Total (including LULUCF)</b>	<b>34 883.79</b>	<b>35 164.60</b>	<b>35 640.80</b>	<b>36 894.20</b>	<b>37 719.68</b>	<b>34 619.45</b>	<b>29 732.08</b>	<b>33 664.08</b>
<i>Total (excluding LULUCF, including indirect emissions)</i>	<i>39 688.50</i>	<i>40 455.60</i>	<i>41 006.56</i>	<i>42 146.51</i>	<i>42 004.85</i>	<i>39 659.93</i>	<i>36 957.76</i>	<i>40 920.57</i>
<i>Total (including LULUCF, including indirect emissions)</i>	<i>34 933.33</i>	<i>35 220.94</i>	<i>35 693.32</i>	<i>36 941.68</i>	<i>37 772.80</i>	<i>34 664.75</i>	<i>29 777.95</i>	<i>33 707.75</i>

Total aggregated GHG emissions, emissions are determined as of March 2025, indirect emissions are reported in the 2025 submission.

**Table ES.3: Summary of the GHG emissions according to the sectors in 1990 – 2021**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Base year (1990)	1991	1992	1993	1994	1995	1996	1997
	Gg of CO <sub>2</sub> equivalents							
1. Energy	56 942.20	50 508.06	45 728.77	41 790.47	39 225.31	38 827.39	38 411.81	38 222.80
2. Industrial Processes	9 427.67	7 225.07	6 844.38	7 886.83	8 171.05	9 028.62	9 405.44	9 445.58
4. Agriculture	5 871.03	5 158.75	4 345.15	3 682.65	3 494.54	3 554.70	3 439.70	3 343.81
5. Land Use, Land-Use Change and Forestry	-8 892.53	-9 714.59	-10 344.22	-10 133.37	-9 573.52	-9 027.04	-8 936.24	-8 724.76
6. Waste	1 184.66	1 199.57	1 194.83	1 198.79	1 121.60	1 124.46	1 127.69	1 152.32

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	1998	1999	2000	2001	2002	2003	2004	2005
	Gg of CO <sub>2</sub> equivalents							
1. Energy	37 699.24	36 973.08	36 086.69	38 036.04	35 702.20	36 617.90	36 156.63	36 675.25
2. Industrial Processes	9 555.64	9 171.96	8 191.62	8 381.92	9 315.95	8 872.47	10 099.17	9 585.07
4. Agriculture	2 992.87	2 863.18	2 787.39	2 892.45	2 888.40	2 723.89	2 576.80	2 603.05
5. Land Use, Land-Use Change and Forestry	-9 791.65	-8 951.35	-8 920.91	-8 242.46	-8 720.91	-8 145.76	-8 153.58	-4 264.26
6. Waste	1 178.55	1 197.10	1 229.36	1 255.69	1 345.20	1 366.42	1 410.47	1 406.11

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2006	2007	2008	2009	2010	2011	2012	2013
	<i>Gg of CO<sub>2</sub> equivalents</i>							
1. Energy	35 920.33	34 258.97	34 678.74	32 106.57	32 409.48	31 856.75	29 591.58	29 413.83
2. Industrial Processes	10 412.72	10 245.83	10 129.73	8 631.05	8 998.01	8 626.83	8 550.70	8 270.40
4. Agriculture	2 379.12	2 418.21	2 472.82	2 274.30	2 474.20	2 236.45	2 205.69	2 322.08
5. Land Use, Land-Use Change and Forestry	-7 358.06	-6 911.26	-5 817.96	-5 588.77	-4 703.66	-5 065.65	-6 166.52	-6 936.73
6. Waste	1 482.95	1 418.32	1 453.02	1 496.93	1 545.35	1 595.55	1 619.65	1 638.14

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	2014	2015	2016	2017	2018	2019	2020	2021
	<i>Gg of CO<sub>2</sub> equivalents</i>							
1. Energy	27 076.55	27 781.52	28 034.09	28 987.41	28 826.42	27 317.54	25 107.26	27 836.04
2. Industrial Processes	8 503.52	8 690.97	8 889.20	9 175.89	9 202.30	8 358.27	7 807.24	9 226.34
4. Agriculture	2 429.97	2 191.16	2 357.62	2 226.09	2 194.58	2 227.29	2 237.62	2 085.21
5. Land Use, Land-Use Change and Forestry	-4 755.17	-5 234.66	-5 313.25	-5 204.83	-4 232.06	-4 995.18	-7 179.80	-7 212.82
6. Waste	1 628.92	1 735.61	1 673.14	1 709.64	1 728.43	1 711.52	1 759.76	1 729.32

*Total aggregated GHG emissions, emissions are determined as of March 2025, indirect emissions are reported in the 2025 submission.*

## ES.4. Indirect Emissions and Precursors of Greenhouse Gases

The Slovak Republic is providing here the estimate of CO, NO<sub>x</sub>, SO<sub>2</sub> and NMVOC emissions for the years 1990 – 2023 originally submitted under the NECD and the CLRTAP on February 15, 2025. The estimation for the year 2023 was available in preliminary format of the NECD submission. The latest data is now included in CRT tables 1990 – 2023 generated using the ETF software, accompanied by the json file. According to the new rules for the reporting of the air pollutants recalling the Article 8(1) and the Annex I of the NECD, annual emission reporting requirements as referred to in the first subparagraph of the Article 8(1) for the years after the year 2017 was set for the emissions inventory in February, 15 and for the informative inventory reports (IIR) or emissions data resubmission in March, 15, respectively.

Major changes in comparison with the 2024 submission are among others for example:

- In the Energy sector, emissions of NO<sub>x</sub>, NMVOC, SO<sub>x</sub> and CO in the categories 1.A.3.b and 1.A.5.a changed based on update of the methodology and due to update of the COPERT model.
- In the IPPU sector, emissions of NMVOC in the category 2.H.2, changed as a result of the activity data update. Emission of SO<sub>x</sub> were recalculated in category 2.D.3.i due to update of the COPERT model.
- In the Agriculture sector, in categories 3.D.a.2.a and 3.D.a.2.c emissions of NO<sub>x</sub> changed as a result of correction of activity data. In category 3.B.3, emission of NO<sub>x</sub> changed based on the correction of Nitrogen excretion values. Emissions of NMVOC changed based on correction of calculation error in the distribution of swine population to the categories. In the category 3.D.e, NMVOC emission changed based on the EU recommendation.
- In the Waste sector, emissions of NO<sub>x</sub>, NMVOC, SO<sub>x</sub> and CO changed in the categories 5.C.1.b.i, 5.C.1.b.i.i and 5.C.1.b.v due to improvement to tier 2 methodology and implementation and reconsideration of abatement technologies into calculation according to the EMEP/EEA GB version 2023. Emission of NMVOC were also recalculated in the categories 5.D.1 and 5.D.2 due to reallocation of the emissions from biogas burning to the energy category in compliance with the GHG inventory. Emission of SO<sub>x</sub> were recalculated in category 5.C.1.b.i.i.i due to the reconsideration of abatement technologies used in the sources according to the EMEP/EEA GB version 2023.

These changes are resulted to the methodological changes in the NECD inventory and are reflected in the February 15, 2025 NECD submission and consequently provided in the GHG inventory submission 2025. According to the analyses, there are no larger inconsistencies (+/-5%) in the reporting under NECD (or CLRTAP) (submitted on 15/02/2025) and the GHG inventory (submitted on 15/03/2025). Due to differences in methodology, small inconsistencies occurred in the emissions from forest fires that are not included in the NECD inventory and emissions of NO<sub>x</sub> in manure management are not included directly in the GHG inventory (indirect N<sub>2</sub>O emissions are calculated based on NO<sub>x</sub> emissions in the category 3.B.2 – Manure Management). More information can be found in [Chapter 10.1](#).

**Table ES.4: Summary of the indirect GHG emissions according to the gases and the sectors in 2023**

EMISSIONS	TOTAL	ENERGY	INDUSTRY	AGRICULTURE	LULUCF	WASTE
	Gg					
NO <sub>x</sub>	51.193	38.619	5.838	6.038	0.345	0.353
CO	271.498	165.544	81.034	NO	12.311	12.610
NMVOC	76.257	41.764	26.458	7.340	0.274	0.421
SO <sub>2</sub>	13.617	8.684	4.893	NO	0.006	0.034

---

Emissions of main pollutants are available in public databases:

- [ŠÚ SR](#) in the STATdat database.
- [SHMÚ website](#) – Air Emission Accounts data for the years 2008 – 2023 are available as the aggregates in format of separate PDF files for particular gases.



---

# CHAPTER 1. INTRODUCTION

## 1.1. Background Information on Greenhouse Gas Inventories and Climate Change

### 1.1.1. Climate Change

The greenhouse effect of the atmosphere is similar to the effect that may be observed in greenhouses, however the function of glass in the atmosphere is taken over by the "greenhouse gases" (international abbreviation GHGs). Short wave solar radiation is transmitted freely through the greenhouse gases, falling to the earth's surface and heating it. Long wave (infrared) radiation, emitted by the earth's surface, is caught by these gases in the major way and partly reemitted towards the earth's surface. Because of this effect, the average temperature of the surface atmosphere is 33°C warmer than it would be without the greenhouse gases. Finally, this enables the life on our planet.

The most important greenhouse gas in the atmosphere is water vapour (H<sub>2</sub>O), which is responsible for approximately two thirds of the total greenhouse effect. Its content in the atmosphere is not directly affected by human activity, in principle it is determined by the natural water cycle, expressed in a very simple way, as the difference between evaporation and precipitation. Carbon dioxide (CO<sub>2</sub>) contributes to the greenhouse effect by 30%, methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and ozone (O<sub>3</sub>); all three together contribute by 3%. The group of synthetic (artificial) substances – chlorofluorocarbons (CFCs), their substitutes, hydrofluorocarbons (HFCs, HFCS) and others such as fluorocarbons (PFCs) and SF<sub>6</sub>, also belong to the greenhouse gases. There are other photochemical active gases as well, such as carbon monoxide (CO), oxides of nitrogen (NO<sub>x</sub>) and non-methane organic compounds (NM VOC), which do not belong to the greenhouse gases, but contribute indirectly to the greenhouse effect of the atmosphere. They are registered together as the precursors of ozone in the atmosphere, as they influence the formation and disintegration of ozone in the atmosphere.

According to the global climatologic classifications, the Slovak Republic is located in the mild climate zone with mean monthly precipitation totals equally distributed over the whole year. The Atlantic Ocean affects more the western part of the country and the continental influence is more typical for the eastern part. The Mediterranean climate influences mainly the south of the central part of Slovakia by higher precipitation totals in autumn. A regular rotation of spring, summer, autumn and winter seasons is typical for the country. However, the overall increase of GHGs emissions concentration caused significant climatic changes in the temperature, water regime and extreme weather events in Slovakia.

Detail climatic measurements at several meteorological stations and more than 200 precipitation gauges since 1881 has enabled us to prepare the study on climate change and variability for the period of 1881 – 2023. It is also possible to separate natural causes of climate changes from those induced by enhanced atmospheric greenhouse effect (using global and regional climatic analyses).

Slovakia has seen a significant increase in above-normal temperature years since 1991, with 2018 and 2019 being extremely warm. In the period 2001 – 2022, dry, rainfall-free periods have been shown to occur more frequently, which, combined with warmer average climatic conditions, leads to more frequent and more widespread soil drought. A major problem in Central Europe and Slovakia is the significant change in the temporal and spatial distribution of precipitation and snow cover. Precipitation in the warm part of the year occurs more often in the form of intense torrential downpours and in the cold part of the year more often in liquid form. The climate change scenarios described in this report assume comparable increases in monthly and annual temperatures of 1.5 to 4.7°C in Slovakia. While the temperature scenarios are very similar in all Slovak locations, the precipitation scenarios show some regional differences. Higher increases in annual precipitation totals are expected in the north of the

country, with summer decreases in precipitation more significant in the southern lowlands. A comparable increase (decrease) is also projected for daily maximum precipitation totals.

Information on climate change and adaptation measures in Slovakia can be found in [the First Biennial Transparency Report of Slovakia to the UNFCCC](#) published in December 2024.

### 1.1.2. Greenhouse Gas Inventories

This National Inventory Document (NID) of Slovakia for the submission to the EU, the UNFCCC and to the Paris Agreement includes data of the anthropogenic emissions by sources and removals by sinks of all greenhouse gases (GHGs) not controlled by the Montreal Protocol, i.e. carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), perfluorocarbons (PFCs), hydrofluorocarbons (HFCs), and sulphur hexafluoride (SF<sub>6</sub>). Emissions of nitrogen trifluoride (NF<sub>3</sub>) did not occur in Slovakia and appropriate notation key was used in inventory. These gases contribute directly to climate change owing to their positive radiative forcing effect. The Slovak GHG inventory includes also the three indirect greenhouse gases (nitrogen oxides reported as NO<sub>2</sub>, carbon monoxide reported as CO, non-methane volatile compounds reported as NMVOC) and sulphur oxides reported as SO<sub>2</sub> ([Chapter ES.5](#)).

In addition, the indirect CO<sub>2</sub> and N<sub>2</sub>O emissions resulting from atmospheric oxidation of NH<sub>3</sub>, CH<sub>4</sub> and NMVOC emissions from non-biogenic sources are also included in the inventory in the sectoral tables (IPPU and Agriculture). The indirect CO<sub>2</sub> emissions have been evaluated and included in the IPPU sector. Indirect N<sub>2</sub>O emissions resulting from a deposition of nitrogen due to emissions of nitrogen oxides (NO<sub>x</sub>) and ammonia (NH<sub>3</sub>) are estimated and indirect N<sub>2</sub>O emissions from agricultural sources are included in the national total emissions.

GHG Emissions inventory of Slovakia is fully consistent with the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (the IPCC 2006 GL) and according to the 2019 IPCC Refinements since the base year and reporting arrangement is consistent with the Modalities, Procedurals Guidelines according to the [Decision 24/CP.19. Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention](#).

According to the emissions inventory submitted in March 2025, the Slovak Republic total anthropogenic emissions of greenhouse gases expressed as CO<sub>2</sub> equivalent decreased by more than 51% without LULUCF and with indirect emissions, compared to the base year 1990. This achievement is the result of impacts of several processes and factors, mainly:

- Recovery and investments after the Covid-19 pandemic impacts on transport, industry and services and do not followed the increasing trend of GDP (further decoupling) after 2021.
- Higher share of services on the GDP.
- Technological and fuels restructuring and change in structure of industries pushed by increasing of fossil fuels' and electricity prices.
- Higher share of gaseous fuels and biofuels (biomass) on consumption of primary energy resources.
- Gradual decrease in energy consumption for certain energy intensive sectors (increase of energy efficiency).
- Impact of air protection legislation, which regulates directly or indirectly generation of greenhouse gas emissions.
- Increase share of the renewable energy sources on final consumption.
- In 2023, further decrease in emissions of EU ETS occurred due to high fuel prices, several operators phased-out or reduced production. This, along with other factors, caused the changes in the share of allocated emissions in the EU ETS and the ESD/ESR; in the EU ETS (47%) and the ESD/ESR (53%) ([Table 1.1](#)).

- Implementation of strict policies and measures in climate change and international agreements up to 2030/2040 focused mostly on the EU ETS categories.
- Less intensive winter seasons, lower fuel consumption for heating in 2023/24.
- Higher share of biomass in the residential heating sector.

In May 2004, the Slovak Republic joined the European Union. Relevant European legislation has brought additional positive direct and indirect effects to the reduction of GHG emissions, mainly in the Energy sector. The introduction of emission trading system will allow the implementation of further reduction measures in all installations included in the EU ETS.

**Table 1.1: Total GHG emissions in the EU ETS and ESD/ESR for the years 2019 – 2023**


YEAR	2023	2022	2021	2020	2019
	Gg of CO <sub>2</sub> equivalents				
<b>Total greenhouse gas emissions without LULUCF and with indirect emissions</b>	36 114.74	36 880.44	40 920.57	36 957.76	39 659.93
<b>Total verified EU ETS emissions</b>	16 994.18	17 418.25	20 898.87	18 170.00	19 903.84
<b>CO<sub>2</sub> emissions from 1.A.3.A civil aviation</b>	1.56	1.48	1.29	0.88	1.83
<b>Total verified ESD/ESR emissions</b>	19 119.00	19 460.71	20 020.41	18 786.88	19 754.26

**Table 1.2** and **Figure 1.1** show the most significant trend indicator of GDP and GHG emissions decoupling which was achieved in Slovakia in past years. In addition, development in the last inventory year (2023) is an evidence of continuation of decoupling process started in the 1997 and continuing after COVID-19 and fuel prices increasing (due war in Ukraine). This is a signal of total reconstruction of Slovak economy and industry. It is also expected, that similar trend will continue in the future, while there are planned investments in energy saving and efficiency of building, electromobility, alternative fuels policy and step by step building a carbon neutral economy.

**Table 1.2: Decrease of carbon intensity per GDP in the Slovak Republic in 2008 – 2023**

<b>YEAR</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>
<b>CO<sub>2</sub> emission in Tg</b>	41.43	37.68	38.46	38.05	35.96	35.62	33.71	34.53
<b>GDP in Bio € at ESA 2015 prices</b>	75.85	71.68	76.54	78.51	79.74	80.30	82.47	86.74
<b>Carbon Intensity in Tg/GDP</b>	0.55	0.53	0.50	0.48	0.45	0.44	0.41	0.40
<b>YEAR</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>	<b>2022</b>	<b>2023</b>
<b>CO<sub>2</sub> emission in Tg</b>	34.98	36.18	36.18	33.85	31.18	35.25	31.59	30.77
<b>GDP in Bio € at ESA 2015 prices</b>	88.43	90.97	94.67	96.82	94.32	99.72	100.17	101.55
<b>Carbon Intensity in Tg/GDP</b>	0.40	0.40	0.38	0.35	0.33	0.35	0.32	0.30

**Figure 1.1: Comparison of CO<sub>2</sub> emissions per GDP (carbon intensity) in 1995 – 2023**



The Slovak Statistical Office, Dpt. of National Accounts. Within the revision of annual national accounts (base year 2015).

### 1.1.3. International Agreements

#### International agreements under the UN:

United Nations Framework Convention on Climate Change (UNFCCC):

- Adopted on May 9, 1992 in New York
- Adopted by the Slovak Republic on May 19, 1993
- Ratified by the Slovak Republic on August 25, 1994
- Entry into force for the Slovak Republic on November 23, 1994

The aim of the Convention is to stabilize the atmospheric concentrations of greenhouse gases to a safe level that enables adapting of ecosystems and to prevent the dangerous consequences of the impact of anthropogenic activity.

Kyoto Protocol (KP):

- Adopted on December 11 1997 in Kyoto, Japan
- Adopted by the Slovak Republic on February 26, 1999
- Entered into force for the Slovak Republic on February 16, 2005
- Amendment to KP adopted on December 8, 2012 in Doha, Qatar

Slovakia fulfilled its commitments for both first and second commitment period with emissions significantly lower than the reduction targets. The obligation to provide the supplementary information report with our GHG inventory as required under article 7 paragraph 1 of the Kyoto Protocol is no longer applicable. Indeed, the end of the compliance review following the True-up period report of the second commitment period of the Kyoto protocol marked the end of the associated reporting obligations. True-up period report and review report for the second commitment period of the Kyoto Protocol for Slovakia can be find here <https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review/reporting-and-review-under-the-kyoto-protocol/second-commitment-period/reporting-and-review-process-for-the-true-up-period-of-the-second-commitment-period-of-the-kyoto>.

Paris Agreement (PA):

- Adopted on 12 December 2015 in Paris
- Adopted by the Slovak Republic on April 22, 2016
- Ratified by the Slovak Republic on September 28, 2016
- Entered into force for the Slovak Republic on November 4, 2016

The Paris Agreement central aim is to strengthen the global response to the threat of climate change by keeping a global temperature rise this century well below 2°C above pre-industrial levels and to pursue

---

efforts to limit the temperature increase even further to 1.5°C. Monitoring, reporting and reducing emissions, including adaptation to climate change, is mandatory for all countries, not just those listed in Annex 1 to the Convention. Emissions reduction action plans, defined as nationally determined contributions (NDCs), set targets for reducing greenhouse gas emissions by 2025 or 2030, along with adaptation to the climate change. Countries should review and tighten their NDCs every 5 years to achieve carbon neutrality by 2050. Slovakia submitted in December 2024, the First Biennial Report under the Paris Agreement and fulfil the PA requirement.

#### International agreements under the EU:

The European Union (EU) considers climate change as one of the four environmental priorities. On November 28, 2018, the European Commission presented its Long-Term Strategy for a prosperous, modern, competitive and climate-neutral economy by 2050. The Low-Carbon Development Strategy of the Slovak Republic until 2030 with a View to 2050 was adopted by the Government of the Slovak Republic by the Resolution No 104/2020. The European Commission launched the European Climate Pact in December 2020, an EU-wide initiative inviting people, communities and organisations to participate in climate action and build a greener Europe. As part of the European Green Deal, the Climate Pact offers a space for everyone to share information, debate and act on the climate crisis, and to be part of an ever-growing European climate movement. The Commission's proposal to cut greenhouse gas emissions by at least 55% by 2030 and 90% by 2040 sets Europe on a responsible path to become climate neutral by 2050.

The Regulation (EU) 2018/1999 of the European Parliament and of the Council of 11 December 2018 on the Governance of the Energy Union and Climate Action (EnGov):

The Regulation (EU) 2018/1999 together with Commission implementing Regulation (EU) 2020/1208 on structure, format, submission processes and review of information reported by Member States pursuant to Regulation (EU) 2018/1999 integrated rules to ensure planning, monitoring and reporting of progress towards its 2030 climate and energy targets and its international commitments under the Paris Agreement have been adopted. The Regulation established a governance mechanism for the implementation of strategies and measures designed to meet the objectives and targets of the Energy Union and the EU's long-term greenhouse gas emission commitments under the Paris Agreement, in particular the EU's ambition to achieve climate neutrality by 2030. Slovakia submitted the 2021 – 2030 National Energy and Climate Plan under the Regulation on the Governance by the end of 2019 and by 2024 updated NECP in line with article 14 of the Governance Regulation.

The Low-Carbon Development Strategy of the Slovak Republic until 2030 with a View to 2050 (LCDS), adopted in March 2020, aims to identify measures, including additional measures, to achieve climate neutrality in the Slovak Republic by 2050. The aim of the LCDS is to outline options for a comprehensive long-term (30-year) strategic roadmap for moving to a low-carbon economy, which will be completed by achieving climate neutrality by 2050. The LCDS identifies key policies and measures that will lead to achieving the headline target of the Paris Agreement – keeping the increase in global temperature this century to well below 2°C and pursuing efforts to limit the temperature increase to 1.5°C above pre-industrial levels. The LCDS aims to select and analyse cost-effective measures in terms of the scope of emission reductions and the economic and social impact. The measures envisaged in the near future, detailed, and modelled in the strategy under the WEM and WAM scenarios raised the fact that climate neutrality in Slovakia cannot be achieved by 2050 with them. Therefore, the strategy also includes additional measures (called NEUTRAL) which should move Slovakia closer to its goal by 2050. Whether this happens will be analysed in detail in the near future as part of the updating process. The implementation of the measures will require the active involvement of the relevant sectors, the interconnection and consolidation of the individual sectoral and crosscutting policies, and society-wide engagement. Consistent horizontal implementation of measures that are in harmony with the objective of achieving climate neutrality by the middle of this century and in line with this strategy is to be ensured

---

by the Council of the Government of the Slovak Republic for the European Green Deal, the adoption of which is expected together with this Strategy.

Consistent horizontal implementation of measures in line with the objectives of climate neutrality by 2050 and in line with the LCDS is to be ensured by the Council of the Government of the Slovak Republic for the European Green Deal and Low-Carbon Transformation, adopted by the Government Resolution No 699 of November 4, 2020.

Thanks to the new approved environmental policy Greener Slovakia – Strategy of the Environmental Policy of the Slovak Republic until 2030 (the Envirostrategy 2030), Slovakia determined a way of how to face the biggest environmental challenges and address the most serious environmental problems. The Slovak Government approved the Envirostrategy 2030 on February 27, 2019.

## 1.2. Description of the National Inventory Arrangements

### 1.2.1. Institutional, Legal and Procedurals Arrangements

The Ministry of Environment of the Slovak Republic (MŽP SR) is responsible for development and implementation of national environmental policy including climate change and air protection objectives. It has the responsibility to develop strategies and further instruments of implementation, such as acts, regulatory measures, economic and market based instruments for cost efficient fulfilment of adopted goals. All ministries and other relevant bodies annotate both, the conceptual documents as well as legislative proposals. Following the commenting process, the proposed acts are negotiated in the Legislative Council of the Government, approved by the Government, and finally by the Slovak Parliament. The Ministry of Environment of the Slovak Republic is the main body to ensure conditions fulfilment and to monitor progress of the Slovak Republic for meeting all commitments and obligations of climate change and adaptation policy.

According to the Governmental Resolution No 3/2025 Coll. from 15<sup>th</sup> January 2025, Government established the inter-ministerial Council of the Slovak Republic for Sustainable Development and its Financing (CSDF). This Council replaced previous Committee for Agenda 2030 on Sustainable Development, Committee for European Green Deal and Governmental Committee for Recovery and Resilience Plan. New Council is chaired by the deputy prime minister for Recovery and Resilience Plan of the Slovak Republic; other members were minister of the Environment of the Slovak Republic, minister of Investments, Regional Development and Informatization of the Slovak Republic, minister of Finance of the Slovak Republic and minister of Foreign Affairs of the Slovak Republic. Other members of the Council are ministers, representatives of professional industry associations, and unions from the private and public sectors. [Figure 1.2](#) provides in depth overview diagram showing the institutional arrangements concerning climate policy and its implementation.


The creation of the Council increases the coherence and efficiency of decision-making at the government level in such important areas as sustainable development, the European Union Green Deal and the Recovery and Resilience Plan. At the same time, there will be a closer link between the Sustainable Development Goals and the spending of public resources. The establishment of the Council also establishes the Executive Board for the Recovery and Resilience Plan of the Slovak Republic, which performs the following tasks:

- monitors progress in achieving milestones and targets within all components of the Recovery Plan,
- performs an advisory role in relation to the strategy for implementing the Recovery Plan,
- at the proposal of the National Implementation and Coordination Authority, expresses its opinion on the implementation of reforms and investments in accordance with the Recovery Plan,

- adopts positions in the form of resolutions, proposes recommendations and draws attention to risks in the implementation of individual investments and reforms of the Recovery Plan,
- if necessary, also deals with issues of synergies and complementarities between the Recovery and Resilience Mechanism (hereinafter referred to as the “Mechanism”) and European Union funds, as well as between the Mechanism and other European Union support instruments.

The members of this Executive Board for the Recovery and Resilience Plan of the Slovak Republic are representatives of the implementers of the Recovery Plan within the Slovak Republic. In addition to establishing working groups, the Council will also establish an expert advisory committee, which will be composed of prominent experts in the areas of individual pillars of sustainable development with international experience and professional practice acquired in the relevant area. Further information - available only in Slovak: <https://rokovania.gov.sk/RVL/Material/30347/1>.

**Figure 1.2: Institutional arrangements in climate change policy and its implementation**



On the EU level, according to the Regulation on the Governance of the Energy Union and Climate Action by 15 March 2021, and every two years thereafter, Member States shall report to the Commission information on their national climate change adaptation planning and strategies, outlining their

---

implemented and planned actions to facilitate adaptation to climate change, including the information specified in Part 1 of Annex VIII and in accordance with the reporting requirements agreed upon under the UNFCCC and the Paris Agreement.

#### National System for GHG Emissions Inventories of the Slovak Republic (NS SR)

Articles 4 and 12 of the UNFCCC require the Parties to the UNFCCC to develop, periodically update, publish, and make available to the Conference of the Parties their national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled under the Montreal Protocol. Moreover, the commitments require estimation of emissions and removals as a part of ensure that Parties are in compliance with emission limits, that they have a national system for estimation of sources and sinks of greenhouse gases, that they submit an inventory annually, and that they formulate national programs to improve the quality of emission factors, activity data, or methods. The obligation of the Slovak Republic to create and maintain the National System of the Slovak Republic (NS SR) which enables continual monitoring of greenhouse gases emissions is given in the Decision 18/CMA.1 Modalities, Procedures and Guidelines (MPG) for the transparency framework for action and support referred to in Article 13 of the Paris Agreement.

Setting up the NS SR of emissions in compliance with the MPG requirements was framed with functions which it should fulfil according to the decision 18/CMA.1. The basic characteristics of the NS SR are as follows:

- To ensure linkages and co-operation among involved institutions, bodies and individuals to perform all activities for monitoring and estimation of GHG emissions from all sectors/categories according to the UNFCCC guidelines and relevant decisions and according to the approved IPCC methodologies. To enable using of all relevant data from national and international databases for preparing and improving GHG emission inventory.
- To define role and competencies of all involved stakeholders including the role of National Focal Point to the UNFCCC.
- To define and regularly implement quality assurance and quality control (QA/QC) process in two lines; both internally and externally by appropriate body.
- To ensure ongoing process of development capacities; financial, technical and expert sources in relation to QA/QC but also in relation to new tasks rising from the international process.

The National System of the Slovak Republic was established and officially announced by Decision of the Ministry of Environment of the Slovak Republic on 1<sup>st</sup> January 2007 in the official bulletin: Vestník, Ministry of Environment, XV, 3, 2007. In agreement with paragraph 30(f) of Annex to Decision 19/CMP.1, which gives the definitions of all qualitative parameters for the NS SR, the description of quality assurance and quality control plan according to Article 5, paragraph 1 is also required. The revised report of the NS SR dated on November 2008 was focused on the changes in the institutional arrangement, quality assurance/quality control plan and planned improvements. The regular update of the NS SR with all qualitative and quantitative indicators is provided on website.

The latest update was published in accordance with the publication of the first Biennial Transparency Report under the Paris Agreement, submitted in December 2024.

#### The role of responsible ministries in the national system

The MŽP SR is responsible for implementation of national environmental policy including climate change and air protection. It serves also as the National Focal Point to the UNFCCC.

District and regional environmental offices are decision-making bodies according to Act No 525/2003 Coll. These are located at eight regional and 46 district administration offices. The four inspectorates of the Slovak Environmental Inspection carry out inspection and enforcement activities. According to the



---

Act No 146/2023 Coll. on Air Protection, competencies and decision-making process concerning large, medium and small pollution sources are given to regional and district levels and municipalities.

Act No 414/2012 Coll. on Emission Trading as amended is the legal instrument directly oriented towards the control of GHG emissions. According to this Act, competencies with respect to emission allowance trading are given to the MŽP SR and the regional and district environmental offices.

#### [Slovak Hydrometeorological Institute as the single national entity](#)

The Slovak Hydrometeorological Institute (SHMÚ) [www.shmu.sk](http://www.shmu.sk) is authorised by the MŽP SR to provide environmental services, including annual GHG inventories according to the [approved statute](#). The range of services, competencies, time schedule and financial budget are updated and agreed annually. All details of the SHMÚ activities are described in the Plan of Main Tasks. The plan, commented by all stakeholders, is published after approval at the [website of the SHMÚ](#). Deadline for the approval of this plan by the ministry is 31<sup>st</sup> December each year. In 2024, organisational changes occurred and the [structure of SHMÚ](#) was updated. Presented changes have no impact on the [NS SR](#) (update in 2024). Establishment of the Department of Emissions and Biofuels (OEaB) was based on organisational changes provided in January 2017. The OEaB has two main tasks: emission inventories and projections (GHG, NECD, and CRLTAP) and National System of Biofuels. The OEaB is also responsible for developing and maintaining the National Emission Information System (the [NEIS](#)) – the database of stationary sources to monitor the development of SO<sub>2</sub>, NO<sub>x</sub>, CO emissions at regional level and to fulfil reporting commitments under the national regulations and EU Directives. The NEIS software product is constructed as a multi-module system, corresponding fully to the requirements of current legislation. The NEIS database contains also some technical information about the sources like fuel consumption and use for the estimation of the sectoral approach. The Single National Entity is a part of the OEaB with the defined structure and overall responsibility for compilation and finalization of the inventory reports and their submission to the UNFCCC Secretariat and the European Commission according to the announcement. The SNE was officially appointed by the Decision of the Director General of the SHMÚ No 16/2011 in August 2011 and amended by the Decision of the Director General of the SHMÚ No 8/2012 in September 2012. The SNE coordinates the NS SR.

On [Figure 1.3](#) is visible a structure of the NS SR, where the Council CSDF is an intergovernmental body responsible for implementation of climate change policy on cross-ministerial level and composition of the NS SR with updated list of internal and external experts is presented in the [Table 1.3](#)

#### [Responsibilities of expert organisations](#)


Contracts with the external institutions and the sectoral experts are fully in a competence of the SNE after previous approval by the MŽP SR. The SNE is fulfilling inventory tasks fully in line with approved annual the Plan of Main Tasks and with financial resources allocated by the MŽP SR. To specify main objectives for given year, kick-off workshop with participation of the MŽP SR, SHMÚ and external co-operating bodies and experts is organised regularly, usually at the beginning of February each year. This workshop is also an official forum for closing and summing up outcomes from the previous year and preparing the activities, including the QA/QC plan and responsibilities for the current year.

The main institutions involved in the compilation of the GHG inventory are:

- Ministry of Environment of the Slovak Republic;
- Slovak Hydrometeorological Institute;
- Statistical Office of the Slovak Republic;
- Slovak Technical University, Faculty of Chemical Engineering
- National Forest Centre – Ministry of Agriculture and Rural Development;
- Research Institute on Soil Protection Bratislava - Ministry of Agriculture and Rural Development.

Supporting institutions, founded by the Ministry of Environment to perform specific tasks linked to inventory activities, play an important role. These include the Slovak Hydrometeorological Institute, the Water Research Institute and the Slovak Environmental Agency. There are also other relevant subjects for data providing, which are listed in sectoral chapters (*Table 1.3*).

**Figure 1.3:** Structure and responsibilities of the NS SR



**Table 1.3:** List of the sectoral experts in the NS SR

INTERNAL EXPERTS - SHMÚ		
INSTITUTION	NAME	RESPONSIBILITY
Dept. of Emissions and Biofuels	Ms. Janka Szemesová	NS SR coordinator
Dept. of Emissions and Biofuels	Ms. Alexandra Nadžadyová	Data manager of quality Biofuels expert
Dept. of Emissions and Biofuels	Mr. Ján Horváth	Deputy of NS SR coordinator Energy expert
Dept. of Emissions and Biofuels	Mr. Marcel Zemko Mr. Jozef Orečný Mr. Roman Mach	Emission projections experts Buildings sector emissions
Dept. of Emissions and Biofuels	Ms. Michaela Câmpian Ms. Petra Kršáková	Other pollutant experts
Dept. of Emissions and Biofuels	Ms. Patrícia Navrátilová	Agricultural expert
Dept. of Emissions and Biofuels	Ms. Monika Jalšovská	NEIS expert
Dept. of Emissions and Biofuels	Mr. Roman Mach	Data manager Uncertainty analyses QA activity
Dept. of Water Quality	Ms. Lea Mrafková	GHG inventory in wastewater sector
Dept. of Climate Service	Mr. Peter Kajaba	Climatological data

EXTERNAL INSTITUTIONS/EXPERTS		
INSTITUTION	NAME	RESPONSIBILITY
Astraia	Mr. Ján Judák	Reference approach
National Forest Centre Zvolen	Mr. Ivan Barka Mr. Tibor Priwitzer Mr. Pavel Pavlenda	GHG inventory in Forest Land and LULUCF coordinators
Animal Production Research Centre	Ms. Zuzana Palkovičová Mr. Ondrej Pastierik Mr. Miroslav Záhradník	GHG inventory in agriculture – animal production
Research Institute on Soil Protection Bratislava National Agricultural and Food Institute	Mr. Michal Sviček Mr. Pavol Bezák Ms. Kristína Buchová	Data provider in agriculture sector – soils, LULUCF Cropland and fertilizers
Central Control and Testing Institute in Agriculture	Mr. Štefan Gáborík Ms. Maggioni-Brázová Ildikó	Data provider in the Agricultural sector – soil nutrition
Faculty of Chemical Technology of the Slovak Technical University Bratislava	Mr. Vladimír Danielik Mr. Juraj Labovský	GHG inventory in industrial processes and solvent use sectors and energy – sectoral approach Consultation in fuel balance Consultation for EU ETS
Faculty of Chemical Technology of the Slovak Technical University Bratislava	Mr. Igor Bodík	GHG inventory in waste – wastewater
Independent Expert	Mr. Marek Hrabčák	GHG inventory in waste – SWDS
Statistical Office of the Slovak Republic – Department of Cross-sectoral Statistics	Ms. Maria Lexová	Statistical energy data provider
Slovak Association for Cooling and Air Conditioning Technology Ministry of Environment		F-gases data provider
Grassland and Mountain Agriculture Research Institute	Mr. Štefan Pollák	GHG inventory in Grassland

### 1.2.2. Inventory Planning, Preparation and Management

The preparation of emission inventories within the NS SR for GHG emissions is decentralized according to the definition of Article 13 of the PA. The individual sectors are fully under the responsibilities of the external institutions and sectoral experts, who are authorized to evaluate the emissions inventory within the delegated sectors. The preparation of the inventory includes three stages – inventory planning, preparation and management.

During the inventory planning are set up roles and responsibilities, specifying processes and resources according to internal and external QA/QC plans. These plans are updated and evaluated annually by the quality manager of the NS SR and approved by the MŽP SR. The inventory preparation process starts with the collection of activity data, emission factors and all relevant information needed for estimation of emissions, followed by choice of methods, data processing and then archiving.

For the inventory management, reliable data management to fulfil the data collecting and reporting requirements is necessary. The inventory management includes a control system for documents and data and for their archives.

### 1.2.3. Quality Assurance/Quality Control and Plans

This section presents the quality management and inventory process. Category – specific QA/QC details with improvements and recommendations are discussed in the relevant sectoral chapters of this NID.

---

## Quality management

The Slovak Hydrometeorological Institute has built and introduced the quality management system (QMS) according to the requirements of EN ISO 9001:2008 standard of conformity. In the frame of introduction of the QMS for the SHMÚ as a global standard, the certification itself proceeds according to the partial processes inside of the SHMÚ structure. The process of Emission Inventories was the subject of internal and external audits during the March 2010 by the certification body ACERT, accredited by Slovak National Accreditation Service. The quality manager has completed several trainings regarding the QMS. The recertification process is taking place every two years.

The objective of the NS SR is to produce high-quality GHG inventories. In the context of GHG inventories, a high quality provides, that both the structures of the national system (i.e. all institutional, legal and procedural arrangements) for GHG emissions and removals and the inventory submissions (i.e. outputs, products) comply with the requirements, principles and elements arising from the UNFCCC in line with the MRV principles. The IPCC Guidelines for the GHG emissions inventory 2006 were fully implemented. The IPCC Guidelines Refinements 2019 were considered for possible utilisation in inventory where the methodology was missing in previous Guidelines. The starting point for accomplishing a high-quality GHG inventory is consideration of the expectations and requirements directed at the inventory. The quality requirements set for the annual inventories – transparency, consistency, comparability, completeness, accuracy, timeliness and continuous improvement – are fulfilled by implementing the QA/QC process consistently. *Figure 1.4* shows a model for the timeline steps provided in inventory process, QA/QC and verification procedures.

The SHMÚ implemented a policy of continuous training process for internal and external experts. Experts are trained during workshops of the NS SR, which are held two times per year. The minutes of the workshop and all relevant documents are available to the sectoral experts of the NS SR. The latest meeting was held on June 27-28, 2024<sup>2</sup> and the other ways of communication within the NS SR are via e-mail, phone call, visits and meetings.

The sectoral experts apply the QA/QC methodology according to the Quality Manual, collect data from providers and process emission inventory for a given sector – they provide partial reports with information on quality and reliability of data on activities and emissions. The quality manual including e.g. guidelines, QA/QC plans, templates and checklists is available to all experts of the NS SR via the Internet. The set of templates and checklists consists these documents:

- ✓ QA/QC Plan (external, internal)
- ✓ Matrix of Responsibility
- ✓ General QC
- ✓ Source Category-specific QC
- ✓ Quality Assurance
- ✓ Archive Document
- ✓ Improvement plan
- ✓ Recommendation list

All documents after filling out by experts are approved by responsible person of inventory system and then are archived. The data manager has the overall responsibility for documentation, formal contact with the sector experts and approval activities, taking over the sectoral reports and archiving them.

---

<sup>2</sup> In the framework of the project EMISIE for the implementation of the IPCC 2019 Refinements


### Inventory planning (PLAN)

The inventory planning stage includes the setting of quality objectives and elaboration of the QA/QC plans for the coming inventory preparation, compilation and reporting work. The setting of quality objectives is based on the inventory principles. The quality objectives regarding all calculation sectors for inventory submissions are the following:

- ✓ Timeliness
- ✓ Completeness
- ✓ Consistency
- ✓ Comparability
- ✓ Accuracy
- ✓ Transparency
- ✓ Improvement

The quality objectives and the planned QC and QA activities regarding to all sectors are set in QA/QC plans (internal and external). In these documents, deadlines and responsibilities are described (included in **Annex 4** in **Tables A4.1** and **A4.2**). These plans updates and evaluates the quality manager of the NS SR and following are approving by the MŽP SR.

**Figure 1.4: PDCA cycle (Plan, Do, Check, Act)**



### Quality control procedures (DO)

The experts perform the general and category-specific QC procedures during inventory preparation, calculation and compilation.

General quality control includes routine checks of correctness, completeness of data, identification of errors, deficiencies and documentation and archiving of the inventory material. The sectoral experts must adopt adequate procedures for development and modification of the spreadsheets to minimise emission calculation errors. Checks ensure compliance with the established procedures as well as allow detecting the remaining errors. Parameters, emission units and conversion factors used for the calculations must be clearly singled out and specified.

Category-specific QC includes reviews of the source categories, activity data and emission factors focusing on key categories and on categories where significant methodological changes or data revision have taken place. Experts fill QC forms during the compilation of inventory; results from QC activities are documented and archived.

#### Quality assurance (CHECK)

Quality assurance is performed after application QC checks concerning the finalised inventory. QA procedures include reviews and audits to assess the quality of inventory and the inventory preparation and reporting process, determine the conformity of the procedures taken and to identify areas where improvements could be made. These procedures ongoing on different levels, including basic reviews of the draft reports, general public review, external peer review, internal audit, EU and UNFCCC reviews.

With uploading to the SHMÚ website, printing and distribution of the final inventory document feedback from public is appreciated. The sectoral experts and the members of inventory team are participating in various seminars, meetings, conferences and sector-specific workshops during the year. The activities of inventory members and results of national inventory of GHG emissions are reported there. A broader range of researchers and practitioners in non-government organizations, industry and academia, trade associations as well as the public have the opportunity to contribute to the final documents. Comments received during these processes are reviewed and, as appropriate, incorporated into the reports or reflected in the inventory estimates.

Independent experts from the MŽP SR and the sectoral experts not directly involved into inventory cycle (except of above-mentioned activities) now perform QA. Each sector has a different reviewer:

<b>GENERAL PART</b>	Ms. Miroslava Dančová Mr. Mário Gnida Ms. Lenka Zetochová	MŽP SR SHMÚ
<b>ENERGY</b>	Mr. Mário Gnida Mr. Marek Sokolík	MŽP SR Institute of Environmental Policy
<b>TRANSPORT</b>	Mr. Leoš Pelikán Ms. Zuzana Kačmárová	Centrum of Transport Research in Brno, Czech Republic <sup>3</sup>
<b>IPPU</b>	Mr. Jozef Škultéty	MŽP SR
<b>AFOLU</b>	Ms. Lenka Malatinská Ms. Hana Fratričová Ms. Kristína Buchová	MŽP SR MPaRV SR VÚPOP <sup>4</sup>
<b>WASTE</b>	Ms. Zuzana Jonáček Mr. Michal Patassi	SHMÚ MŽP SR

When checking the data quality of each sector, the NS SR coordinator, quality manager of the NS SR, data manager of the NS SR and other stakeholders must conduct the following general activities:

<sup>3</sup> In the framework of the Agreement on Mutual Cooperation signed in 2023

<sup>4</sup> Institute for Soil Protection

---

Checking: Check whether the data in the sectoral reports (calculations and documents) for each sector conform both to the general and specific procedures.

Documentation: Write down all verification results filling out a checklist, including conclusions and irregularities that have to be corrected. Such documentation helps to identify potential ways to improve the inventory as well as store evidence of the material that was checked and of the time when the check was performed.

Follow-up of corrective actions: All corrective actions necessary for documenting the activities carried out and the results achieved must be taken. If such check does not provide a clear clue concerning the steps to be taken, the quality control, bilateral discussion between expert and the NS SR coordinator will take place.

Data transference: All checked documents (including the final questionnaire and all annexes) shall be put into the project file and copies shall be forwarded to all the NS SR experts. Since the data quality supervision procedures must be observed all the time, it is not mandatory to conduct all checks annually during the inventory preparation. Certain activities, such as verification of the electronic data quality or project documentation for checking whether all documents have been provided, must be carried out every year or at least at set intervals. Some checks may be conducted only once (however, comprehensively) and then only from time to time.

Part of the QA procedures is bilateral cooperation with Czech Republic. The first meeting took place in July 2013 and since then is repeated every year. Team of GHG inventory experts from the SHMÚ and the Czech Hydrometeorological Institute (CHMI) met to exchange information and experience relating to the preparation of GHG inventory. In the last meeting, the experts from Slovakia, Czech Republic, Poland, Hungary and Austria attended. This last meeting with the Czech Hydrometeorological Institute (NIS CZ) took place in June 2024 in Bratislava (Slovakia) and the next meeting is scheduled for May 2025. Meeting includes expert discussion on sectoral level for sectors energy, IPPU, agriculture and waste. Separate meeting for LULUCF was held on national level on June 19, 2024.

In addition to the activities regarding the regional knowledge transferring in emissions inventories, the QA procedures focusing on introducing changes and improvements on national level are organised regularly. National experts, not directly involved in the NS SR, are invited to provide comments and discuss methodological issues.

### Verification activities

Independent verification procedure was introduced on the level of the Ministry of Environment (coordinator) with inclusion of other responsible experts from relevant ministries. The nominated experts and stakeholders involved in the verification and approval process for the selected parts of the emission inventory are responsible for the official and legislative agreement of the presented results and ensure harmonisation within several international reporting.

Verification refers to the collection of activities and procedures that can be followed during the planning and development, or after the completion of an inventory, that can help to establish its reliability for the intended applications of that inventory. The used parameters and factors, the consistency of data is checked regularly. Completeness checks are undertaken, new and previous estimates are compared every time. The sectoral expert for uncertainty checks data entry into the database many times. If possible, activity data from different data sources are compared and thus verified. Comprehensive consistency checks between national energy statistics and IEA time series. Checking the results of the EU's internal review for the EU-27 and analyse its relevance for Slovakia.

Confidential information is provided to the NS SR experts based on the bilateral agreements but cannot be reported on individual level (only aggregated) in emissions inventory.

### Inventory improvement (ACT)

The main aim of the QA/QC process is continuous improvement of the quality of inventory. The outcomes and experiences from the annual reviews are the main sources for the preparation of recommendation lists and improvement plans based on this recommendation lists.

The recommendation lists and improvement plans are updated annually after the regular UNFCCC and/or EU compliance reviews take place. As the Slovakia is one of the Member States of the European Union, the separate review regime is undertaken under the EU Effort Sharing Regulation (ESR) in spring every year. These outcomes and recommendations are included in the improvement plan, too. Detailed recommendation lists and improvement plans are prepared by the sectors and delivered to the sectoral experts for consideration and prioritisation of planned activities for the next inventory cycle. These plans are including in [Annex 4](#). According to the latest annual review on GHG emissions inventories 2022 (final ARR delivered on 4<sup>th</sup> April 2023), several ERT recommendations focused on general part of the inventory were implemented. These are connected with the key category analysis and uncertainty improvements.

Prioritisation process is based on recommendations raised during the previous UNFCCC and ESR reviews. Prioritisation for improvements is given to those categories of the GHG emissions inventory, where higher uncertainty is a result of the assessment. The latest examples can be found in categories of swine in agriculture or in 1.B.2 of fugitive methane emissions. The underlying assumptions used for estimating uncertainties applied on EF and AD are mostly based on the default values provided in the IPCC 2006 GL and/or expert judgment. The prioritisations are performing on annual basis also by quantitative assessment of uncertainty assessment (UA) for the base year and the latest inventory year. This approach is a part of the annual QA/QC system since 2017 submission. According to the previously identified outcomes made for tabular comparison of the key categories and tier method used, it was recognised, that the tier 1 approach (fugitive emissions of methane, direct soil emissions) was used several key categories. These categories are selected as the high priority of important to move to higher tier method. During the last years, the prioritisation of the Improvement Plan was focused on the Energy sector and the harmonisation of different data sources for energy balance and implementation of the higher tiers for fugitive emissions based on the IPCC 2019 Refinements. The methodological tiers for significant categories (bases on the UA results) are continuously improving, also for example in the Agricultural sector (change methodology from tier 1 up to tier 2 for enteric fermentation and manure management in swine and in direct soil emissions). In the Waste sector, the high priority in this inventory was put on distribution of the sewage sludge and implementation of the QA/QC activities. The improvement of the uncertainties in the LULUCF sector finished in 2022 and are fully implemented in 2024 submission and undated in 2025 ([Chapter 6](#)).

#### 1.2.4. Changes in the National Inventory Arrangements

During the preparation cycle of the GHG emissions inventory submitted in 2025, no significant changes in the arrangement or structure of the NS SR occurred. The NS SR is operational, functioning and fulfilling all main tasks and obligation in the line with the approved plans. NS SR is continuing in the process of strengthening capacity among the national system in line with the improvement and prioritization plans. The uncertainties calculations were previously based on external cooperation, now an internal expert is responsible for all sectors across inventory. In addition, a new expert was involved in the cropland category to strengthen new calculations on land-based matrix and new expert was involved into agricultural team. During previous years, the several new institutions were involved in the inventory, aiming to focus on QA activities, new internal (SHMÚ) expert on emission projections and emissions estimation in household sector.

In previous inventory submission (2024), outcomes of the [Project EMISSIONS](#) concerned to the implementation of the IPCC 2019 Refinement Guidelines was included in sectoral inventory. New methodological approaches were implemented on sectoral level in line with the recommendations from



the Refinement and considering national circumstance. A part of the project was also improving internal database and archiving system and ensuring higher consistency in dataflows. This was performed by the MESAP system (internal database IS) what is currently under testing and development at the NS SR.

### 1.2.5. Inventory Preparation, and Data Collection, Processing and Storage and Archiving

The compilation of the emission inventory starts with the collection of activity data. A comprehensive description of the inventory preparation for GHG emissions is described in methodologies for the individual sectors. The methodologies are updated annually within the improvement plan and recommendation list and they are archived after formal approval at the [web page](#) of the NS SR and by the sectoral experts and the NS SR coordinator. The most important source of activity data is the Statistical Office of the Slovak Republic and the sectoral statistics of the relevant ministries. The NEIS database is also important reference source of data on fuels and other characteristics of stationary air pollution sources. The OEaB of the SHMÚ operates the NEIS. Other important sources are listed in [Table 1.5](#) below and full catalogue of activity data is listed in the [NS SR description](#).

**Table 1.5:** List of important information sources for inventory preparation

SECTOR	SOURCE OF INPUT DATA
ENERGY	<a href="#">Energy Statistics of the SR</a> , <a href="#">NEIS</a> , <a href="#">www.spp.sk</a> , <a href="#">www.transpetrol.sk</a> , EU ETS Reports, Reports of the EU ETS verifiers
INDUSTRIAL PROCESSES AND PRODUCT USE	Association of cement and lime producers, Association of refrigeration and air conditioning engineers, Association of paper producers; EU ETS Reports, Reports of the EU ETS verifiers, Association for coating and adhesives, solvent distributors, <a href="#">Research Institute for Crude Oil</a>
AGRICULTURE	<a href="#">Green Report of the Ministry of Agriculture of the SR</a> - Agriculture, Institute for Fertilisers Research, <a href="#">List of Livestock to the 31. 12. 2023</a> , <a href="#">Crop yields data for crops and vegetables in 2023</a>
LULUCF	<a href="#">Green Report of the Ministry of Agriculture of the SR</a> - Forest, Cadastral Office
WASTE	Population (mid-year), Statistical Yearbook of Slovakia Table 3-3; Real Wage Index, Statistical Yearbook of Slovakia Table 1; Municipal Waste, industrial waste landfilled, Waste in the Slovak Republic in 2023; <a href="#">Database of disposal sites</a> ; Municipal Waste, industrial waste composted, industrial waste incinerated Waste in the Slovak Republic in 2019; Incinerators, <a href="#">Enviroportal</a> ; Generated, discharged BOD, COD, N, Environment in the SR (selected indicators in 2013 – 2023); Protein Consumption, Statistical Yearbook of Slovakia Table 5-8, State of Environment report 2023; Sludge, database of wastewater treatment plants, SHMÚ.

Collected input data are compared and checked with the international statistics (Eurostat, IAE, FAO and others). In some cases, the collected input data are compared with the results from models (e.g. in road transport it is COPERT model, model for the Waste sector, etc.).

Archiving of inventory documents and database is in the competence of the quality and data managers of the NS SR. Archiving of database is in the competence of the NS SR coordinator. Documents and emission inventories are archived at three levels. Official documents, methodologies and reports are archived and stored at the [web page](#) of the NS SR. The archiving is controlled by rules for archiving systems in organizations at the SHMÚ level. The documents needed for the quality management systems are archived in electronic form at the webpage of the SHMÚ (intranet). Documents required signature are printed and archived according to the archiving regulation of the Institute. Printed documents are archived in central archive of the SHMÚ and at the OEaB.

---

An archive system allows information to be easily reproduced, allows safeguards against data and information loss, and allows reproducibility of the estimates. The archive system includes relevant data sources and spreadsheets, reproduce the inventory and review all decisions about assumptions and methodologies. The archiving system checklist contains these archiving activities: documenting methods used, including those used to estimate uncertainty and data sources for each category; expert comments; revisions, changes in data inputs or methods and recalculation, also reason and source of changes; documenting the used software for calculation of emission. Each new inventory cycle benefits from effective data and documents management during development of the previous inventory.

Archived information includes all emission factors and activity data at the most detailed level, and documentation of how these factors and data have been generated and aggregated for the preparation of the inventory. This information also includes internal documentation on QA/QC procedures, external and internal reviews, documentation of annual key categories and key category identification, and planned inventory improvements and recommendations. All information on archiving is recorded in Archiving System. In addition, internal document about good practise in archiving were prepared. In this document, the exact way of archiving, procedures and steps is described.

### 1.2.6. Brief General Description of Methodologies and Data Sources Used

The methodologies used for the preparation of greenhouse gas inventory in the Slovak Republic are consistent with the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006 GL). In line with the Quality Improvement Plans of the NS SR, methodologies and parameters have been implemented fully in accordance with the 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories by the end of 2023 and therefore, this submission is fully in line with the IPCC 2019 Refinement. Detailed descriptions of used methodologies can be found as the sector specific ones in the following chapters of this Report. Regarding the tier approaches used in the NS SR, the detailed information can be found in CRT tables and sectoral chapters. The increasing tier of methodologies is one of the priorities mostly for key categories. This is also included in the improvement plan. In the view of provided recalculations, the higher tier method was implemented in the Agriculture, IPPU and Energy sectors.

Additional sources of activity data for the major sectors are as follows:

Energy:

The Statistical Office of the Slovak Republic:

- *Energ. P 2-01*: Yearly company statement on energy process of fuel enrichment.
- *Energ. P 3-01*: Yearly company statement on the consumption of fuels, electricity and heat for production of selected commodities.
- *Energ. P 4-01*: Yearly company statement on the production of heat and electricity.
- *Energ. P 5-01*: Yearly company statement of retail trade in solid fuels.
- *Energ. P 6-01*: Yearly company statement on sources and distribution of fuels.
- *Energ. P 1-01*: Yearly company statement of manufacture branches.

Transport:

Road transportation:

- *SLOVNAFT a. s. Bratislava*: Production and selling of gasoline and diesel fuel.
- *The Ministry of Economy of the Slovak republic*: Fuel sales of gasoline, diesel and biofuels.
- *SAPPO – Slovak association of petrochemical industry*: Gasoline, diesel and LPG selling data.
- *Slovak Gas Trading Company SPP Inc.*: Selling of compressed natural gas at gas stations.

- *SAD, a. s. Zvolen; ARRIVA Slovakia; DP Košice, a.s. Košice; DPB a.s. Bratislava; SAD Prievidza, a.s.*: CNG consumption data from bus transportation companies.
- *Presidium of the Police Force of the Slovak Republic, the Department of Documents and Registration of the Presidium*: Numbers of road vehicles for each year.
- *Ministry of Transport and Construction of the Slovak Republic*: Cumulative mileage data, odometers data.
- *Slovak Technical Control Stations*: Information on mileages.

#### Railways:

- *Železničná spoločnosť Slovensko, a. s.*: Fuel consumption data and selected operation capacity of combustion engine driven locomotives in personnel railway transport.
- *Železničná spoločnosť Cargo Slovakia, a. s.*: Fuel consumption data and selected operation capacity of combustion engine driven locomotives in railway freight service.
- *CER Slovakia a. s.*: Fuel consumption data and selected operation capacity of combustion engine driven locomotives in railway freight service.

#### Navigation:

- *Slovak navigation and harbours Inc. Bratislava & Norwardia*: Diesel oil selling data from custom storage to navigation companies in Slovak harbours.
- *Small companies from lakes and dams*: Fuel consumption data during the season.

#### Aviation:

- *EUROCONTROL*: Fuel consumption, LTO cycles and emissions.

#### IPPU:

- *Operators*: Manufacturers, importers, exporters and service, assembling organizations reported over by refrigerant.

#### Agriculture:

- *The Research Institute for Animal Production Nitra*: Expert guaranty of emission inventory
- *The Statistical Office of the Slovak Republic*: Number of the livestock, sowing areas, harvested areas, harvested yield.
- *The Breeding Services*: Detailed dividing of cattle and sheep
- *The Research Institute for Animal Production*: Animal production data.
- *The Central Controlling and Testing Institute in Agriculture*: Synthetic and organic fertilizers (sewage sludge, compost) applied to the soils, liming and urea application on soils, liming and urea application on the soil, pH of soils.

#### Waste:

- *COHEM SAŽP (Waste Management Centre of the Slovak Environmental Agency)*: Industrial solid waste data.
- *ÚRSO – Regulatory Office for Network Industries*: Data on methane recovered from SWDSs.
- *ACE (the Association of Experts on Waste Water Treatment)*: Data on sewage sludge management.
- *Duslo a. s.*: Data on ISW incineration.
- Websites of several companies and institutions are also used for the inventory: *OLO, KOSIT, Slovnaft, Duslo, NsP Prievidza, Fecupral, Ecorec*.

### 1.2.7. Brief Description of Key Categories

Key categories were assessed by Approach 1 by the level of emissions in years 1990 and 2023 and the trend in emissions for the year 2023 with and without LULUCF categories and those key categories have been chosen, whose cumulative contribution is less than 95%. The identification includes all reported greenhouse gases CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs and SF<sub>6</sub> and all IPCC source categories with LULUCF categories (in absolute values) performed with the detailed categorization. The detailed key categories were assessed and are listed in [Annex 1](#) of this Report.

In 2023, the Slovak Republic determined using the Approach 1 by the level assessment, 28 key categories with LULUCF and 24 key categories without LULUCF. In 2023, the Slovak Republic determined using the Approach 2 by the level assessment, 16 key categories with LULUCF and 19 key categories without LULUCF. In 2023, the Slovak Republic determined using the Approach 1 by the trend assessment, 32 key categories with LULUCF and 27 key categories without LULUCF. In 2023, the Slovak Republic determined using the Approach 2 by the trend assessment, 19 key categories with LULUCF and 25 key categories without LULUCF. List of key categories is almost identical for the base year 1990 and for the latest inventory year. The most important key categories are fuel combustion in energy sector for CO<sub>2</sub>, road transport, forest land, direct N<sub>2</sub>O emissions from agricultural soil or methane emissions from SWDS.

### 1.2.8. General Uncertainty Evaluation

The uncertainty assessment by the Approach 1 is enclosed in [Annex 3](#) to this report. Quantification of emissions uncertainty by level and trend assessment was calculated by using Approach 1 method published in the IPCC 2006 GL. The Approach 1 with the LULUCF estimated the 11.62% level uncertainty and the 6.14% trend uncertainty in 2023. Approach 1 without LULUCF estimated the 2.76% level uncertainty and the 1.15% trend uncertainty in 2023. According to the previous recommendations, Slovakia is using hybrid combination of Approaches 1 and 2 in this submission for calculation of total uncertainty of the inventory ([Annex 3](#)). Uncertainty analyses performed by the Approach 1 in transport was carried out using Table 3.2 for uncertainty calculation and country specific uncertainties for activity data and emission factors were inserted into calculation table.

The Slovak Republic provided also Approach 2 for uncertainty analyses according to Chapter 3 of the IPCC 2006 GL for the complete Energy and Waste sectors for the year 2015 (latest results). The methodology and results were published and described in previous SVK NIR 2018. Based on the latest Improvement Plan ([Chapter 1.2](#)), Monte Carlo calculation in the IPPU sector was updated in this submission and the results can be found in the [Chapter 4.2.1](#) of this Report. Approach 2 in the Agriculture sector is provided in this submission. Uncertainty evaluation is based on Monte Carlo method. Results and methodology are described in the [Annex A.5.1](#) of this Report. Approach 2 in the LULUCF uncertainty analyses was updated in this submission, too. Uncertainty evaluation is based on Monte Carlo method. Results and methodology are described in the [Annex A.6.2](#) of this Report.

Uncertainty assessment by Approach 1 on level assessment for the base year 1990 represents 2.86% without LULUCF and 8.91% with LULUCF. More information is in [Annex 3](#).

### 1.2.9. Completeness

Assessment of completeness is one of the elements of quality control procedure in the inventory preparation on the general and sectoral level. The completeness of the emission inventory is improving from year to year and the updates are regularly reported in the NIDs. The completeness checks for ensuring time series consistency is performed and the estimation is completed in recent inventory submission (2024). The improvements were performed in the previous inventory submissions such as estimation of GHG emissions for the agriculture and transport.

The list of categories reported by the notation keys is provided in CRT table 9. Whole overview of notation keys with detailed explanation is provided in [Table A2.1](#) with information on notation keys used

for each sector was prepared. More information can be found in [Annex 2](#) of this Report. Information is divided to the sectors and categories. Several categories are reported as not occurring (NO) due to the not existence of the emission source or the source is out of threshold and measurement range. If the methodology does not exist in the IPCC Guidelines, the notation key not applicable (NA) was used. Several NE key categories have been reported in 2025 submission for 1990 – 2023. Three reasons for not estimated (NE) categories are:

- no methodology is available;
- potential emissions/removals will under the threshold level of emissions in comparison to GHG emissions total;
- insufficient activity data (mostly for indirect GHG emissions like CO<sub>2</sub>, SO<sub>2</sub> or NMVOC).

**Table 1.6: List of NEs in the 2025 submission**

GAS	SECTOR	CATEGORY	DESCRIPTION
CO <sub>2</sub>	Energy	1.B Fugitive emissions from fuels > Oil and natural gas and other emissions from energy production > Oil > Refining/storage	Change of notation according to FCCC/ARR 2019 recommendation E.38; emissions are not estimated because the 2006 IPCC guidelines do not include methodologies to estimate these emissions.
CH <sub>4</sub>	Energy	1.B Fugitive emissions from fuels > Oil and natural gas and other emissions from energy production > Oil > Distribution of oil products	Change of notation according to FCCC/ARR 2019 recommendation E.38; emissions are not estimated because the 2006 IPCC guidelines do not include methodologies to estimate these emissions.
CH <sub>4</sub>	IPPU	2.C Metal Industry/2.C.1 Iron and Steel Production/2.C.1.b Pig Iron	Used methodology does not allow to distinguish the emissions
CH <sub>4</sub>	IPPU	2.C Metal Industry/2.C.1 Iron and Steel Production/2.C.1.e Pellet	Used methodology does not allow to distinguish the emissions
CH <sub>4</sub>	IPPU	2.D Non-energy Products from Fuels and Solvent Use/2.D.1 Lubricant Use	No methodology is provided neither in the IPCC 2006 GL not in IPCC 2019 Refinement.
CH <sub>4</sub>	IPPU	2.D Non-energy Products from Fuels and Solvent Use/2.D.2 Paraffin wax use	No methodology is provided neither in the IPCC 2006 GL not in IPCC 2019 Refinement.
N <sub>2</sub> O	IPPU	2.D Non-energy Products from Fuels and Solvent Use/2.D.1 Lubricant Use	No methodology is provided neither in the IPCC 2006 GL not in IPCC 2019 Refinement.
N <sub>2</sub> O	IPPU	2.D Non-energy Products from Fuels and Solvent Use/2.D.2 Paraffin Wax Use	No methodology is provided neither in the IPCC 2006 GL not in IPCC 2019 Refinement.
CO <sub>2</sub>	Agriculture	General	Indirect CO <sub>2</sub> emissions are not estimated in agriculture due to a lack of available methodology on atmospheric oxidation.
N <sub>2</sub> O	Agriculture	General	Part of the indirect emissions of N <sub>2</sub> O are included in the sectoral tables for manure management and agricultural soils indirect emissions from other than agricultural sources are not estimated.
N <sub>2</sub> O	Agriculture	3.D Agricultural Soils/3.D.1 Direct N <sub>2</sub> O Emissions From Managed Soils/3.D.1.6 Cultivation of Organic Soils	The emissions are under the threshold of significance. See NID Chapter Agriculture.

Categories included elsewhere (IE) are listed also in CRT table 9 with the explanations of reallocation.


Both direct and indirect GHGs as well as precursor gases are covered by the inventory of the Slovak Republic. The geographic coverage is complete; the whole territory of the Slovak Republic is covered by the inventory.

# CHAPTER 2. TRENDS IN GREENHOUSE GAS EMISSIONS

## 2.1. Description and Interpretation of Emission Trends for Aggregated GHG Emissions

The GHG emissions presented in the National Inventory Document 2025 were updated and converted by using the newest available methods, national conditions and data published by the Statistical Office of the Slovak Republic and other official statistical authorities. The improvements for the categories included in the Improvement Plan and prioritisation according to Recommendation Lists were implemented in previous submission. Total GHG emissions were 36 114.74 Gg CO<sub>2</sub> eq. in 2023 (without LULUCF and with indirect emissions). This represents a reduction by more 51% in comparison with the reference (base) year 1990. In comparison with 2022, the emissions decreased by 2%. Total GHG emissions in the Slovak Republic decreased in 2023 in comparison with the previous year by almost 800 kt, which was affected by the decrease in the Energy and IPPU sectors (mostly in the EU ETS sources) because of decreasing of industrial production in Slovakia. Total GHG emissions excluding the LULUCF sector have been decreasing continually from the base year and more stable trend in the recent years, dropped significantly in the years 2019 and 2020 due to special circumstances connected with the COVID-19 and other important changes made in Slovak economy. Then during the year 2021, emissions increased due to recovery of economy and afterwards due to Ukraine war, emissions decreased in 2022 affected by the increasing prices for fossil fuel. Significant changes in methodologies and emission factors are implemented in the frame of trying to keep consistency with the European Emission Trading System (EU ETS). **Table 2.1** shows the aggregated GHG emissions. In the period 1990 – 2023, the total greenhouse gas emissions in the Slovak Republic did not exceed the level of the base year 1990. **Figure 2.1** shows trends in the gases without LULUCF in relative expression.

**Figure 2.1:** The aggregated GHG emission trends



*Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of January 2025.*


This important reduction of emissions has resulted above all from the strong although temporary decrease in economy activities, followed by restructuring of economy joined with implementing new and more effective technologies, reducing the share of the intensive energy industry and increasing share of services in GDP generation. Transport (mostly the road transport), with increasing emissions is an important exception.

Total anthropogenic greenhouse gas emissions by gases in the years 1990 – 2023 are depicted in **Table ES.2** in this Report.

Beside the basic macroeconomic indicators as GDP, GDP per capita, foreign and domestic trade development, inflation, employment, there are also mentioned the data on the amount of investment in environmental protection and activities in the area of science and research, without specifying their orientation. The economic crisis that began in 2008 has brought a significant weakening of the external demand, causing a decreasing dynamic of the Slovak export, manufacturing, labour market and total domestic demand. The debt crisis in the Eurozone that broke out in 2012 again caused a decline in external demand. Emission situation in Slovakia can be considered and evaluated separately. While the EU ETS sources/sectors is going to further reduction of their emissions, the emissions in the non-EU ETS sources (ESR sectors/sources) is mostly stabilised or negative. Regulations included in the EU ETS push sources via economical instruments (Modernisation Fond) into larger investments and reduction of CO<sub>2</sub> emissions. In addition, the Slovak economy introduced changes in energy industry and steel production (phase-out of the furnace in the U.S. Steel company) what have positive effect on emissions in the EU ETS part of inventory. On the other hand, non-EU ETS sources representing agriculture, small industry, transport, waste and other small sources have not effective mitigation measures in place and the sectors policies are not targeting emissions reduction in a sufficient way. Therefore, the Ministry of Environment prepared the new Climate Change legislation, what introduces the sectoral targets with the shared responsibility among the ministries and the private sectors.

The indicators can assess the current economic and emission situation in Slovakia. While the indicator of carbon intensity can be changed much more rapidly in the situation of a high economic growth, GHG per capita is a different case where you can get very impressive results even without any measures, just by higher population growth rate. However, this is not the case of the Slovak Republic right now. It will take much longer time to change numerator by the impact of new technologies implementation namely in combination with high dynamic of development in the energy intensive industries. However, the indicator reached the lowest level in 2020. This was caused by combination of above mentioned measures and special situation with COVID-19, Ukraine war and fuel prices policy in the last few years.

**Figure 2.2:** Total GHG emissions in Gg of CO<sub>2</sub> eq. per capita in 1990 – 2023



Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.

---

## 2.2. Description and Interpretation of Emission Trends by Gas

Population of the Slovak Republic as of December 31, 2023 was 5 424 687 and has slightly decreasing. Average residential density is 110.7 inhabitants per km<sup>2</sup>. The population is concentrated in towns in lowlands and the main basins. Mountain areas are randomly populated. Unemployment rate in the Slovak Republic was 5.6% at the end of 2023 (according to the national statistics), what is lower than the previous years. The capital Bratislava is the largest city in the Slovak Republic with the number of inhabitants 475 500.

Total anthropogenic emissions of carbon dioxide excluding LULUCF and including indirect emissions have decreased by more than 50% in 2023 compared to the base year (1990). Nowadays the amount is 30 807.32 Gg of CO<sub>2</sub> without LULUCF and with indirect emissions. Compared to the previous inventory year 2022, emissions decreased by more than 2%. The reason for the decrease in CO<sub>2</sub> emissions in 2023 is caused by decreasing of CO<sub>2</sub> emissions in almost all energy and industry categories. Mainly in energy industry, manufacturing industry and in metal industry. In 2023, CO<sub>2</sub> emissions including the LULUCF and including indirect emissions significantly decreased compared to the previous year and decreased by 55% compared to the base year.


Total anthropogenic emissions of methane without LULUCF and with indirect emissions decreased compared to the base year (1990) by more than 58% and currently the emissions are 3 409.35 Gg of CO<sub>2</sub> eq. In absolute value, CH<sub>4</sub> emissions were 121.76 Gg without LULUCF. Methane emissions from the LULUCF sector are 0.54 Gg of CH<sub>4</sub> caused by forest fires. These emissions, however negligible, are decreasing due to lower number of forest fires in Slovakia. Trend of methane emissions is affected by the implementation of new waste legislation and measures in fugitive emissions and agriculture. Compared to the previous inventory year 2022, the amount of emission is decreased by more than 2%, mostly due to declining emissions in energy and IPPU sectors.

Total anthropogenic emissions of N<sub>2</sub>O without LULUCF decreased compared to the base year (1990) by more than 57% and currently the emissions are 1 445.47 Gg of CO<sub>2</sub> eq. Emissions of N<sub>2</sub>O in absolute value were 5.45 Gg without LULUCF. Emissions of N<sub>2</sub>O from the LULUCF sector are 0.10 Gg. Compared to the previous inventory year 2022, the emission increased by almost 14%, the increased was caused by Agricultural soils.

Total anthropogenic emissions of F-gases 452.60 Gg, from it 437.89 Gg of HFCs, 0.01 Gg of PFCs and 14.70 Gg of SF<sub>6</sub> in CO<sub>2</sub> eq. Emissions of HFCs decreased since 1995 due to the decrease in consumption and the replacement of PFCs and HFCs substances. Since that time, first decrease had occurred in the 2016 inventory year and repeated in 2018 and significant decrease continue in 2023. Decrease occurred in all F-gases and this is effect of implemented legislation of the EU in line with F-gases regulation ([Chapter 4](#)). Emissions' trend of PFCs has been decreasing and emissions of SF<sub>6</sub> has been slightly decreasing due to the decreasing consumption in industry. Decrease of F-gases emissions beginning in 2016 was caused by the biannual interval of servicing equipment. Despite this facts, emission of F-gases decreased compared to previous year 2022.



**Figure 2.3:** Emission trends by gas for the years 2000 – 2023 relative to the 1990 level (relative in %)




Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.

## 2.3. Description and Interpretation of Emission Trends by Category

The major share of CO<sub>2</sub> emissions comes from the Energy sector (fuel combustion, transport) with the 77.9% share from the total carbon dioxide emissions in 2023 inventory, 21.8% of CO<sub>2</sub> is produced in the IPPU sector and negligible amount is produced in the Agriculture (0.2%) and the Waste (0.01%) sectors. The energy related CO<sub>2</sub> emissions from waste incineration are included in the Energy sector. The 43.2% of CH<sub>4</sub> emissions is produced in the Waste sector (SWDS), 23% of methane emissions is produced in the Energy sector and 33.4% in the Agriculture sector. Almost 67% of N<sub>2</sub>O emissions is produced in the Agriculture sector (nitrogen from soils), 7.3% in the IPPU sector (nitric acid production), 13.6% in the Waste sector and 12% in the Energy sector. F-gases are produced exclusively in the IPPU.

**Figure 2.4:** Emission trends by gas in the sectors in 2023



Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.

Aggregated GHG emissions from the Energy sector based on the sectoral approach (combustion) data in 2023 were estimated to be 24 927.36 Gg of CO<sub>2</sub> eq. including transport emissions (7 735.54 Gg of CO<sub>2</sub> eq.), which represent the decrease by 56% compared to the base year and decrease compare to previous year by 2.6%. Transport decreased by 0.55% compared to 2022 and in comparison with the base year increased by more than 13%.

---

Total emissions from the IPPU sector were 7 292.59 Gg of CO<sub>2</sub> eq. in 2023, which was decreased by more than 22% compared to the base year and the decreased by 3% compared to the previous year. This sector covers also emissions from solvents use and indirect CO<sub>2</sub> emissions from solvents NMVOC emissions.

Emissions from the Agriculture sector were estimated to be 2 181.85 Gg of CO<sub>2</sub> eq. It is almost 63% decrease in comparison with the base year and the increase compared to the previous year was more than 10%. The Agriculture sector is the sector with the most significant decrease compared to the base year 1990, because of the decreasing trend in cattle numbers and fertilisers use.

Emissions from the Waste sector were estimated to be 1 671.86 Gg of CO<sub>2</sub> eq. The decrease is more than 4% compared to the previous inventory year and the time series are stable for last years. Compared to the base year, the increase was more than 41%, because of increased methane emissions from solid waste disposal sites. The emissions from waste incineration with energy use are included into the Energy sector, categories 1.A.1.a, 1.A.2.f and 1.A.2.c.

Structural changes in the Energy sector and the implementation of economic instruments have played an important role in achieving the status, when the trend of GHG emissions does not copy the fast GDP growth. In this context, the most important measure seems to be the adoption of the national legislation on air quality, which was approved in 1991 and it has initiated the positive trend in the reduction of the emissions of basic air pollutants and indirectly GHG emissions. At the same time, the consumption of primary energy resources as well as total energy has decreased.

Total anthropogenic greenhouse gas emissions by the sectors in the years 1990 – 2023 are depicted in [Table ES.2](#) in this Report.


According to the [Joint Research Centre of the European Commission](#), the highest reduction in the energy intensity values during the 15-years period from 2000 to 2014 was found in the Slovak Republic, which has undergone a growth rate of 82.5%.<sup>5</sup> This positive development is the result of the successful restructuring of industry, the introduction of energy-efficient production processes in industry and effective energy-saving measures in household by superseding home appliances with more efficient variants (due to several support programmes focused on households). Energy intensity in 2023 decrease in comparison with the previous year, due to decrease of the GDP caused by the economic reasons and lower total inland energy consumption. The latest year development estimated the long-term trend in energy intensity per GDP and final decarbonisation of economy.

Transport is a significant source of emissions in the Energy sector, with 28% share in total FEC (Final Energy Consumption) in the Slovak Republic. The proportion of transport is growing each year and the adopted policies and measures have no positive impact on increasing trend of emissions from transport. Emission balances in road transport are modelled according to method COPERT 5 version. GHG emissions from non-road transport are balanced by the use of EMEP/EEA 2023 methodology according to individual transport types (air, water and rail). The share of rail and water transports is decreasing from year to year, while the share of air transport increased rapidly in previous years, especially due to the increasing activity of low cost airlines, but the trend is stabilised recently. Slovak transport policy started to support railways and other alternative mode of transport (public, car sharing, etc.), but the effect of investments will be visible later.

---

<sup>5</sup> [Joint Research Centre: Energy Consumption and Energy Efficiency Trends in the EU-28 2000-2014 2016](#), p. 19.

**Figure 2.5:** The trend of energy intensity (right y axis) in the period 2007 – 2023 (estimated by the revised statistical approach NACErev.2)



Fugitive methane emissions from the extraction (only 0.4% share in total FEC) and distribution of fossil fuels were important, as the Slovak Republic is an important transit country regarding the transport of oil and natural gas from the third countries (East Europe, Asia) to Europe. Raw materials are transported through high-pressure pipelines and distribution network and they are pumped in pipeline compressors. During previous years, massive investments were introduced into transmission network to reduce fugitive emissions and losses. Further improvements were implemented by the specific distribution companies of oil and natural gas to the pipeline system (exploration, transit, distribution, etc.) in line with the international requirements. Side effect of these changes caused reducing fugitive emissions in this sector. New data and methodological approach for fugitive emissions from natural gas transmission was implemented into previous and current submissions.

The IPPU sector includes all GHG emissions generated from technological processes producing raw materials and products with the 28% share in total FEC in the Slovak Republic. Within the preparation of the GHG emission balance in the Slovak Republic, consistent emphasis is put on the analysis of individual technological processes and distinction between the emissions from fuel combustion in heat and energy production and the emissions from technological processes and production. Most important emission sources are balanced separately, emission and oxidation factors are re-evaluated, as well as other parameters entering the balancing equations and the results are compared with the verified emissions for CO<sub>2</sub> emissions. Fundamental emissions inventory in solvents is based on the balance of non-methane volatile organic compounds (NMVOC) according to EMEP/EEA 2023 methodology. Emissions are recalculated according to the stoichiometric coefficients to CO<sub>2</sub> emissions. Indirect emissions of CO<sub>2</sub> are estimated since the base year and allocated in the IPPU sector according to the IPCC 2006 GL.


The Agriculture sector with more than 1% share in total FEC in the Slovak Republic is the main source of methane and N<sub>2</sub>O emissions in the GHG emissions balance in the Slovak Republic. The emissions balance is compiled annually based on the sectoral statistics and in recent years based on a new regionalisation of agricultural areas of the Slovak Republic. The Ministry of Agriculture of the Slovak Republic issues annual statistics “Green Report”, part agriculture and food industry on a yearly basis. In recent year, the increasing trend of services and other (non-industrial) activities on GDP is visible. This has positive impact on the emissions. Slovakia is also providing to the EUROSTAT national accounts inventory of GHGs and pollutants according to the NACE rev.2 classification of economic activities. However, the methodology is different from the GHG inventory preparation, emissions trend shows interlinkages with the shift of GDP share of the economic sectors on total GDP of Slovakia.

The area of forest in the Slovak Republic covers approximately 40% of the territory and wood harvesting is historically an important economic activity. Since 1990, sinks from the LULUCF sector have remained at the level of 8-10% of total GHG emissions, but in the recent years, sinks increased on 15% level of the total GHG emissions. Historically stable trend was disrupted in 2004 by a wind calamity in the High Tatras, which resulted in increased harvest of wood damaged by the calamity and pests and consequently in the decrease in total sinks to the half of former volumes. The lower harvest and better management of forest caused increasing of sinks in the last years.

Several significant changes and re-evaluations of the applied methods have been carried out in the Waste sector, followed by recalculations in all categories of waste treatment. Methane emissions from solid waste disposal sites (SWDS) have the largest share on total emissions. Waste balance methodology has been revised and tier 2 approach FOD (First Order Decay) methodology has been used for the recalculations of the time series since 1950. The trend of methane emissions has been stabilised depending on the adopted legislation in municipal waste landfills, lower production of waste and higher share of recycling. A more detailed description of the methodology as well as with the Monte Carlo uncertainty analysis is described in the references.<sup>6</sup> The disaggregation of emissions from municipal waste incineration into two groups, i.e. waste incineration with and without energy utilisation, was another important change with respect to the quality improvement of the emission inventory. The emissions from waste incineration with energy utilisation were reported under the Energy sector, sub-category 1.A.1.a (other fuels). The emissions from waste incineration without energy utilisation are reported within the Waste sector, but are negligible in the present year. The comparison of the 2023 sectors share with the base year is shown on following **Figure 2.6**. The significant decrease is visible in the Energy and Agricultural sectors (without transport) and increase in the Waste and IPPU sectors and transport. Emissions from international aviation and shipping are excluded from the national totals and therefore not presented here.

International bunker emissions of the inventory are the sum of the aviation bunker and maritime bunker emissions. These emissions are reported as memo items; but excluded from national totals. Emissions of greenhouse gases from international aviation increased constantly between 1992 and 2008. Between 2009 and 2014, international bunker emissions decreased, partly reflecting the economic recession. Total GHG emissions from international transport represents 170.97 Gg of CO<sub>2</sub> equivalents in 2023, after dramatically decrease, practically stopping of air transport caused by Covid-19 pandemic situation in 2020-2021, emissions increase in 2023. Emissions from international aviation have more than 95% share.

**Figure 2.6:** The share of the individual sectors in total GHG emissions in 2023 (right) and 1990 (left)



Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.

<sup>6</sup> Szemesová, J.; Gera, M.: Emission estimation of solid waste disposal sites according to the uncertainty analysis methodology, Bioclimatology and Natural Hazards, ISBN 978-80-228-17-60.


### 2.3.1. Change in Emissions from Key Categories

Key categories are defined as the sources or removals of emissions that have a significant impact on the inventory as a whole, in terms of the absolute level of emissions, the trend, or both. The Slovak Republic prepared key categories analysis for 2023 and 1990 emission sources in line with the IPCC 2006 GL by using Approach 1. The quantitative analyses include combined uncertainty (on emission factors and activity data) and recognized key categories by level assessment with and without the LULUCF sector (**Chapter 1.2.12** and **Annex 1** of this Report).

CO<sub>2</sub> emissions from the category 1.A.3.b - Road Transportation are the largest key source remains accounting for 24% of total CO<sub>2</sub> emissions without LULUCF in 2023. Between 1990 and 2023, CO<sub>2</sub> emissions in road transportation increased by 3.04 Mt of CO<sub>2</sub>, which is 67% increase due to an increase in fossil fuel consumption (liquid) in this key category (**Figure 2.7**). Since 1990, the large increase in road transportation related CO<sub>2</sub> emissions was recognized. **Figure 2.7** shows that, solid fuels from the category 1.A.1 - Energy Industries, solid fuels is the key category without LULUCF (8.8%) with the largest decrease (78%; 10 Mt of CO<sub>2</sub>) is between 1990 and 2023. The main explanatory factors of emissions decrease are in improvements in energy efficiency and (fossil) fuel switching from coal to gas. CO<sub>2</sub> emissions from the category 1.A.2 - Manufactured Industry, solid fuels in the Energy sector are the third largest key source in the Slovak Republic, accounting for 10.1% of total GHG emissions in 2023. Between 1990 and 2023, emissions from this category showed the decrease by 65%.

CO<sub>2</sub> emissions from fuels in the category 2.C.1 - Iron and Steel Production are the largest key category without LULUCF in the IPPU sector, accounting for 11.8% of total CO<sub>2</sub> emissions in 2023. Emissions decreased by 10% in the comparison with the base year. A shift from solid and liquid fuels to mainly natural gas took place and an increase of biomass and other fuels has been recorded.


**Figure 2.7: Absolute change of CO<sub>2</sub> emissions by large key categories 1990 to 2023**



*Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.*

Methane emissions account for almost 9% of total GHG emissions in 2023 and decreased by almost 59% since 1990 to 121.76 Gg CH<sub>4</sub> without LULUCF in 2023. The three largest key sources (5.A - Solid Waste Disposal at 34%, 3.A - Enteric Fermentation at 30% and 1.B.2 Fugitive emissions from fuels - oil, NG and Other - Natural gas at 10% of total CH<sub>4</sub> emissions in 2023) account for more than 74% of CH<sub>4</sub> emissions in 2023. **Figure 2.8** shows that the main reasons for declining CH<sub>4</sub> emissions were reductions in enteric fermentation mainly caused by the decreased of animal numbers and use reductions in fugitive emissions and coal mining. **Figure 2.8** shows significant decrease in the categories 3.A and 3.B and increase in 5.A caused by the change of IPCC methodology used for solid waste disposal sites which considers time layer since 1960. Slight increase occurred also in the category 5.B - Biological Treatment of Solid Waste, due to changing in waste management praxis in Slovakia.


**Figure 2.8: Absolute change of CH<sub>4</sub> emissions by large key categories 1990 to 2023**



Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.

N<sub>2</sub>O emissions are responsible for 4% of total GHG emissions and decreased by 58% to 5.45 Gg of N<sub>2</sub>O without LULUCF in 2023 (Figure 2.9). The three largest key sources causing this trend in agriculture are 3.D.1 - Direct N<sub>2</sub>O Emissions from Managed Soils 32%, 3.D.2 - Indirect N<sub>2</sub>O Emissions from Managed Soils at 21% and 3.B - Manure Management at 15% of total N<sub>2</sub>O emissions in 2023. The main reason for large N<sub>2</sub>O emission cuts were reduction measures in the nitric acid production and decreasing agricultural activities (Figure 2.9). N<sub>2</sub>O emissions increased in biological treatment of waste and other products manufactured categories. This increase was caused by increase of operationalise and production.

**Figure 2.9: Absolute change of N<sub>2</sub>O emissions by large key categories 1990 to 2023**



Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of March 2025.

Fluorinated gas emissions account for 1.3% of total GHG emissions. In 2023, emissions were 452.6 Gg CO<sub>2</sub> eq., which was 111% above 1990 levels. The largest key source is 2.F.1 - Refrigeration and Air Conditioning and accounts for 96% of fluorinated gas emissions in 2023. HFC emissions from the consumption of halocarbons showed large increases between 1990 and 2023. The main reason for this is the phase-out of ozone-depleting substances such as chlorofluorocarbons under the Montreal Protocol and the replacement of these substances with HFCs (mainly in refrigeration, air conditioning, foam production and as aerosol propellants). On the other hand, PFC emissions decreased substantially since the base year. The decrease has started in 1996 and peaked in 1999 and 2001.

### 2.3.2. Main Reasons for Emission Changes in 2022 – 2023

Total GHG emissions in the Slovak Republic decreased by 2% in 2023 in comparison with the previous year, which was affected by the decrease in the Energy and IPPU sectors. This decrease demonstrates the economic and industrial impact of the energy prices policy, restrictions against the import of fossil fuels and raw materials from Russian Federation and development of electricity prices for industry. Several industrial subjects phased-out or reduced production or transformed. There were several significant changes in methodologies and emission factors implemented in the latest submission, particularly in fugitive emissions, agriculture and waste sectors. More changes were connected with the implementation of the 2019 IPCC Refinement.

The main reason for emission changes in 2021 – 2023 were as follows:

- CO<sub>2</sub> emissions decrease in the Energy sector - category 1.A.1 – Energy Industry (0.35 Tg of CO<sub>2</sub>) caused by decrease energy and heat production.
- CO<sub>2</sub> emissions decrease in the Energy sector - category 1.A.4 – Other Sectors (0.3 Tg of CO<sub>2</sub>) caused by decrease industrial production of heavy metals and chemistry.
- CO<sub>2</sub> emissions decrease in the Energy sector - category 1.A.3 – Transport (0.04 Tg of CO<sub>2</sub>) caused by increasing road transportation, mainly diesel-driven cars and transit.
- CO<sub>2</sub> emissions decrease in the IPPU sector – category 2.C – Iron and Steel Production (0.2 Tg of CO<sub>2</sub>).

In addition, methane emissions decreased in the Energy sector - category 1.A.4 – Other Sectors (0.2 Tg of CH<sub>4</sub>) and N<sub>2</sub>O emissions increased in the Energy sector - category 1.A.4 – Other Sector (0.5 Tg of N<sub>2</sub>O).

### 2.3.3. Key Drivers Affecting Emission Trends in LULUCF

The increasing trend of removals in forest land-use category is evident in the Slovak Republic since 1970. The increasing trend of removals cropland land-use category was recorded at the same time. Grassland areas decreased from 1980 to beginning of 1990 and since this year, decreasing trend of removals was recorded up to 2005. Since 2005, moderately downward trend has been taking place. Settlements land-use category has continual increasing trend during the whole period. This situation is mostly caused by development of transport infrastructure, industrial areas, municipal development and raising the standards and infrastructure in country and is very often connected with decreasing of the cropland and other land categories. Wetland represents 1.9% (94 kha) of the Slovak territory and it is considered constant, not involving any land use conversions. The LULUCF sector with net removals -7 775.62 Gg of CO<sub>2</sub> eq. in 2023 is very important sector and comprises from several key categories.

The major share from the LULUCF sector in 2023 represents removals in CO<sub>2</sub> with the contributions of the categories provided in the [Table 2.1](#). N<sub>2</sub>O emissions from the disturbance associated with the land-use conversion to Cropland, Grassland, Settlements and Other Land were reported in this submission. In addition, removals from the harvested wood products were estimated in this submission. The emissions of other pollutants originate from forest fires and controlled burning of forest. The estimated amount of NO<sub>x</sub> emissions was 0.35 Gg and the estimated amount of CO emissions was 12.31 Gg in 2023 ([Table 2.1](#)).

**Table 2.1:** Summary of total emissions and removals according to the categories in 2023

CATEGORY	Net CO <sub>2</sub>		CH <sub>4</sub>	N <sub>2</sub> O	NO <sub>x</sub>	CO	NM VOC	SO <sub>2</sub>
		Gg	Gg		Gg			
<b>4. LULUCF</b>	NO	-7 817.81	0.54	0.10	0.35	12.31	0.27	0.01
<b>A. Forest Land</b>	NO	-7 009.5	0.54	0.03	0.35	12.31	0.27	NO
<b>B. Cropland</b>	NO	-654.30	NO	0.03	NO	NO	NO	NO
<b>C. Grassland</b>	NO	-27.85	NO	0.00	NO	NO	NO	NO
<b>D. Wetlands</b>	NO	NO	NO	NO	NO	NO	NO	NO
<b>E. Settlements</b>	NO	76.24	NO	0.02	NO	NO	NO	NO
<b>F. Other Land</b>	NO	88.78	NO	0.02	NO	NO	NO	NO
<b>G. HWP</b>	NO	-291.90	NO	NO	NO	NO	NO	NO
<b>H. Other</b>	NO	NO	NO	NO	NO	NO	NO	0.01

Aggregated GHG emissions without LULUCF and indirect emissions; emissions are determined as of January 2025.

#### 2.3.4. Description and Interpretation of Emission Trends for Indirect GHG and SO<sub>2</sub>

Information can be found in **Chapter ES.5** of this Report.



---

**CHAPTER 3. ENERGY (CRT 1) .....50**

- 3.1. Overview of the Energy Sector..... 54
- 3.2. Fuel Combustion (CRT 1.A)..... 55
  - 3.2.1. *Overview of Fuel Combustion* ..... 55
  - 3.2.2. *Uncertainty Analyses of the Fuel Combustion* ..... 59
  - 3.2.3. *Category-specific QA/QC and Verification Process*..... 60
  - 3.2.4. *Category-specific Recalculations* ..... 62
  - 3.2.5. *Category-specific Improvements and Implementation of Recommendations* . 62
  - 3.2.6. *Energy Industries (CRT 1.A.1)*..... 64
  - 3.2.7. *Manufacturing Industries and Construction (CRT 1.A.2)*..... 71
  - 3.2.8. *Transport (CRT 1.A.3)*..... 76
  - 3.2.9. *Other Sectors (CRT 1.A.4)*..... 99
  - 3.2.10. *Non-Specified (CRT 1.A.5)* ..... 102
- 3.3. Comparison of the Sectoral Approach with the Reference Approach (CRT 1.AC) 104
- 3.4. Feedstocks and Non-energy Use of Fuels (CRT 1.AD) ..... 108
- 3.5. Fugitive Emissions from Fuels (CRT 1.B) ..... 110
  - 3.5.1. *Overview of Fugitive Emissions from Fuels*..... 110
  - 3.5.2. *Uncertainties and Time-series Consistency* ..... 111
  - 3.5.3. *Category-specific QA/QC and Verification Process*..... 112
  - 3.5.4. *Category-specific Recalculations* ..... 112
  - 3.5.5. *Category-specific Improvements and Implemented Recommendations*..... 115
  - 3.5.6. *Solid Fuels (CRT 1.B.1)*..... 115
  - 3.5.7. *Oil and Natural Gas and Other Emissions from Energy Production (CRT 1.B.2)*...  
..... 119
- 3.6. International Bunker Fuels (CRT 1.D.1) ..... 123
  - 3.6.1. *International Aviation (CRT 1.D.1.a)*..... 123
  - 3.6.2. *International Navigation (CRT 1.D.1.b)* ..... 124

## CHAPTER 3. ENERGY (CRT 1)

This Chapter was prepared using GWP<sub>100</sub> taken from the [5<sup>th</sup> Assessment Report of the IPCC](#) by the sectoral experts and institutions involved in the National Inventory System of the Slovak Republic:


INSTITUTE	CHAPTER	SECTORAL EXPERT
Faculty of Chemical and Food Technology, Slovak Technical University	Chapter 3.2 (except 3.2.8) Chapter 3.3	Juraj Labovský
Faculty of Chemical and Food Technology, Slovak Technical University	Chapter 3.4	Vladimír Danielik
Slovak Hydrometeorological Institute, Department of Emissions and Biofuels	Chapter 3.2.8 Chapters 3.5 & 3.6	Ján Horváth
External consultant	Chapter 3.3	Ján Judák

A significant decline in energy intensity was recorded in the previous years in Slovakia. The gross domestic energy consumption decreased by almost 16% since 2010. This decrease is associated with a decrease in solid and liquid fuels consumption for heating and with the significant decrease in natural gas consumption, while the electricity consumption is relatively stable. On the other hand, significant increase of biomass is visible. The share of different fuels on the primary energy consumption is as follow: natural gas 21.5%, nuclear fuel 46.7%, coal 21.2%, crude oil 5.9% and renewable sources (RES) more than 4.7% in 2023. Based on the National Energy Strategy up to 2030, an increase of nuclear and RES share on the total energy consumption is expected. A slight increase is projected in natural gas consumption in transport up to 2030 (transition fuel to zero-carbon 2050).


The most indicative decoupling trend in GHG emissions and GDP is visible directly in sector energy (fossil fuels consumption). The decrease in the consumption of solid fuels is 83% in comparison with the base year 1990. The consumption of liquid fuels decreased by 21% and the decline in gaseous fuels is 35%. By comparison, the consumption of biomass was 5 times higher in 2023 than in 1990. General trend in total consumption of fossil fuels is declining by 52% due to the increase in energy efficiency. [Figure 3.1](#) shows GHG emissions trend in Gg of CO<sub>2</sub> equivalents by categories for time series. Basic key categories 1.A.1 – Energy Industries, 1.A.2 – Manufacturing Industries and Construction and 1.A.4 – Other Sectors (services and households) have the most significant influence on the overall emission trends.

The Energy sector is the main contributor to overall GHG emissions with its share of 69% and 24 927.36 Gg of CO<sub>2</sub> eq. in 2023. Within this sector, [Figure 3.2](#) shows significant contributors (and key categories) to the emissions as follow: transport with its share of 30.3%, fuel combustion in the large (share 25%) and medium stationary sources of pollution (share 23%), pollution from small sources of residential heating systems (share 18.8%) and fugitive methane emissions from transmission/transport/distribution, processing and storage of oil and natural gas (share 2.7%).

**Figure 3.1:** Trend in aggregated GHG emissions by categories in Gg of CO<sub>2</sub> eq. within the Energy sector in 1990 – 2023



**Figure 3.2:** The share of aggregated GHG emissions by categories within the Energy sector in 2023




**Table 3.1** and **Figure 3.3** show overall emissions trends since the base year 1990 according to gases and major categories. The majority of emissions is reported in the category 1.A – Fuels Combustion (more than 90% in all years) and major gas is carbon dioxide (more than 90% in all years). According to this analysis, prioritization in the inventory preparation and improvements is set for the key categories within 1.A (using higher tier approach in key categories) and mostly focused on CO<sub>2</sub> gas (developing country/plant specific EFs for CO<sub>2</sub>).

**Table 3.1:** GHG emissions by categories within the Energy sector in particular years

YEAR	CO <sub>2</sub> EMISSIONS			CH <sub>4</sub> EMISSIONS			N <sub>2</sub> O EMISSIONS		
	Energy	1.A	1.B	Energy	1.A	1.B	Energy	1.A	1.B
<i>Gg of CO<sub>2</sub> eq./year</i>									
1990	53 349.7	53 272.9	76.86	3 362.93	507.58	2 855.35	229.54	229.37	0.18
1995	36 309.1	36 236.5	72.56	2 362.23	312.42	2 049.80	156.10	155.89	0.21
2000	34 008.6	33 938.4	70.16	1 920.85	264.33	1 656.52	157.24	156.99	0.25
2005	34 737.2	34 662.2	74.98	1 749.61	358.43	1 391.18	188.42	187.26	1.16
2010	30 895.6	30 824.0	71.66	1 335.03	355.40	979.63	178.80	178.60	0.20
2011	30 420.5	30 348.2	72.31	1 252.13	334.43	917.70	184.14	183.91	0.23
2012	28 355.3	28 291.5	63.82	1 047.58	359.43	688.15	188.71	188.49	0.22

YEAR	CO <sub>2</sub> EMISSIONS			CH <sub>4</sub> EMISSIONS			N <sub>2</sub> O EMISSIONS		
	Energy	1.A	1.B	Energy	1.A	1.B	Energy	1.A	1.B
	<i>Gg of CO<sub>2</sub> eq./year</i>								
2013	28 183.9	28 115.7	68.25	1 034.91	345.26	689.65	194.98	194.75	0.23
2014	26 007.7	25 933.8	73.95	875.13	227.13	648.00	193.68	193.46	0.21
2015	26 694.7	26 622.9	71.84	877.00	304.82	572.18	209.81	209.58	0.23
2016	26 875.3	26 793.2	82.04	952.33	322.25	630.08	206.50	206.33	0.17
2017	27 850.7	27 760.2	90.45	926.85	311.96	614.89	209.86	209.64	0.22
2018	27 784.5	27 689.5	94.99	840.79	264.36	576.43	201.10	200.88	0.22
2019	26 289.2	26 205.6	83.59	835.19	270.56	564.63	193.13	192.93	0.21
2020	24 110.9	24 030.5	80.38	819.88	269.11	550.77	176.48	176.25	0.23
2021	26 792.7	26 715.9	76.83	847.22	270.65	576.57	196.08	195.86	0.22
2022	24 609.4	24 521.2	88.20	791.31	252.40	538.91	187.43	187.22	0.21
2023	23 972.4	23 890.9	81.54	782.47	218.69	563.78	172.49	172.28	0.21

**Figure 3.3:** Trend in aggregated emissions by gases within the Energy sector in 1990 – 2023 (Gg of CO<sub>2</sub> eq.)



Sectoral approach based on bottom-up methodology is the most appropriate method for energy balance and for emissions estimation in the Slovak Republic. The sectoral approach is based on direct information from the large and medium stationary sources included in the EU ETS and completed with the statistical information provided by the Statistical Office of the Slovak Republic (ŠÚ SR) on the level of the statistical units (enterprises) – confidential data. Sectoral approach is compared with the reference approach based on top-down data published by the ŠÚ SR in the National Energy Balance (publicly available). The inter-annual fluctuation is very low and small discrepancies can occur in the fuel characteristics and using average parameters such as the calorific values or oxidation factors.

Fugitive GHG emissions in the period 1990 – 2023 were calculated based on the coal production from underground mines, obtained from the official statistical sources, mine companies (HBP, a.s., Baňa Dolina, a.s. and Baňa Čáry, a.s.), oil and NG production and transport companies, the ŠÚ SR and the Ministry of Economy of the Slovak Republic. A significant decrease in methane emissions in this category is visible in 2020, the situation was stabilised in 2021 while after 2021 further decrease was recorded. This is caused by the decrease of amount of coal mined and natural gas in transiting (therefore also fugitive emissions decreased inter-annual). In the end of the year 2023 the last coal mine was closed. This decrease was milder by methane emissions from abandoned mines.

The overview of categories according to the IPCC 2006 GL and 2019 Refinement to IPCC 2006 GL relevant for the Slovak Republic in the Energy sector is listed in [Table 3.2](#).

**Table 3.2: Reported emissions and tier approach in the Energy sector in 2023**

CATEGORY		DESCRIPTION / EMISSIONS / TIER					
1.A.1 Energy industries							
1.A.1.a	Public electricity and heat production	Electricity, combined heat and power generation, industrial and municipal waste incineration with energy use, cogeneration					
1.A.1.a.i	<i>Electricity generation</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.1.a.ii	<i>Combined heat and power generation</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.1.a.iii	<i>Heat plants</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.1.a.iv	<i>Other (waste incineration, methane cogeneration)</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.1.b	Petroleum refining	Refineries, petrochemical oil processing					
		CO <sub>2</sub>	T3	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.1.c	Manufacture of solid fuels and other energy industries	Coke production, coal manufacturing					
1.A.1.c.i	<i>Manufacture of solid fuels</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.1.c.ii	<i>Oil and gas extraction</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2 Manufacturing industries and construction							
1.A.2.a	Iron and steel	Iron, steel and ferroalloy production, manufacturing of iron ore					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.b	Non-ferrous metals	Non-ferrous metals production, casting					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.c	Chemicals	Chemical products manufacturing and production					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.d	Pulp, paper and print	Paper and pulp production, printing,					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.e	Food processing, beverages and tobacco	Food industry					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.f	Non-metallic minerals	Glass, cement, lime and magnesite production, brickworks, asphalt mixing plant, bating and electroplating					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g	Other						
1.A.2.g.i	<i>Manufacturing of machinery</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g.ii	<i>Manufacturing of transport equipment</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g.iii	<i>Mining (excluding fuels) and quarrying</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g.iv	<i>Wood and wood products</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g.v	<i>Construction</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g.vi	<i>Textile and leather</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.2.g.viii	<i>Other (industry not included above)</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.3 Transport							
1.A.3.a	Civil aviation - domestic aviation	CO <sub>2</sub>	T2	CH <sub>4</sub>	T3	N <sub>2</sub> O	T3
1.A.3.b	Road transport						
1.A.3.b.i	<i>Cars</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T3	N <sub>2</sub> O	T3
1.A.3.b.ii	<i>Light duty trucks</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T3	N <sub>2</sub> O	T3
1.A.3.b.iii	<i>Heavy duty trucks and buses</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T3	N <sub>2</sub> O	T3
1.A.3.b.iv	<i>Motorcycles</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T3	N <sub>2</sub> O	T3
1.A.3.b.v	<i>Other/Urea Based Catalysts</i>	CO <sub>2</sub>	M	-	-	-	-
1.A.3.c	Railways	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.3.d	Domestic navigation - domestic shipping	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.3.e	Other transport						
1.A.3.e.i	<i>Pipeline transport</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1

CATEGORY		DESCRIPTION / EMISSIONS / TIER					
1.A.4 Other sectors							
1.A.4.a	Commercial/Institutional	Commercial and institutional building, hospitals, schools					
1.A.4.a.i	<i>Stationary combustion</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.4.b	Residential	Sale fuels for households					
1.A.4.b.i	<i>Stationary combustion</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.4.c	Agriculture/Forestry/Fishing	Farms and forest organizations, slaughters					
1.A.4.c.i	<i>Stationary</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.4.c.ii	<i>Off-road vehicles and other machinery</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.5 Other							
1.A.5.a	Stationary	Compress and petrol stations, paint shops, wastewater treatment plants, crematory					
		CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.A.5.b	Mobile	Military aviation					
		CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.B.1 Solid fuels							
1.B.1.a	Coal mining and handling	Underground mines for brown coal, brown coal processing					
1.B.1.a.1.i	<i>Underground mines - mining activities</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T2	-	-
1.B.1.a.1.ii	<i>Post-mining activities</i>	-	-	CH <sub>4</sub>	T1	-	-
1.B.1.a.1.iii	<i>Abandoned underground mines</i>	CO <sub>2</sub>	T2	CH <sub>4</sub>	T1	-	-
1.B.1.b	Solid fuel transformation	Charcoal production and coke production					
		CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.B.2 Oil and natural gas and other emissions from energy production							
1.B.2.a	Oil						
1.B.2.a.2	<i>Production</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	-	-
1.B.2.a.3	<i>Transport</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	-	-
1.B.2.a.4	<i>Refining / Storage</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	N <sub>2</sub> O	T1
1.B.2.b	Natural gas						
1.B.2.b.2	<i>Production</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	-	-
1.B.2.b.3	<i>Processing</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	-	-
1.B.2.b.4	<i>Transmission and storage</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T3	-	-
1.B.2.b.5	<i>Distribution</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T3	-	-
1.B.2.b.6	<i>Other</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T1	-	-
1.B.2.c	Venting and flaring						
1.B.2.c.1	Venting						
1.B.2.c.1.ii	<i>Gas</i>	CO <sub>2</sub>	T1	CH <sub>4</sub>	T3	-	-
1.B.2.c.2	Flaring						
1.B.2.c.2.i	<i>Oil</i>	-	-	-	-	N <sub>2</sub> O	T1
1.B.2.c.2.ii	<i>Gas</i>	-	-	-	-	N <sub>2</sub> O	T1

### 3.1. Overview of the Energy Sector

The Energy sector covers emissions from fossil fuels combustion (CRT 1.A) and fugitive emissions from mines, oil and natural gas (CRT 1.B). The inventory of emissions from fuel combustion includes direct GHG emissions (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) and indirect GHG emissions (NO<sub>x</sub>, CO, NMVOCs), as well SO<sub>2</sub> emissions. Point sources, transport and other fuels combustion are included, too. The inventory of fugitive emissions from mines, oil and natural gas includes CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and NMVOCs emissions from brown coal mining, oil and natural gas refining and storage, the emissions from venting and flaring at oil refineries as well as, the emissions from natural gas transmission and distribution. The emissions from international bunkers (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, SO<sub>2</sub> and indirect gases) and CO<sub>2</sub> emissions from biomass are included in memo items and not included into national total.

---

## 3.2. Fuel Combustion (CRT 1.A)

### 3.2.1. Overview of Fuel Combustion

Fossil fuels combustion for energy and heat production (including transport) is the most important source of GHG emissions in the Slovak Republic. The GHG emissions in this sector represent the major share of total GHGs emissions in CO<sub>2</sub> equivalents. It is especially category of public energy production for power and heat supply, industrial energy production for electricity and heat supply for technological processes, road transport and district heating – heat supply for the residential sector (block of flats and dwellings), public and services buildings and other objects of the non-productive sector.

Total aggregated emissions from fuel combustion, including transport, based on sectoral approach represented 24 281.83 Gg of CO<sub>2</sub> eq. in 2023.

Beginning a year 2014, a minor temporary increase in CO<sub>2</sub> emissions was observed. This increase can be attributed to the economic growth of Slovakia. However, since the year 2017, the emissions further decreasing and this trend is continuing until present inventory years. The increase in liquid fuels consumption is most notably in transport. In 2023, the consumption of fuels decreased the first time in the transport sector without any major external disturbance (e.g. economic crisis, pandemic). The increase of biomass and other fuels (waste) consumption was notable. Emissions decreased more sharply in 2020 than in the previous period. There are several cumulative reasons for this decrease. Due to Covid-19 pandemic, a significant decrease in transport was observed. Similar decrease was observed in the category 1.A.4.a (services), especially in solid fuels. In addition, iron and steel production was significantly reduced. During the past five years, there have been significant fluctuations in the iron and steel production of the largest producer in Slovakia - the U. S. Steel, s. r. o. Košice. U. S. Steel, idled one of its three blast furnaces (June 2019 - January 2021). The inter-annual decrease in CO<sub>2</sub> emissions between 2018 and 2020 in fuel combustion was more than 13%. In the beginning of 2021, all furnaces in U. S. Steel, s. r. o. were put back into operation and the emissions returns to the values of 2018. Due to very high energy prices, low market demand and a sharp increase in steel imports, the steel production was significantly reduced in 2022. During the year 2022, up to two blast furnaces were gradually shut down. The reduction of CO<sub>2</sub> emission in the U. S. Steel, s. r. o. Košice was more than 1 650 kt CO<sub>2</sub>, which represents a decrease more than 18% in comparison with previous year. In 2023 the iron and steel production was partially recover. Due to a small market size of Slovakia, the iron and steel production can significantly influenced the emission trend in overall.

On the other hand, notable increase of CO<sub>2</sub> emission was observed in services and in households in 2021. The increase of fuel consumption in households sector was caused by colder climatic conditions in 2021, which was also represented by increase of heating degree days across all regions. In 2022, the fuel consumption in services and households decreased and returned to the values before 2021. In 2023, the decrease in emissions continued, and compared to 2022, a 9.4% year-on-year decrease can be observed.

Significant reduction of natural gas consumption was caused by technical problems in a large-scale power plant in Malženice. After the general maintenance (April 2022), the operation of the power plant could not be resumed due to the damage and subsequent shutdown of the generator. The combined cycle power plant outage lasted almost 10 months and the reduction of natural gas consumption was more than 385 mil.m<sup>3</sup>. In 2023, electricity production was restored and emissions in the 1.A.1.a.i sector increased by more than 355 kt of CO<sub>2</sub>.

In the last ten years, biomass consumption has been relatively constant. A significant change in this trend occurred in 2023. The decline in biomass consumption was at the level of 12.8% compared to 2022. The causes of this high drop can be identified in the Pulp, paper and print and household sectors. In the case of pulp, paper and print, the decrease is caused by a decrease in production and the related


decrease in the need for black liquor and solid biomass. In the case of a household, the reason is milder winters and a shorter heating season.

**Table 3.3** shows trend in GHG emissions by categories within the sectoral approach in particular years indicated the significant decrease in emissions followed by decrease in fuel consumption and switch of fuel's share (increasing of gas and biomass, decreasing of liquid and solid fuels) which is showed on **Figures 3.4** and **3.5**.

**Table 3.3: GHG emissions by categories in the 1.A - sectoral approach in particular years**

YEAR	1.A.1 ENERGY INDUSTRIES	1.A.2 MAN. INDUSTRIES AND CONST.	1.A.3 TRANSPORT	1.A.4 OTHER SECTORS	1.A.5 OTHER
	Gg of CO <sub>2</sub> eq./year				
1990	19 076.50	16 094.81	6 816.32	11 543.22	478.98
1995	11 917.42	11 809.02	5 490.92	7 208.06	279.39
2000	12 342.73	9 434.03	5 721.59	6 713.60	147.82
2005	12 125.38	8 576.38	7 693.08	6 717.37	95.72
2010	9 491.57	7 664.18	7 421.48	6 710.90	69.85
2015	8 076.34	6 768.99	7 294.17	4 933.84	63.93
2019	7 378.25	6 327.49	8 122.83	4 756.87	83.68
2020	6 752.18	5 930.99	7 061.47	4 662.12	69.11
2021	7 308.61	7 032.32	7 523.36	5 254.07	64.06
2022	6 407.46	5 922.85	7 778.60	4 789.54	62.38
2023	6 394.26	5 666.82	7 735.54	4 419.52	65.69


**Figure 3.4: Trend in fuels consumption within 1.A category in TJ in 1990 – 2023**



High-level dependency on import of primary energy sources (PES) is a limiting factor for the Energy sector in Slovakia and subsequently for the complete economic (mostly industrial) development of country. Net imports of PES are covered by almost 90% of the total energy demand.



**Figure 3.5: The share of fuels' consumption within category 1.A in 1990 and in 2023**



Energy Industries (CRT 1.A.1), Manufacturing Industries and Construction (CRT 1.A.2), Transport (CRT 1.A.3), Other Sectors (CRT 1.A.4) and Other (CRT 1.A.5) categories include emissions from fuel combustion in large and medium point sources (power plants, boilers and industrial plants with boilers and/or other combustion installations). Detailed emission trends by subcategories in particular years are presented in [Table 3.4](#).

**Table 3.4: GHG emissions by categories in the sectoral approach in particular years**

YEAR	1.A.1 ENERGY INDUSTRIES			1.A.2 MANUFACTURING INDUSTRIES AND CONSTRUCTION		
	1.A.1.a	1.A.1.b	1.A.1.c	1.A.2.a	1.A.2.b	1.A.2.c
	Gg of CO <sub>2</sub> eq./year					
1990	14 758.96	2 998.22	1 319.32	2 689.75	1 262.08	2 664.26
1995	8 403.78	2 209.76	1 303.87	2 454.39	534.68	3 067.04
2000	8 924.68	2 169.08	1 248.97	2 782.45	287.47	1 663.57
2005	8 677.58	2 098.93	1 348.87	3 397.87	188.47	875.43
2010	6 267.69	1 915.27	1 308.61	3 752.60	199.50	562.26
2015	4 969.16	1 817.08	1 290.11	2 874.94	139.23	484.56
2019	4 469.70	1 735.06	1 173.48	2 448.92	101.75	473.68
2020	3 960.11	1 814.27	977.80	2 185.10	98.06	474.47
2021	4 382.71	1 841.78	1 084.11	3 170.55	116.65	476.70
2022	3 323.16	1 897.53	1 186.77	2 480.03	81.68	450.31
2023	3 239.62	1 887.00	1 267.63	2 637.89	50.22	399.16

YEAR	1.A.2 MANUFACTURING INDUSTRIES AND CONSTRUCTION				1.A.3 TRANSPORT		
	1.A.2.d	1.A.2.e	1.A.2.f	1.A.2.g	1.A.3.a	1.A.3.b	1.A.3.c
	Gg of CO <sub>2</sub> eq./year						
1990	2 341.32	1 144.12	3 429.37	2 563.91	3.77	4 585.89	410.95
1995	1 215.04	761.49	1 838.65	1 937.72	2.68	4 112.70	220.30
2000	704.68	570.09	1 502.56	1 923.20	2.67	4 142.50	170.19
2005	547.82	436.90	1 390.07	1 739.82	7.85	6 240.52	115.43
2010	419.70	306.52	1 182.18	1 241.43	5.17	6 499.42	91.27
2015	499.68	329.64	1 248.30	1 192.64	3.68	7 005.91	93.72
2019	450.62	345.82	1 461.37	1 045.33	1.84	7 628.04	90.06
2020	406.97	342.71	1 423.93	999.74	0.89	6 806.56	80.63
2021	313.10	321.78	1 439.37	1 194.15	1.30	7 303.85	91.28
2022	261.08	310.37	1 326.05	1 013.32	1.49	7 664.14	91.48
2023	190.52	307.78	1 116.29	964.96	1.57	7 617.99	89.65


YEAR	1.A.3 TRANSPORT		1.A.4 OTHER SECTORS			1.A.5 OTHER	
	1.A.3.d	1.A.3.e	1.A.4.a	1.A.4.b	1.A.4.c	1.A.5.a	1.A.5.b
	Gg of CO <sub>2</sub> eq./year						
1990	0.02	1 815.69	4 166.56	7 220.89	155.78	407.24	71.73
1995	0.02	1 155.22	2 433.87	4 606.04	168.15	213.72	65.67
2000	0.02	1 406.20	1 570.15	4 771.32	372.13	130.58	17.24
2005	0.03	1 329.24	2 259.75	4 002.61	455.01	76.64	19.08
2010	0.33	825.29	2 571.84	3 732.13	406.93	54.08	15.78
2015	6.28	184.58	1 502.50	2 990.75	440.59	46.62	17.31
2019	4.21	398.67	1 350.47	3 079.14	327.26	72.42	11.26
2020	5.40	167.99	1 164.60	3 133.55	363.97	58.01	11.10
2021	5.88	121.05	1 464.26	3 413.28	376.53	52.87	11.19
2022	5.35	16.14	1 374.86	3 094.03	320.64	53.74	8.64
2023	5.37	20.95	1 343.35	2 802.45	273.72	61.34	4.35

The share of GHG emissions from stationary combustion (categories 1.A.1, 1.A.2, 1.A.4 and 1.A.5) to GHG emissions in Energy sector was 68.1% in 2023 (without transport). The share of solid fuels decreased from 42.14% in 1990 to 12.74% in 2023. By comparison, the consumption of biomass was 6.5 times higher in 2023 than in 1990. The share of biomass fuels increased from 1.45% in 1990 to 16.38% in 2023. General trend in total consumption of fuels is declining. Total consumption of fuels decreased by 43 % in comparison with base year ([Figure 3.5](#)). The highest share on GHG emissions has category 1.A.1.a - Public Electricity and Heat Production (13.31%), followed by 1.A.4.b - Residential (11.52%) and 1.A.2.a - Iron and Steel (10.84%) categories ([Figure 3.6](#)). The major share has category 1.A.3.b - Road transport (31.30%) which is the most important key category with one of the highest share on emissions in overall trend and in Energy sector. There is a significant decrease in CO<sub>2</sub> emissions in the category 1.A.2.c - Chemicals caused by the 99% decrease of solid fuels consumption. This decrease is significant and occurred in whole time series. However, the sharpest decrease occurred between 2001 and 2002. In 2001, there were only five plants in Slovakia, which used solid fuel as source of energy in chemical industry. In 2002, one of these plants stopped (significantly reduced) the production (ENERGETIKA, s. r. o. Strážske decreased by 355 Gg of CO<sub>2</sub> in solid fuels) and two others chemical plants reduced the production and also the consumption of solid fuels (CHEMES, a. s.,

HUMENNÉ decreased by 43 Gg of CO<sub>2</sub> in solid fuels, Duslo Šála, a. s. decreased by 43 Gg of CO<sub>2</sub> in solid fuels). In 2021, there was further significant decrease in the consumption of solid fuels. The main consumer of solid fuels (CHEMES, a. s) stopped the use of anthracite and residual fuel oil for heat production. In 2023, the consumption of solid fuels was practically the same as in the previous year.

A significant decrease can be observed also in categories 1.A.4.a - Services and 1.A.4.b - Households. This decrease is caused mainly by reduction of solid fuels combustion. The reduction of CO<sub>2</sub> emission from combustion of solid fuels is more than 90.8% percent in 1.A.4.a and 97.1% in 1.A.4.b in comparison with the base year. On the other hand, there is an increase of 57.0% in emission from natural gas in category 1.A.4.b in comparison with the base year.

**Figure 3.6:** The share of emissions in CO<sub>2</sub> eq. on different subcategories within 1.A in 2023



### 3.2.2. Uncertainty Analyses of the Fuel Combustion

According to the previous recommendations, Slovakia is using hybrid combination of Approaches 1 and 2 in this submission for calculation of total uncertainty of the inventory (**Annex 3** of this Document). Uncertainty analyses performed by the Approach 1 in the IPPU sector was carried out using Table 3.2 (IPCC 2006 GL) for uncertainty calculation and country specific uncertainties for activity data and emission factors were inserted into calculation table.

The Slovak Republic provided and published also Approach 2 for uncertainty analyses according to the Chapter 3 of the IPCC 2006 GL for the complete Energy and IPPU sectors for the year 2015. The methodology and results were described in previous SVK NIRs 2017 and 2018. The latest Monte Carlo simulation was performed in this sector for the year 2015. Due to capacity reasons and according to the QA/QC plan in this sector, new calculation of Monte Carlo uncertainty (approach 2) in the Energy sector and categories (including transport) will be performed in the next submission. For more information, please see the **Chapter 1.2** of this Document. Results of the Monte Carlo simulations were almost identical since this exercise was performed (since 2011).

CO<sub>2</sub> emissions from categories 1.A.1, 1.A.2, 1.A.3, 1.A.4 and 1.A.5 (liquid, solid and gaseous fuel combustion) are the most important key categories and they have a decisive effect on the management of level and trend uncertainties. The emission balance of other GHGs (CH<sub>4</sub>, N<sub>2</sub>O) from these categories was estimated by using the IPCC default methodology and default emission factors consistent with previous reporting.

AD, caloric value, EF and their uncertainties are available by the sectoral experts based on national circumstances. It helps to verify the correctness of aggregated uncertainty computation by Monte Carlo technique. EFs are expressed in t C/TJ. The changes and reallocations made in previous year are

included to the current inventory. The new categories 1.A.2.g and 1.A.3.e are added to analysis. Contrary, the subcategory 1.A.5.b was removed from analyses.

From the background data structure, differences between Approach 1 and Approach 2 (based on the IPCC 2006 GL) are concentrated to the correlation among inputs parameters in this case, because formulas, which are applied in the Approach 2, use only multiplication and addition operation. In this time, Approach 2 is computed without correlation, therefore Approach 1 and Approach 2 are well comparable. Approach 2 offers more reliable statistical results and shows more information about statistical structure of analysed uncertainty.

### 3.2.3. Category-specific QA/QC and Verification Process

The sector specific QA/QC plan is based on the general QA/QC rules and activities in specific categories. Information used in the process of preparation GHG emissions inventory of the energy sector was obtained from the different data sources:

- Statistical Office of the Slovak Republic, Department of Cross-Cutting Statistics (energy balance),
- National Emission Information System (database of all stationary emission sources),
- Emission Trading System (reports from operators and from verifiers),
- Questionnaires that were sent to the producers (in case of doubt).

More information on general QA/QC activities within the NS SR is included in the [Chapter 1.2](#) of this Document.

Emission balance in the Energy sector was prepared in the model taking into consideration also fuel balance in transport and IPPU. The sector specific QC activities were performed directly during calculation when checking several data sources for the emissions factors and other parameters. Activity data verification is processing with the cooperation of the ŠÚ SR and the NEIS experts including operators (or verifiers) in some cases. As it was already mentioned, the main source of activity data (and also NCVs and EFs) in current submission are verified EU ETS reports (plant level) and disaggregated data provided by the ŠÚ SR (enterprise level). New database system developed for fuels and emissions balance in the GHG inventory allows to perform several QC check more or less automatically.

In the category 1.A.1, more than 90% of emissions are cover by the EU ETS reports. The EU ETS activity data are compared against two independent sources: the NEIS database and disaggregated fuel consumptions provided by the ŠÚ SR. The Slovak Republic is providing information on the actual or estimated allocation of the verified emissions included in the EU ETS to the national GHG inventory. Further details can be found in the [Table 3.5](#). The emission from EU ETS are balanced foe energy and IPPU sector.

**Table 3.5:** *Actual allocation of the verified emissions reported by installations and operators under Directive 2003/87/EC for the year 2023*

CATEGORY	GAS	GHG INVENTORY EMISSIONS	VERIFIED EMISSIONS UNDER DIRECTIVE 2003/87/EC	VERIFIED EMISSIONS/ INVENTORY EMISSIONS
		<i>Gg of CO<sub>2</sub> or CO<sub>2</sub> eq.</i>		<i>Ratio in %</i>
Greenhouse gas emissions (total emissions without LULUCF for GHG inventory and without emissions from 1A3a Civil aviation, total emissions from installations under Article 3h of Directive 2003/87/EC)	Total GHG	34 928.04	16 994.18	50.30%
CO <sub>2</sub> emissions (total CO <sub>2</sub> emissions without LULUCF for GHG inventory and without emissions from 1A3a Civil aviation, total	CO <sub>2</sub>	30 803.38	16 947.45	55.24%

CATEGORY	GAS	GHG INVENTORY EMISSIONS	VERIFIED EMISSIONS UNDER DIRECTIVE 2003/87/EC	VERIFIED EMISSIONS/ INVENTORY EMISSIONS
		Gg of CO <sub>2</sub> or CO <sub>2</sub> eq.		Ratio in %
emissions from installations under Article 3h of Directive 2003/87/EC)				
1.A Fuel combustion activities, stationary combustion	CO <sub>2</sub>	16 546.29	10 105.8	61.08%
1.A.1 Energy industries	CO <sub>2</sub>	6 394.26	5 759.74	90.08%
1.A.2 Manufacturing industries and construction	CO <sub>2</sub>	5 666.82	4 346.06	76.69%
1.A.3 Transport	CO <sub>2</sub>	7 735.54	20.95	0.27%
1.B Fugitive emissions from fuels	CO <sub>2</sub>	81.54	NO	NA
2.A Mineral products	CO <sub>2</sub>	1 991.90	1 970.77	98.94%
2.B Chemical industry	CO <sub>2</sub>	918.20	1 101.35	119.95%
2.C Metal production	CO <sub>2</sub>	3 748.67	3 748.58	100.00%
2.B Chemical industry (Nitric acid production)	N <sub>2</sub> O	46.73	46.73	100.00%
3.C Metal production (Aluminium production)	PFCs	5.34	5.34	100.00%

\* Ratio verified emissions/inventory emissions is higher than 1, because downstream of CO<sub>2</sub> for urea production is reported in inventory.

Based on analyses, total GHG emissions verified under the EU ETS represent 48.65% on the total GHG emissions (without LULUCF and CO<sub>2</sub> from domestic aviation) based on March 15, 2025 inventory submission. The share of the EU ETS emissions is comparable with the share of the EU ESR emissions in the Slovak Republic, in a latest years the share of EU ETS emissions is decreasing. This progress was analysed and the resulting outcomes refer to increasing of energy effectivity and decreasing of emissions in large point sources included in the EU ETS scheme. The number of installations fell under the threshold to be included into the scheme and therefore, the EU ESR emissions increased inter-annually.

Total CO<sub>2</sub> emissions verified under the EU ETS represent 55.02% on the total CO<sub>2</sub> emissions (without LULUCF and domestic aviation) based on March 15, 2025 inventory submission.

Total N<sub>2</sub>O emissions verified under the EU ETS represent 0.13% on the total GHG emissions without LULUCF and domestic aviation) based on March 15, 2025 inventory submission. On N<sub>2</sub>O emissions, the share is 3.17%.

Total PFCs emissions verified under the EU ETS represent 100% on the total PFCs emissions based on March 15, 2025 inventory submission. Due to industrial circumstances with the aluminium production in Slovakia, in 2023 PFCs emissions decreased rapidly. More information [IPPU Chapter](#).

Basic QC procedures, which are performed for all recorded EU ETS data, can be distinguished in following categories:

- Comparison of aggregated site-specific data with the national statistics and/or EUROSTAT;
- Comparison of data across similar sites in individual CRT categories;
- Review significant changes in year-over-year estimates for individual plants, categories and subcategories;
- Comparison of direct measurements with estimates using a factor;
- Comparison of default factors to site or plant-level factors.

Information on activity data of the non-EU ETS sources obtained from the ŠÚ SR is compared and validated with the NEIS database. The NEIS database is referenced data set, not used directly for fuels

and emissions balance in the national inventory, but considered in the QC process of activity data and other available parameters.

The QC activities directly provided during data collection in the NEIS database run at two levels. The first level is verification provided by the regional environmental offices according to the national law and the second level is provided by the SHMÚ, the Department of Emissions and Biofuels. The process of data verification in the NEIS database must be completed by the end of July year x-1.

The background documents are archived by the sectoral experts and in central archiving system of NS SR at SHMÚ.

In line with the national rules applied in the EU ETS, annual publication of emission factors and NCVs used in the sectoral and reference approach of the GHG emissions inventory is publicly available.

This procedure is a part of the QC activity applied particularly in this sector and checked by the verifiers and operators of ETS sources.

Also according to agreement with CDV (Centrum dopravného výskumu) from 2023, there is a QA/QC cross-check between Slovak and Czech in transport sector emissions estimation, including parameters and factors.

### 3.2.4. Category-specific Recalculations

Sectoral experts made revisions of the methodological approach and used activity data also in 2025 submission. After analysis, improvement in biomass consumption in households introduced in this submission led to recalculation for years 2022 and 2023.

In line with the Improvement and Prioritization Plan for 2025 the following change was implemented in 2025 submission.

**Table 3.6:** Description of recalculation/reallocation implemented in 2025 submission

RECOMMENDATION NO.	CATEGORY	DESCRIPTION	REFERENCE
1.	1.A.3.b	Recalculation based on update of calculation model and EFs	<a href="#">Chapter 3.2.8</a>
2.	1.A.4.b	Recalculation of biomass consumption in the household sector	<a href="#">Chapter 3.2.9</a>
3.	1.B.1.a	Implementation of the IPCC 2019 RF to the IPCC 2006 GL and correction of calculations	<a href="#">Chapter 3.5</a>

**Ad.1.:** Please see [Chapter 3.2.8](#), implementation of new version of the COPERT model was included in this submission.

**Ad.2:** The mathematical model for estimating of biomass consumption in households has been improved. Recalculations were made in sector 1.A.4.b based on data from the new 2021 Census. The changes concerned the number of apartments connected to district heating system. This resulted in changes of biomass consumption for this sector. Detail information about the recalculation and comparison with previous data is provided in the [Chapter 3.2.9](#).

**Ad.3:** Please see [Chapter 3.5](#), implementation of the methodology based on the IPCC 2019 Refinement to the IPCC 2006 Guidelines.

### 3.2.5. Category-specific Improvements and Implementation of Recommendations

According to the draft ARR 2022 delivered on 28<sup>th</sup> February 2023, the [ERT recommendation E.2](#) regarding the category 1.A.4 Other sectors – solid fuels – methane emissions to estimate and report CH<sub>4</sub> emissions from solid fuels for category 1.A.4 using at least a tier 2 methodology (in accordance

---

with the 2006 IPCC Guidelines) if the emissions are identified as key. This issue is reflected below in this chapter and in the [Chapter 3.2.9](#).

In addition, during the inventory preparation, following room for improvements was identified for future submissions:

- Households represent serious issue related to achievement of the reduction commitments for the PM<sub>2.5</sub> emissions of the Slovak Republic. Air pollution and high emissions burden are mainly caused by the individual combustion of solid fuels in households, which produces emissions of total suspended particles (TSP) and their fractions (PM<sub>10</sub>, PM<sub>2.5</sub> and BC). This impacts also GHG emission inventory. Further cooperation with the Ministry of the Environment is in place; a new project LIFE for improvement of regional air quality requires also regional data on emissions from small sources. Therefore, additional statistical survey realised in 2022, improved emissions data on regional level, mostly included in biomass from households. More information can be find on [website](#).
- Regarding the growing demand for better quality of emissions data and missing input data required for further improvement of methodology, balances and inventories, the Slovak Hydrometeorological Institute, Department of Emission and Biofuels applied for the EUROSTAT subvention for the road transport data collection. The grant project began in 2021 and finished in March 2023. The results were implemented in road transport. More information can be find on [website](#).
- According to [the ERT recommendation E.2 from the draft ARR 2022](#), the ERT identified room for improvement in moving to higher tier approach (tier 2) in CH<sub>4</sub> and N<sub>2</sub>O emissions estimation for key fuels in energy. However, due to lack of information and absence of relevant study or report about types and numbers of combustion equipment in households and services (at most), this was not implemented, yet. More advanced and country specific EFs for non-CO<sub>2</sub> gases are essential for full implementation of higher tier. Moreover, in our Improvement Plan, an improvement in AD estimation has currently a higher priority. Moving to higher tier in category 1.A.4 is currently very difficult, as it covers large number of small sources. Category 1.A.4 covers two main sub-categories: households and services (agriculture is practically negligible). During last three years, several significant improvements in households' emissions inventory (1.A.4.b) were performed. These improvements were described and documented in previous submissions. Results were also published in several scientific journals and there is planning to be published also in future. This project was conducted together with the ŠÚ SR and the results were already implemented in the official statistical Energy Balance of the Slovak Republic. Statistical surveys in households were focused on the fuel consumption and energy balance in households with individual heating. This was used as inputs in mathematical model calculated fuels consumption in households. It was mainly focused on solid fuels and biomass; however, several improvements were performed also in other areas. The similar approach is planned to develop also for services, but this is budget related. The primary objective is to be able estimate the fuel consumption in buildings (non-residential). A relevant methodology to estimate the energy demand precisely and transparently and/or fuels consumption in non-residential buildings is the most essential. Therefore, according to the prioritization plan, the moving to higher tier method for CH<sub>4</sub> and N<sub>2</sub>O emissions was postponed and improvement of activity data in the category 1.A.4.a was prioritized in the Improvement Plan.

---

### 3.2.6. Energy Industries (CRT 1.A.1)

The CRT category energy industries 1.A.1 consists of the following subcategories: Public Electricity and Heat Production (CRT 1.A.1.a), Petroleum Refining (CRT 1.A.1.b) and Manufacture of Solid Fuels and Other Energy Industries (CRT 1.A.1.c). These subcategories are further divided based on the IPCC 2006 GL.

**Public electricity and heat production (1.A.1.a)** - this allocates GHG emissions from power installations to produce electricity and heat and the combined heat-power installations (CHP). Total volume of fuels reported here was 58 270.65 TJ in 2023. The most significant gas reported here was carbon dioxide, which represented 3 202.95 Gg of CO<sub>2</sub> in 2023. Total CH<sub>4</sub> emissions were 0.49 Gg and total N<sub>2</sub>O emissions were 0.086 Gg in 2023.

After significant decrease of emissions in years 2013 – 2014, trend was stabilized. Between 2018 and 2020 the decrease in CO<sub>2</sub> emissions was more than 5%. In 2022, a further significant decrease can be observed, when CO<sub>2</sub> emissions declined inter-annually by more than 30%. The decrease of solid fuels is continuous and visible in many facilities allocated in this category. Most important decrease in solid fuels was caused by thermal power plant in Vojany in year 2019, where the decrease of the semi-anthracite coal was more than 50%. The largest electricity producer in Slovakia (Slovenské elektrárne, a. s.) is undergoing the process of phase-out the coal consumption and replacing it with biomass. Similarly, one of the largest heat plants in Eastern Slovakia (TEKO, a. s.), reduced coal consumption in 2021 by more than 33% in comparison with year 2019 and more than 42% in comparison with year 2018. The decline in coal consumption in TEKO, a. s continued in 2022 and reached a value of 30% in comparison with year 2021. The sharp decline in solid fuels ended in 2023 and consumption stabilized at a value comparable to 2022.

On the other hand, natural gas consumption in this sector has a growing trend. The sharp increase of natural gas consumption in 2019 was caused by ZSE Elektrárne, s. r. o. power plant, it operates the combined cycle power plant near Malženice city in the Western Slovakia and currently, it is the biggest combined cycle power plant in Slovakia. Technically, it is based on a joint shaft connecting a gas turbine with 284 MW of capacity and a steam turbine with 152 MW of capacity, jointly total of 430 MW. The power plant was put into operation in 2010 and put out of operation due to unfavourable conditions on energy markets in 2013. Since August 2018, the power plant has new owner and was put into operation, again. A significant change in trend occurred at the beginning of 2022. Reduction of natural gas consumption in Malženice was caused by technical problems. After the general maintenance (February - April 2022), the operation of the power plant could not be resumed due to the damage and subsequent shutdown of the generator. The combined cycle power plant outage lasted almost 10 months and the reduction of natural gas consumption was more than 385 mil. m<sup>3</sup>. In 2023 the production was recovered and the electricity production increased from 31 to 932 MW/h. The interannual increase of emissions in category 356 Gg of CO<sub>2</sub>.

GHG emissions in the category 1.A.1.a are disaggregated into subcategories (electricity generation, combined heat and power generation, heat plants and other). This reporting is based on information provided by the ŠÚ SR (modules ENER 719 – ENER 721).

The category 1.A.1.a.ii includes also emission from cogeneration gas from mining activity for the years 2007 – 2014 (no CH<sub>4</sub> emissions from cogeneration occurred since 2015); and emissions from municipal solid waste incineration with energy use. These gases are used for electricity and heat production and therefore are reported in Energy sector. Methane emissions from waste incineration with energy use are excluded from the category 5.C – Incineration and Open Burning of Waste.

**Petroleum refining (1.A.1.b)** - GHG emissions from the refineries are allocated in the category 1.A.1.b. Refineries process crude oil into a variety of hydrocarbon products. The biggest refinery Slovnaft, a. s. is the only petroleum refining company operating in Slovakia, processing approximately 5.23 million tons



---

of crude oil in year 2023 (5.39 million tons of crude oil in 2022). This company is the most important supplier of petrol and diesel fuels in Slovakia (60% of market). Emissions from the petroleum refining, concern all combustion activities required to support the refining of petroleum products.

Within 1.A.1.b, the main emissions sources of fuel balance are oil, refinery gas and natural gas, which are used for heating and as sources of hydrogen for oil products processing (hydrocracking). Fuels are allocated to liquid and gaseous fuels categories. No solid fuel is combusted here.

Total volume of fuels allocated in 1.A.1.b expressed in energy units represented 27 754.51 TJ in 2023, practically identical to previous year (27 552.01 TJ in 2022). Total CO<sub>2</sub> emissions were 1 882.94 Gg. Total CH<sub>4</sub> emissions were 0.06 Gg and total N<sub>2</sub>O emissions were 0.0092 Gg.

**Manufacture of solid fuels and other energy industries (1.A.1.c)** - Total volume of fuels allocated in the 1.A.1.c expressed in energy units represented 6 740.87 TJ in 2023. Total CO<sub>2</sub> emissions were 1 267.26 Gg in 2023. Total CH<sub>4</sub> emissions were 0.0067 Gg and total N<sub>2</sub>O emissions were 0.0007 Gg.

### **Methodological Issues – Activity Data**

Tier 2 or/and tier 3 approaches are used for the majority of CO<sub>2</sub> combustion sources and country-specific emission factors are used for all fuels. CO<sub>2</sub> emissions estimation was performed based on the bottom-up approach. This is especially visible in the categories 1.A.1, 1.A.2 and 1.A.5 where emissions originated from the point sources (different approach is used in households and transport). For these sources, simple equations that combine activity data with emission factors are used.

The most important and essential methodological change in the sectoral approach was performed in 2013. Before year 2013, the primary source of activity data was the NEIS database.<sup>1</sup> Main reason for the mentioned modification was to increase the transparency of the sectoral approach.

The actual submission used activity data from verified reports of operators included in the EU ETS and individual statistical data of economical subjects in details (NACE rev.2 classification<sup>2</sup>) provided by the ŠÚ SR. The share of emission sources covered by the EU ETS in 1.A.1 is 89.7% and in 1.A.2 is 86.3%. The remaining sources allocated here are balanced by using ŠÚ SR data. After verification of the EU ETS reports by accredited verifiers, the EU ETS reports (in NIMs<sup>3</sup> formats) are released to the NS SR expert team. In the first step, the EU ETS reports are processed and transferred into internal database system (see below) in May, year-1. Activity data are directly linked to the specific IPCC categories based on the NACE rev.2 classification (provided by the ŠÚ SR).

This approach is used also for proxy inventory for the year-1. As in May, the official data from the ŠÚ SR are not available; the EU ETS reports are validated against the ŠÚ SR and the NEIS data from previous year, to check the time series consistency and trends. After releasing official data from the ŠÚ SR and the NEIS (October – November, year-1), the validation procedure (focused on identification of gaps in data) is repeated and all potential issues are recorded and prepared for further analyses. The EU ETS reports are directly used to prepare the background for the sectoral approach in 1.A.1, 1.A.2, 1.A.4 and 1.A.5. The EU ETS reports incorporate at least two levels of verification. The EU ETS reports are verified by the accredited verifiers in accordance with the legislative requirements before submission to the competent authority (districts offices). There are only five plants, which used measurement-based approaches. Emissions from measured-based approach are not directly used in the GHG inventory due to ensure consistency of the methodology and emission factors across IPCC categories. Therefore,

---

<sup>1</sup> The NEIS is the database of stationary sources, which collects the data on air pollutants and fuels consumption from the large and medium sources of air pollution in the Slovak Republic. These data are available in consistent time series since 2000, when the system NEIS was put in operation.

<sup>2</sup> Pan-European classification system of economic activities

<sup>3</sup> NIMs – National Implementation Measures.

---

these operators are directly contacted to provide further details on fuels consumption, characteristics and other relevant inputs for emissions calculation. Emissions are calculated by the sectoral experts of the NS. Calculated emissions can differ from the measurements, these differences are further analysed in the cooperation with the operators, verifiers and the Ministry of Environment of the Slovak Republic and used for emission inventory.

The activity data for the less energy efficient plant (not covered by the EU ETS = non-ETS) are obtained from the disaggregated energy balance data on plant level provided by the ŠÚ SR.<sup>4</sup> Official (verified) data from the ŠÚ SR are released to the SHMÚ in November year-1. These data are formed by several modules (Energ 719-721 and Energ 723-725). All modules are processed automatically and the information on fuels consumption is mapped to appropriate IPPC category. In similar manner, the fuel types used in individual modules are allocated to corresponding IPCC fuels' categories. This allows emissions estimate for all non-ETS plants. Data is completed with the EU ETS data and used for the sectoral approach balance in the categories 1.A.1, 1.A.2, 1.A.4 and 1.A.5.

The emissions balance in the categories 1.A.1, 1.A.2, 1.A.4 and 1.A.5 is done by combination and summation of activity data from the EU ETS reports and the ŠÚ SR database provided on plant level. This procedure is performed automatically by the internal database system. This system contains unmodified information about the fuel consumption and allows comparison of data from different sources. All fuels are linked automatically to the corresponding IPCC fuels categories. Individual plants in database are allocated to specific IPPC category based on the NACE rev.2 classification. This allows disaggregation of emissions into individual IPPC categories without modifying the original dataset.

In chemical industry, petroleum industry and iron & steel production, the allocation procedure is more complicated, and it is performed manually (plant specific) in a collaboration with the IPPU experts (detailed information is provided in the [Chapter 4](#) of this Document and in the [Annexes 4](#)). The material and emissions data flows are too complicated to split of technological (IPPU) and combustion emissions (Energy sector). Therefore, models on plant level are included in the main database. Models are prepared by the IPPU and energy sectoral experts and their methodological description is provided in appendix of this NID. The results of these models are presented in the form of simple input-output balance and the activity data from the EU ETS reports (or data from the ŠÚ SR) are replaced by the activity data calculated by the models. The background information for preparing models are obtained directly from the plant operators or the EU ETS verifiers. Data is validated against information from the standard databases and cross-checked by the energy and IPPU (or waste) experts. The cross checking is used to eliminate the issues with double counting, underestimated emissions or discrepancies with the IPCC 2006 GL. Based on the recent improvement in the EU ETS reporting, the comparisons were made for the apparent consumption of different fuels on plant (installation) level and for the allocation of production categories and harmonization with NACE rev.2 classification of installations.

For illustration, [Table 3.7](#) compares the share of GHG emissions in the individual IPPC categories based on the EU ETS data and the ŠÚ SR database. Very interesting is also comparison of the number of plants by the IPPC categories.

---

<sup>4</sup> These data are officially provided based on agreement between the MŽP SR, the SHMÚ and the ŠÚ SR.

**Table 3.7: Distribution of CO<sub>2</sub> emissions estimated by a different type of source of activity in 2023**

CATEGORY	CO <sub>2</sub> EMISSIONS		NUMBER OF COMPANIES	
	EU ETS	ŠÚ SR	EU ETS	ŠÚ SR
	%		No.	
1.A.1 Energy Industries	90.1	9.9	22	183
1.A.2 Manufacturing Industries and Construction	76.4	23.6	46	1 826
1.A.4 Other Sectors	0.0	100.0	0	556
1.A.5 Other (Not specified elsewhere)	0.0	100.0	0	70

Based on the information provided in **Table 3.7** is visible, that the EU ETS share of CO<sub>2</sub> emissions in 1.A.1 is 90.0% and in 1.A.2 is 79.7%. Due to high “EU ETS CO<sub>2</sub> emissions” share, it is possible to compare the activity data between three independent sources (EU ETS, ŠÚ SR and NEIS).

For fuel combustion in 1.A.1.b - Petroleum Refining, a plant specific, tier 3, bottom-up approach was used. Activity data obtained directly from the Slovnaft, a. s. (data on the amount of fuel combusted in individual sources, plant specific emission factors) was used for calculation of GHG emissions and compared with the information provided by the ŠÚ SR and the NEIS database.

In 1.A.1.b, emission factors for liquid fuels are plant specific. The emissions estimation is based on the tier 3 while the material and energy balances are provided directly by operator. This information is formed by monthly consumption of individual fuel types and emissions sources used in each operation unit in refinery. The CO<sub>2</sub> EFs and NCVs are evaluated experimentally in the company’s laboratory using the national standards. Certified measurements of emission factors for natural gas were provided by the Slovak Gas Company (SPP, a. s.). The main sources of fuel balance are oil, refinery gases, petroleum coke and natural gas, which are used for heating and as sources of hydrogen for oil products processing. Consumptions provided by the ŠÚ SR, NEIS and operator correlated very well. Refinery gas, for which country specific NCV and EF are used, is a mixture of various gases of different quality. The main type of refinery gas used in Slovnaft, a. s. a source of energy is fuel gas H1 produced by mixing natural gas and waste gases from the technological operations in mixers. The refinery gas and the imported natural gas are blended (in blenders H1 and H2) and distributed through the refinery fuel system. Natural gas is used to stabilize the pressure and qualitative parameters of fuel gases. The next part of balanced gasses are fuel gases from local networks, especially from production units R5 (FG-R5) and RHC (FG-RHC) and waste gases from pressure swing adsorption (PSA-HPP and PSA-V-KHK). Emission factors of these gasses are based on the statistical evaluation of the chromatographic analyses performed every month. These analyses are performed in the laboratory of quality control of the refinery, accredited by STN EN ISO 17025:2005. Residual fuel oils are liquid distillation residues from refinery processes. Samples of the fuel are analysed in the quality control laboratory, which meets accreditation standards ISO/IEC 17 025. Based on the analysis, the NCV, sulphur content and nitrogen content are estimated. The analyses are performed every day enabling the estimation of monthly averages of qualitative parameters.

Moreover, information provided by operator is practically identical to information, which is background for the EU ETS. Therefore, there is good (practically absolute) correlation between emissions reported under the EU ETS and the national inventory. This approach was introduced in submission 2013 and slightly modified based on the recommendations provided by the ERT in previous reviews. The emissions originally allocated in the 1.A.1.b were split and reallocated into three new subcategories. Emissions from ethylene production were shifted into 2.B.8.b and emissions from hydrogen production into 2.B.10. The background for mentioned disaggregation is based on the consumption of fuels in individual units for production of plastics and units producing hydrogen. This information is provided directly by the operator. In 2024 submission, the emissions from hydrogen production were reallocated from category 2.B.10 to 1.A.1.b. The reason for reallocation was implementation of the IPCC 2019 Refinement, where the new guidance for hydrogen production is presented. Based on the used

---

approach, the emissions from hydrogen production within a refinery as an intermediate product are primarily to support Energy sector activities and allocated here.

Greenhouse gas emissions were calculated for each emissions source by multiplying the fuel consumption (provided by the operator) by the respective emission factor. For calculation of CO<sub>2</sub> emissions, plant specific emission factors were used. CH<sub>4</sub> and N<sub>2</sub>O emission factors were taken from the IPCC 2006 GL.

#### **Municipal Solid Waste Incineration with Energy Use in the Category 1.A.1.a.i**

Municipal solid waste incineration with energy use is reported in 1.A.1.a.ii as other fuels. No emissions from the municipal solid waste incineration are reported in the category 5.C.1 Municipal Waste Incineration without energy use in the Waste sector because all incinerators of the MSW produce energy or heat in the Slovak Republic. Therefore, notation key “NO” is used in the 5.C.1. The MSW is combusted in two large stationary incinerators situated in Bratislava and Košice. Statistically negligible volume of MSW is incinerated outside of these two large plants. Industrial waste is incinerated mainly in cement and chemical industry, therefore these emissions are reported in the categories 1.A.2.f and 1.A.2.c.

#### **Emission Factors and NCVs**

The country specific calorific values of the fuels are announced by the ŠÚ SR published in the Statistical Yearbook annually. The variations depend on fuel characteristics. If an operator used the plant specific calorific values, it is an obligation to provide supported measurements and inform relevant competent authority. The plant specific data and results of measurements can be found also in the EU ETS reports.

The NCVs taken from the ŠÚ SR and the EU ETS reports are used in inventory. These were calculated as country specific average (annual weighted average NCV):

- NCV of primary and secondary liquid fuels in the RA are the same as statistical values;
- NCV of primary and secondary solid fuels and natural gas applied in the RA are based on the analysis in accredited laboratories;
- NCV values of solid fuels used in the ŠÚ SR and the RA are not significantly different.

According to the direct information on the quantity of fuels combusted (in kt or mil. m<sup>3</sup>) and their specific net calorific values, calculation of fuels consumption in energy unit (TJ) is provided. For fuel combustion and industrial processes, the following numerical data is reported in the EU ETS reports:

- mass or volume of fuels consumption;
- net calorific values of fuel;
- CO<sub>2</sub> emission factors;
- additional process material (carbonates).

Due to the high EU ETS emissions share in 1.A.1, the emission factors are estimated as weighted average of published emission factors for individual fuels in all installations allocated in this category. Averaged emission factors are subsequently used for estimation of CO<sub>2</sub> emission for plants, which are not covered by the EU ETS. CO<sub>2</sub> emission factors in refinery are plant specific (only one installation in 1.A.1.b).

The annual EU ETS reports are an important source of activity-specific and company specific data on CO<sub>2</sub> emissions, fuels and emission factors for major combustion sources (and industrial processes sources) in the national GHG emissions inventory. The EU ETS covers 98 sources in the total CO<sub>2</sub> emissions of 16 994 Gg in 2023.

For each fuel type, the default, country or plant specific emission factor is used and the corresponding emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O are calculated. The CO<sub>2</sub> emission factors are country or plant specific (and also IPCC category specific) derived from the national/plant fuel characteristics. Default carbon

emission factors (t C/TJ) are estimated for individual fuel types based on the international methodologies (IPCC, OECD, IEA) and/or national measurements (expert judgment of the sectoral experts, EU ETS reports, industrial association's measurements, and scientific papers). Carbon emission factors are estimated from fuel composition and available average net calorific values of the most used fuels. Average country specific CO<sub>2</sub> emission factors have been used for natural gas, coal, brown coal according to the source of origin (Slovak, Ukraine, the Czech Republic), coke and coke gas since 2000. The revised emission factors depend on net calorific values and slightly vary from year to year and across to IPCC categories. The emission factors for natural gas and other important fuels are based on precise measurements and calculation published every month by the SPP, a. s., the Slovak Energy Industry, a. s., refinery plant Slovnaft, a. s. (liquid fuels), and the U. S. Steel, s. r. o. for iron and steel production. These EFs are in use for the installations under the EU ETS and for the reporting requirements of the MŽP SR. Carbon content per unit of energy is usually lower for light refined products, such as petrol, than for heavier products such as residual fuel oil.

For natural gas, the carbon emission factor depends on the composition of the gas (in its delivered state it is primarily methane, but it can also include small quantities of ethane, propane, butane, and heavier hydrocarbons). In the Slovak Republic, the emission factor for natural gas (mostly of the Russian origin) is based on precise measurements and calculations published every month by the SPP, a. s. since the year 2000. The same EFs for natural gas are used for the installations covered by the EU ETS annually to ensure consistency across country. The emission factors and composition of NG are published monthly online (**Tables 3.8 - 3.10**). Weighted averages are calculated based on monthly consumption announced by the SPP, a. s. Despite the fact, that the SPP, a. s. is in the present days not exclusive natural gas supplier (approximately 60% of market), the parameters of the NG are consistent in all consumers due to the common origin of natural gas distributed by the SPP, a. s. – Distribution.

**Table 3.8: Composition of natural gas published on-line by the SPP, a. s. in 2023**

MONTH	CH <sub>4</sub>	C <sub>2</sub> H <sub>6</sub>	C <sub>3</sub> H <sub>8</sub>	i-C <sub>4</sub> H <sub>10</sub>	n-C <sub>4</sub> H <sub>10</sub>	i-C <sub>5</sub> H <sub>12</sub>	n-C <sub>5</sub> H <sub>12</sub>	neo-C <sub>5</sub> H <sub>12</sub>	C <sub>6</sub> H <sub>14</sub>	CO <sub>2</sub>	N <sub>2</sub>
	mol %										
I.	93.9236	3.4704	0.8906	0.1352	0.1402	0.0298	0.0222	0.0014	0.0309	0.6070	0.7482
II.	94.1570	3.4434	0.8897	0.1393	0.1367	0.0278	0.0198	0.0011	0.0252	0.4942	0.6655
III.	93.9676	3.4213	0.8354	0.1260	0.1331	0.0300	0.0223	0.0012	0.0334	0.6627	0.7665
IV.	92.8468	3.8068	1.0357	0.1453	0.1671	0.0385	0.0292	0.0013	0.0430	0.9263	0.9598
V.	92.6365	3.9882	1.1384	0.1589	0.1800	0.0400	0.0300	0.0012	0.0420	0.8762	0.9084
VI.	92.2230	4.4258	1.3247	0.1924	0.2070	0.0424	0.0312	0.0009	0.0413	0.7452	0.7656
VII.	92.4014	4.3654	1.2705	0.1846	0.1990	0.0411	0.0302	0.0009	0.0409	0.7238	0.7416
VIII.	93.1432	4.0209	1.2422	0.1898	0.1938	0.0387	0.0273	0.0006	0.0317	0.4891	0.6220
IX.	94.1687	3.4314	1.0860	0.1653	0.1664	0.0312	0.0220	0.0004	0.0199	0.3139	0.5941
X.	93.4282	3.7868	1.1418	0.1692	0.1749	0.0353	0.0253	0.0005	0.0288	0.5136	0.6950
XI.	93.1504	3.7701	1.0390	0.1474	0.1651	0.0365	0.0280	0.0013	0.0425	0.7495	0.8694
XII.	91.9157	4.2894	1.0805	0.1437	0.1815	0.0438	0.0361	0.0017	0.0678	1.1397	1.0984

**Table 3.9: Overview of the EFs and NCVs for natural gas [15°C; 101.325 kPa] published online by the SPP, a. s. in 2023**

MONTH	RELATIVE DENSITY	DENSITY	NCV	COMBUSTION HEAT	WOBBE NUMBER	SULPHUR CONTENT	EF C
	mol %	kg.m <sup>-3</sup>	kWh.m <sup>-3</sup>	kWh.m <sup>-3</sup>	kWh.m <sup>-3</sup>	mg.m <sup>-3</sup>	tCO <sub>2</sub> /TJ
I.	0.5953	0.7295	9.796	10.852	14.07	0.0631	56.02
II.	0.5935	0.7273	9.809	10.867	14.11	0.0667	55.95
III.	0.5949	0.7291	9.775	10.829	14.04	0.0570	56.03
IV.	0.6035	0.7395	9.810	10.865	13.97	0.0166	56.30
V.	0.6051	0.7415	9.855	10.913	14.03	0.0235	56.33

MONTH	RELATIVE DENSITY	DENSITY	NCV	COMBUSTION HEAT	WOBBE NUMBER	SULPHUR CONTENT	EF C
	<i>mol %</i>	<i>kg.m<sup>-3</sup></i>	<i>kWh.m<sup>-3</sup></i>	<i>kWh.m<sup>-3</sup></i>	<i>kWh.m<sup>-3</sup></i>	<i>mg.m<sup>-3</sup></i>	<i>tCO<sub>2</sub>/TJ</i>
VI.	0.6081	0.7452	9.954	11.021	14.13	0.0574	56.38
VII.	0.6067	0.7435	9.942	11.007	14.13	0.0393	56.34
VIII.	0.6016	0.7372	9.940	11.007	14.19	0.0882	56.16
IX.	0.5940	0.7280	9.872	10.935	14.19	0.0893	55.92
X.	0.5992	0.7344	9.887	10.950	14.15	0.0591	56.10
XI.	0.6012	0.7367	9.833	10.890	14.05	0.0203	56.19
XII.	0.6100	0.7476	9.836	10.892	13.95	0.0186	56.51
<b>AVERAGE</b>	-	-	-	-	-	-	56.18

**Table 3.10: Overview of country or plant specific CO<sub>2</sub> EFs in t/TJ used in the category 1.A.1 in 2023**

1.A.1.a	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	74.57	Gas/Diesel oil	20.35	74.62
		Residual fuel oil	20.80	76.27
		Liquefied petroleum gases	17.22	63.14
Solid	97.34	Anthracite	27.15	99.55
		Other bituminous coal	27.56	101.05
		Lignite	26.31	95.81
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	102.38	Other biogas	14.90	54.59
		Sludge gas	14.90	54.59
		Other primary solid biomass	27.30	100.10
		Wood/Wood waste	30.50	111.83
1.A.1.b	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	73.43	Residual fuel oil	21.59	79.15
		Petroleum coke	29.86	109.50
		Refinery gas	15.18	55.66
Gaseous	56.18	Natural gas	15.32	56.18
1.A.1.c	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	63.74	Liquefied petroleum gases	17.22	63.14
		Gas/Diesel oil	20.35	74.61
Solid	201.86	Lignite	26.89	98.60
		Coke oven gas	11.44	41.95
		Blast furnace gas	76.62	280.94
Gaseous	56.18	Natural gas	15.32	56.18

Default CO<sub>2</sub> emission factors from the IPCC 2006 GL are used only for biomass, which almost invariably refers to wood, wood wastes and biogas. The actually used fuels-specific EFs are in [Table 3.10](#).

In addition to CO<sub>2</sub> emissions, the fuel combustion in stationary sources results in the CH<sub>4</sub>, N<sub>2</sub>O, NO<sub>x</sub>, CO and NMVOCs emissions. Of these, CH<sub>4</sub>, and N<sub>2</sub>O account around 0.65% on the total GHG emissions (expressed in CO<sub>2</sub> eq.), in the Fuel combustion sector (CO<sub>2</sub>: 6 353.16 Gg; CH<sub>4</sub>: 15.56 Gg CO<sub>2</sub> eq.; N<sub>2</sub>O: 25.54 Gg CO<sub>2</sub> eq.). These emissions are influenced by many factors, including fuel type, equipment design, and emissions control technology. Therefore, it is inherently more complex and more uncertain than CO<sub>2</sub> emissions estimation. The non-CO<sub>2</sub> EFs are default based on the IPCC 2006 GL.

### Uncertainties and Time-series Consistency

According to the previous recommendations, Slovakia is using hybrid combination of the Approaches 1 and 2 in this submission for calculation of total uncertainty of the inventory ([Annex 3](#) of this Document). Uncertainty analyses performed by the Approach 1 in transport was carried out using Table 3.2 for

---

uncertainty calculation and country specific uncertainties for activity data and emission factors were inserted into calculation table.

The Slovak Republic provided and published also Approach 2 for uncertainty analyses according to the Chapter 3 of the IPCC 2006 GL for the complete Energy and IPPU sectors for the inventory year 2015. The methodology and results were described in previous SVK NIRs 2017 and 2018. The latest Monte Carlo simulation was performed for the 2015 emissions inventory. Due to capacity reasons and according to the QA/QC plan in this sector, new calculation of Monte Carlo uncertainty (Approach 2) in the Energy sector and categories (including transport) will be performed in next submissions. For more information, please see the [Chapter 1.2](#) of this Document. Results of the Monte Carlo simulations were almost identical since this exercise was performed (since 2011).

Time series is consistent in all aspects (methodological approach, country specific EFs and oxidation factor used, fuel characteristics, etc.) to the detailed level of disaggregation (on plant specific level).

### **3.2.7. Manufacturing Industries and Construction (CRT 1.A.2)**

Category 1.A.2 includes CO<sub>2</sub> emissions allocated in: Iron and steel (1.A.2.a); Non-ferrous metals (1.A.2.b); Chemicals (1.A.2.c); Pulp, paper, and print (1.A.2.d); Food processing, beverages, and tobacco (1.A.2.e); Non-metallic minerals (1.A.2.f) and Other (1.A.2.g). Emissions include industrial emissions originating largely from energy and heat production in raw materials and semi-manufactured goods production. The emissions reported here are related to fuel combustion, only. Consumption of fuels used as feedstock and reduction medium is not included in this category as it is allocated in the IPPU sector.

[Iron and steel \(1.A.2.a\)](#) – the iron and steel industry is one of the most energy intensive industrial branches in the Slovak Republic. Total volume of fuels allocated in 1.A.2.a expressed in energy units represented 18 588.43TJ in 2023.

The main iron and steel producer in the Slovak Republic - U. S. Steel, s. r. o. idled one of its three blast furnaces, whose total capacity is 4.5 million tonnes of raw iron a year, on June 2019. It did so in response to the situation on the European steel market which has been massively impacted by steel products imported into the European Union. The shutdown of the blast furnace led to a reduction in CO<sub>2</sub> emissions by more than 860 kt of CO<sub>2</sub> between years 2019 and 2018. In 2020, the decrease in emissions continued and total CO<sub>2</sub> emissions were at the level of 2 179 kt. From January 2021, iron production was resumed at all blast furnaces. As a result of the increase in iron production, there was an interannual increase in emissions at the level of 45% in 2021. Due to very high energy prices, low market demand and a sharp increase in steel imports, the steel production was significantly reduced in 2022. During the year 2022, up to two blast furnaces were gradually shut down. The reduction of CO<sub>2</sub> emissions was more than 1 650 kt CO<sub>2</sub>, which represents a decrease of more than 18%. This sharp fluctuation is reflected in all categories, where the emissions from steel production are allocated (1.A.1.c, 1.A.2.a, 1.A.2.m and 2.C.1). In 2023 the production was partially recovered and the iron production increased by 12%.

One of the most significant companies in this category (OFZ, a. s. Oravský Podzámok), significantly reduced its production, too. The production was limited provisionally until the end of the year 2022 due to continued negative market conditions, in particular high electricity prices and low market prices for ferro-alloys. OFZ consumed large amount of biomass and there was a 50% decrease in biomass consumption in OFZ. The decline in production continued in 2023 as well. Solid fuels decreased interannually by 86 %. Wood waste and solid biomass decrease practically to zero. As a result the and interannual decrease of biomass consumption in sector 1.A.2a is 56%.

Total CO<sub>2</sub> emissions were 2 633.23 Gg. Total CH<sub>4</sub> emissions were 0.074 Gg and total N<sub>2</sub>O emissions were 0.0098 Gg.

---

**Non-ferrous metals (1.A.2.b)** – this source covers combustion-related emissions from non-ferrous metal industry. Total volume of fuels allocated in 1.A.2.b expressed in energy units was 1 463.23 TJ in 2023. There was also a significant decline in emissions in the CRT category 1.A.2.b. The most pronounced declines are seen for natural gas and coal. At the end of 2021, the Slovalco company (producer of aluminium), reduced production volume to 80% of their capacity, and a further reduction in production volume to 60% of capacity was achieved from February 2022. The production of primary foundry alloys was stopped. The complete closure of primary aluminium production after 70 years of production took place at the beginning of January 2023 with the shutdown of the last of the 226 pots. In 2023, the consumption of solid fuels was completely stopped in one of the largest enterprises in this category (Veolia Utilities Žiar nad Hronom, a.s.). Veolia at the same time reduced biomass consumption by more than 50%.

Consumption of liquid fuels is similar in absolute values to previous years, but the mix of liquid fuels has changed. LPG consumption remained unchanged, however the consumption of residual fuel decreased practically to zero. The result of change in fuel mix caused a significant inter-annual change of IEFs.

Total CO<sub>2</sub> emissions were 48.92 Gg, total CH<sub>4</sub> emissions were 0.0206 Gg and total N<sub>2</sub>O emissions were 0.0027 Gg.

**Chemicals (1.A.2.c)** – includes emissions from fuels combustion in chemical industry. Chemical industry produces several different products such as chemicals, plastics or solvents. Total volume of fuels expressed in energy units allocated in 1.A.2.c was 7 000.41 TJ in 2023. In 2015, significant reduction of natural gas consumption occurred, which was caused by the termination of operation of one company with relatively high share of fuels in the period between 2016 and 2020. Natural gas consumption was almost constant. In 2022, a moderate decrease of emissions from gaseous fuels can be observed. The rate of decline increased in 2023. The internal change of gaseous fuels is 12%. This decline can be observed in most large enterprises (Continental Matador Rubber, Fortischem, Chemes).

There is a visible reduction in consumption of solid fuels. This trend is similar than in other categories, where solid fuels are replaced by natural gas and/or biomass. In year 2020, significant reduction in coal consumption occurred in the power plant Chemes, a. s., where the coal consumption decreases by more than 15%. In 2021, the major consumer of solid fuels (Chemes, a. s) stopped using anthracite and biomass. Therefore, the emissions from solid fuels decreased in 2022 practically to zero and the reduction of emissions is more than 99% in comparison with base year. In 2023 the consumption of solid fuels remains practically constant.

Total CO<sub>2</sub> emissions were 398.44 Gg, total CH<sub>4</sub> emissions were 0.01238 Gg and total N<sub>2</sub>O emissions were 0.0014 Gg in 2023.

**Pulp, paper and print (1.A.2.d)** – includes emissions from fuel combustion in pulp, paper and print industry. Total volume of fuels allocated in 1.A.2.d expressed in energy units was 17 623.94 TJ in 2023. There was a visible decrease of inter-annual energy consumption between 2015 and 2016 (27 472.11 TJ in 2015 and 22 926.55 TJ in 2016). It was caused by decrease of fuels consumption in three major plants allocated here. In 2021, a significant interannual change in fuel mix occurred in this category. Major emissions producer (Bukoza Energo) cut coal consumption in half (decrease in coal consumption was more than 60 thousand tons). The reduction in coal consumption was compensated by an increase in biomass consumption (increase in biomass consumption was 10% in 2021). The result of the change in the fuel mix is 10% decrease in emissions. Similar trend was visible also in 2022. The inter-annual increase of emissions from biomass is 3.5% and decrease of emissions from solid fuels was 15%. In 2023 Bukoza Energo completely stopped using other bituminous coal and lignite consumption was reduced by 70%. Similar situation is in Bukocel pulp mill, the most important part of the Bukóza holding. It is the second largest domestic wood processing company. At the end of 2023 it was facing collapse. The consumption of solid and biomass fuels halved compared to 2022. The company Mondi SCP has the largest share in the decrease in biomass. Biomass consumption has



grown for five years in a row until 2022. In 2023, there was a decrease in the consumption of sulphite lyes (black liquor) at the level of 112 kt and other solid biomass decreased by 32 kt. The decrease in biomass in Mondi SCP and Bukocel resulted in an interannual decline in biomass at the level 5 734 TJ (24.5%) in comparison with year 2022.

Total CO<sub>2</sub> emissions were 177.26 Gg, total CH<sub>4</sub> emissions were 0.131 Gg and total N<sub>2</sub>O emissions were 0.0362 Gg in 2023.

**Food processing, beverage and tobacco (1.A.2.e)** – total volume of fuels allocated in 1.A.2.e expressed in energy units represented 5 415.19 TJ in 2023. Total energy is and emissions are practically constant in last 10 years, however the fuels mix has been significantly changed in year 2021. One of the largest source in this category (Slovenské cukrovary) stopped producing heat from coal. The lignite was fully replaced with natural gas (therefore a increase in gaseous fuels is visible in year 2021). Very sharp increase in liquid fuels can be observed (in relative numbers) in 2023. This increase was caused by the start of using fuel oil in the company Považský cukor a. s., which is one of the largest food companies in Slovakia.

Total CO<sub>2</sub> emissions were 307.43 Gg, total CH<sub>4</sub> emissions were 0.0062 Gg and total N<sub>2</sub>O emissions were 0.0007 Gg in 2023.

**Non-metallic minerals (1.A.2.f)** – total volume of fuels allocated in 1.A.2.f expressed in energy units represented 16 681.75 TJ in 2023. The fuels are allocated in solid, liquid, gaseous, other and biomass fuels. From 2018, a significant decrease in solid fuels can be observed in sector 1.A.2.f. Even in 2023, there was a decrease in the consumption of in practically all cement plants (CEMMAC, Carmeuse Slovakia, Považská cementáreň, Danucem). The decrease in gaseous fuels was caused mainly by the decrease in natural gas consumption in the company Slovenské magnezitové závody, where the interannual decrease in consumption was at the level of 26%.

Total CO<sub>2</sub> emissions were 1 10270 Gg, total CH<sub>4</sub> emissions were 0.21 Gg and total N<sub>2</sub>O emissions were 0.0289 Gg.

**Other (1.A.2.g)** - The remaining emissions from fuels combustion in manufacturing and industry were allocated in this category. Total volume of fuels expressed in energy units represented 17 826.91 TJ in 2023. Energy and emissions are comparable to the previous period. A gradual decrease can be observed in all types of fuels. Significant fluctuations in the consumption of solid fuels occurred in 2020 and 2022. These fluctuations were caused by the reduced production of iron and steel. With lower iron and steel production, the share of coke oven gas in the fuel mix decreases. Coke oven gas has a very low value of the emission factor (compared to coal). Therefore, the implied emission factor of solid fuels also showed a significant fluctuation between years (2019 – 2022 and 2022 – 2023).

Total CO<sub>2</sub> emissions were 959.07 Gg, total CH<sub>4</sub> emissions were 0.095 Gg and total N<sub>2</sub>O emissions were 0.01217 Gg in 2023. Based on the IPCC 2006 GL, this category was further split into 8 subcategories. The distribution of individual plants into subcategories was done based on the NACE rev.2 classification. The distribution of emissions along this category is **Table 3.11**.

**Table 3.11: Disaggregation of CO<sub>2</sub> emissions across the subcategories of the 1.A.2.g in 2023**

SUBCATEGORY	CO <sub>2</sub> EMISSIONS	SHARE
	Gg/year	%
1.A.2.g.i Man. of machinery	138.68	14.46
1.A.2.g.ii Man. of transport equipment	171.48	17.88
1.A.2.g.iii Mining and quarrying	8.88	0.93
1.A.2.g.iv Wood and wood products	17.08	1.78
1.A.2.g.v Construction	46.87	4.89
1.A.2.g.vi Textile and leather	18.63	1.94
1.A.2.g.viii Other	557.45	58.12

## Methodological Issues – Activity Data

Detail description of the methodological issues and activity data used for estimation of emissions from fuel combustion is given in the [Chapter 3.2.6](#).

**Iron and steel (1.A.2.a)** - in Slovakia, pig iron and steel are produced in iron and steel integrated plant and by the EAF. Iron and steel integrated production is a complex one with many energy-related installations (coke ovens, heating plant, etc.). To avoid double counting of the primary and secondary fuels from iron and steel industry, the revised estimation was prepared in previous years in cooperation with the IPPU experts. The estimation includes and compares information from the iron and steel industry based on the EU ETS reports of the biggest iron and steel company in the Slovak Republic (U. S. Steel, s. r. o.). Methodology for emissions estimation was prepared by the specific model developed according to the national circumstances to ensure higher quality of estimation, avoiding double counting and properly allocated emissions in the Energy and IPPU sectors. Description of model is provided in details in the [Annex 4.2](#) (Methodology for carbon balance of iron and steel production).

## Emission Factors and NCVs

Detail description of the emission factors and NCVs used for estimation of emissions from fuel combustion is given in the [Chapter 3.2.6](#). Mainly country-specific or plant-specific emission factors are used in the category 1.A.2, although IPCC default emission factors are used for not key fuels. In the case of iron and steel integrated plant, all emission factors (NCVs and oxidation factors) are plant specific. Emission factors for anthracite, cooking coal, other bituminous coal and petroleum coke in the 1.A.2.a are also country specific (estimated as weighted average of sources allocated in this subcategory). The list of actually used EFs is presented in [Table 3.12](#).

**Table 3.12: Overview of country or plant specific CO<sub>2</sub> EFs in t/TJ in the category 1.A.2 in 2023**

1.A.2.a	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	70.07	Residual Fuel Oil	20.80	76.27
		Liquefied Petroleum Gases	17.22	63.14
Solid	156.41	Gas Coke	29.80	109.27
		Other Bituminous Coal	27.39	100.43
		Blast Furnace Gas	76.62	280.94
		Coke Oven Gas	11.44	41.95
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	110.69	Wood/Wood Waste	30.50	111.83
1.A.2.b	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	69.88	Gas/Diesel Oil	20.23	74.18
		Residual Fuel Oil	20.80	76.27
		Liquefied Petroleum Gases	17.22	63.14
Solid	98.88	Other Bituminous Coal	27.39	100.43
		Gas Coke	26.78	98.19
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	109.27	Wood/Wood Waste	30.50	111.83
1.A.2.c	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	63.15	Residual Fuel Oil	20.80	76.27
		Gas/Diesel Oil	20.23	74.18
		Liquefied Petroleum Gases	17.22	63.14
Solid	98.34	Anthracite	27.15	99.55
		Coking Coal	25.68	94.16
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	139.29	Wood/Wood Waste	30.50	111.83
		Other Primary Solid Biomass	27.30	100.10
		Other Biogas	14.90	54.63

		Biogenic waste	39.00	143.00
<b>1.A.2.d</b>	<b>WEIGHTED CO<sub>2</sub> EFs</b>	<b>FUEL TYPE</b>	<b>C EFs</b>	<b>CO<sub>2</sub> EFs</b>
Liquid	71.56	Residual Fuel Oil	20.80	76.27
		Liquefied Petroleum Gases	17.22	63.14
Solid	97.42	Other Bituminous Coal	27.39	100.43
		Lignite	26.78	98.19
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	98.69	Sulphite lyes (black liquor)	26.00	95.33
		Wood/Wood Waste	30.50	111.83
		Sludge Gas	14.90	54.63
		Other Primary Solid Biomass	27.30	100.10
<b>1.A.2.e</b>	<b>WEIGHTED CO<sub>2</sub> EFs</b>	<b>FUEL TYPE</b>	<b>C EFs</b>	<b>CO<sub>2</sub> EFs</b>
Liquid	71.26	Liquefied Petroleum Gases	17.22	63.14
		Gas/Diesel Oil	20.23	74.18
		Residual Fuel Oil	20.80	76.27
Solid	109.06	Anthracite	27.15	99.55
		Brown Coal Briquettes	26.61	97.57
		Gas Coke	26.78	98.19
Gaseous	56.15	Natural gas	15.32	56.18
Biomass	87.34	Other Primary Solid Biomass	27.30	100.10
		Sludge Gas	14.90	54.63
		Other Biogas	14.90	54.63
		Wood/Wood Waste	30.50	111.83
<b>1.A.2.f</b>	<b>WEIGHTED CO<sub>2</sub> EFs</b>	<b>FUEL TYPE</b>	<b>C EFs</b>	<b>CO<sub>2</sub> EFs</b>
Liquid	95.05	Residual Fuel Oil	20.80	76.27
		Petroleum Coke	26.19	96.03
		Liquefied Petroleum Gases	17.22	63.14
		Gas/Diesel Oil	20.23	74.18
Solid	102.07	Anthracite	27.15	99.55
		Other Bituminous Coal	27.39	100.43
		Lignite	26.78	98.19
		Gas Coke	26.78	98.19
Gaseous	56.18	Natural gas	15.32	56.18
Other	90.90	Municipal and Industrial Wastes	24.79	90.90
Biomass	91.23	Wood/Wood Waste	30.50	111.83
		Waste (biogenic)	24.79	90.90
<b>1.A.2.g</b>	<b>WEIGHTED CO<sub>2</sub> EFs</b>	<b>FUEL TYPE</b>	<b>C EFs</b>	<b>CO<sub>2</sub> EFs</b>
Liquid	65.68	Gas/Diesel Oil	20.23	74.18
		Liquefied Petroleum Gases	17.22	63.14
		Residual Fuel Oil	20.80	76.27
Solid	81.27	Blast Furnace Gas	76.62	280.94
		Coke oven Gas	11.44	41.95
		Lignite	26.78	98.19
		Other bituminous coal	27.39	100.43
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	111.83	Other primary solid biomass	27.30	100.1
		Wood/Wood waste	30.50	111.83

---

## Uncertainties and Time-series Consistency

Description of uncertainty is similar to the [Chapter 3.2.6](#) of this Document.

Time series is consistent in all aspects (methodological approach, country specific EFs and oxidation factor used, fuel characteristics, etc.) to the detailed level of disaggregation (on plant specific level).

### 3.2.8. Transport (CRT 1.A.3)

Transport has a very special position in the Energy sector, as it is not included in the EU ETS or other policies or measures, thus transport emissions are very difficult to regulate. The emissions balanced in the transport (1.A.3) include subcategories Domestic Aviation (1.A.3.a), Road transport (1.A.3.b), Railways (1.A.3.c), Domestic Navigation (1.A.3.d) and Pipeline transport (1.A.3.e.i). This Document uses the GWP<sub>100</sub> based on IPCC Assessment report 5. The difference between emission based on GWP<sub>100</sub> IPCC Assessment report 4 (AR4) and 5 (AR5) are shown in the previous NIR 2023.

As previously mentioned in reports, Slovakia has experienced a notable shift from public transport to individual passenger cars. Following the decrease in fuel consumption and emissions during the pandemic year of 2020, there was an overall increase across all categories of road transport, including passenger cars. However, in 2023, a change was observed. For the first time since 1990, without any external disturbances, there was a decrease in the transportation of passenger cars and all other types of road transport. Total aggregated GHG emissions in transport increased in 2023 against the base year by 13.49%, but decreased against the previous year by 0.55%. Road transport emissions rose by 67.13% in 2022 in comparison with the base year.

The emissions from road and non-road transport were calculated by using models, default methodologies and the consistent data series from 1990 – 2023 are presented in CRT Tables. Total GHG emissions in transport were 7 735.54 Gg of CO<sub>2</sub> eq. in 2023. The CO<sub>2</sub> emissions were 7 647.82 Gg, which represent 98.87% share on total transport emissions, the CH<sub>4</sub> emissions were 5.36 Gg of CO<sub>2</sub> eq. with the 0.07% share and N<sub>2</sub>O emissions were 82.35 Gg of CO<sub>2</sub> eq. with the 1.06% share on total transport GHG emissions.

Within transport, the share of road transport was 98.48%, pipeline transport 0.27%, railways 1.16%, domestic aviation represents 0.02% and domestic navigation 0.07% (in CO<sub>2</sub> eq.). Total energy consumption was 111 893.26 TJ of fuels in 2023. Among fuels, the most important are liquid fuels ([Figure 3.7](#)) and gaseous fuels. No solid fuels were used in transport category. Category “other fossil fuels” represents the fossil part of biomass fuels. The time series of GHG emissions are presented in [Table 3.13](#).

Figure 3.7: The share of fuels on different categories within transport in 2023




Table 3.13: Fuel consumption and GHG emissions in transport by subcategories in particular years

YEAR	1.A.3.a DOMESTIC AVIATION				1.A.3.b ROAD TRANSPORT			
	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg/year			TJ	Gg/year		
1990	51.48	3.74	0.000070	0.000102	61 027.37	4 503.02	1.1654	0.1895
1995	36.57	2.66	0.000050	0.000072	54 601.91	4 033.64	1.2323	0.1681
2000	36.50	2.65	0.000050	0.000072	56 107.97	4 077.90	0.9199	0.1466
2005	107.14	7.79	0.000158	0.000212	84 295.59	6 159.74	1.0570	0.1932
2010	70.59	5.13	0.000095	0.000140	92 325.43	6 435.39	0.7401	0.1634
2015	50.31	3.66	0.000069	0.000099	99 465.29	6 930.98	0.3173	0.2492
2016	49.00	3.56	0.000065	0.000097	102 045.72	7 063.71	0.2251	0.2510
2017	46.96	3.42	0.000066	0.000093	104 095.16	7 182.75	0.2765	0.2669
2018	39.21	2.85	0.000054	0.000078	106 591.21	7 338.02	0.1934	0.2699
2019	25.15	1.83	0.000040	0.000050	109 196.65	7 550.01	0.1842	0.2750
2020	12.17	0.88	0.000017	0.000024	98 359.45	6 743.80	0.1509	0.2209
2021	17.82	1.29	0.000025	0.000035	105 501.30	7 227.29	0.1632	0.2717
2022	20.47	1.48	0.000027	0.000040	111 491.04	7 583.93	0.1821	0.2834
2023	21.52	1.56	0.000031	0.000042	110 255.77	7 538.72	0.1856	0.2795

YEAR	1.A.3.c RAILWAYS				1.A.3.d DOMESTIC NAVIGATION			
	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg/year			TJ	Gg/year		
1990	5 024.14	372.29	0.0209	0.1437	0.30	0.02	0.0000021	0.0000006
1995	2 693.37	199.58	0.0112	0.0770	0.27	0.02	0.0000019	0.0000005
2000	2 080.68	154.18	0.0086	0.0595	0.33	0.02	0.0000023	0.0000007
2005	1 411.21	104.57	0.0059	0.0404	0.47	0.03	0.0000033	0.0000009
2010	1 162.77	82.32	0.0048	0.0333	4.49	0.33	0.0000339	0.0000090
2015	1 220.28	84.33	0.0051	0.0349	83.94	6.22	0.0005895	0.0001679
2016	1 250.91	86.53	0.0052	0.0358	64.24	4.76	0.0004522	0.0001285
2017	1 222.54	84.35	0.0051	0.0350	63.32	4.69	0.0004458	0.0001262
2018	1 197.06	82.93	0.0050	0.0342	34.53	2.56	0.0002446	0.0000691
2019	1 174.06	81.02	0.0049	0.0336	56.36	4.17	0.0003974	0.0001127
2020	1 052.53	72.53	0.0044	0.0301	72.25	5.35	0.0005058	0.0000144
2021	1 186.31	82.15	0.0049	0.0339	78.25	5.82	0.0005477	0.0001565

YEAR	1.A.3.c RAILWAYS				1.A.3.d DOMESTIC NAVIGATION			
	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg/year			TJ	Gg/year		
2022	1 193.84	82.29	0.0050	0.0341	71.45	5.29	0.0005002	0.0001429
2023	1 171.64	81.29	0.0049	0.0310	71.73	5.32	0.0005021	0.0001435

YEAR	1.A.3.e.i PIPELINE TRANSPORT			
	FUEL	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg/year		
1990	31 844.87	1 813.95	0.0318	0.0032
1995	20 644.81	1 154.10	0.0206	0.0021
2000	25 523.75	1 404.81	0.0255	0.0026
2005	24 168.60	1 327.92	0.0242	0.0024
2010	1 4961.55	824.47	0.0150	0.0015
2015	3 309.18	184.40	0.0033	0.0003
2016	5 351.33	298.41	0.0054	0.0005
2017	5 730.92	319.11	0.0057	0.0006
2018	5 315.65	295.17	0.0053	0.0005
2019	7 141.84	398.28	0.0071	0.0007
2020	3 009.14	167.83	0.0030	0.0003
2021	2 160.94	120.93	0.0022	0.0002
2022	287.05	16.12	0.0003	0.00003
2023	372.60	20.93	0.0004	0.00004

To estimate CO<sub>2</sub> emissions, country-specific (CS) data were used. The data used to calculate country-specific CO<sub>2</sub> emission factors (EF) included the fuel Net Calorific Value (NCV) and the H:C and O:C ratio for specific fuels. This EF was subsequently applied to every transport sector where these fuels were used. The calculated CS EF is summarized in [Table 3.14](#).

**Table 3.14:** CO<sub>2</sub> country-specific emission factors for selected fuels for the year 2023

FUEL	PETROL	DIESEL OIL	BIO-ETHANOL	BIO-DIESEL
	t/TJ	t/TJ	t/TJ	t/TJ
Emission factor	69.786	74.197	70.605	74.636

**Domestic aviation (CRT 1.A.3.a)** - The inventory evaluation of GHG emissions in domestic aviation was performed for all GHGs, precursors and air pollutants. In the absence of national data on the exact numbers of domestic LTO cycles for the years 1990 – 2004 (only total national + international numbers of LTO cycles are available), summary information from the EUROCONTROL database was used. According to the recommendations of the ERT during previous reviews and following the IPCC 2006 GL, the emissions estimation was based on the fuel sold to national and international civil flights (tier 1 approach as it is not a key category for the Slovak Republic) for the years 1990 – 2004. The Slovak airports, except for the airport in Žilina, where exercises with light aircrafts of the Žilina University predominate, are managed by themselves as separate company. Other smaller civil airports (Nitra, Prievidza, Ružomberok and Lučenec) are operated by aero clubs with predomination of sport flights. Described approach is maintained for a time series 1990 – 2004. For the time series 2005 – 2023, EUROCONTROL data on the number of flights, fuels consumption and share of domestic and international flights was used.

The fuels consumption in domestic aviation decreased in 2023 compared to the base year 1990 by 58.3%. The total jet kerosene consumption was 19.21 TJ and the consumption of aviation gasoline (AvGas) was 2.31 TJ allocated in domestic aviation in 2023 ([Table 3.15](#)). Total GHG emissions from domestic aviation were 1.56 Gg of CO<sub>2</sub> eq. in 2023. There was a visible increase of emissions in years

2002 – 2008 ([Figure 3.10](#)). In 2002, air transport was positively affected by the entry of low-cost companies to the Slovak market, like SkyEurope Airlines, Seagle Air and Danube Wings, but this also caused an increase in emissions. The time series is influenced by the fact, that the Slovak Republic has no official national airlines since the Slovak Airlines are out of business since 2007, SkyEurope since 2009 and close distance of other big international airports in Vienna and Budapest.

**Table 3.15: The fuels consumption and GHG emissions for national flights in particular years**

YEAR	AVIATION GASOLINE				JET KEROSENE			
	CONSUMPTION	EMISSIONS			CONSUMPTION	EMISSIONS		
	TJ	t CO <sub>2</sub>	t CH <sub>4</sub>	t N <sub>2</sub> O	TJ	t CO <sub>2</sub>	t CH <sub>4</sub>	t N <sub>2</sub> O
1990	3.35	236.99	0.002	0.007	48.13	3 501.22	0.068	0.095
1995	2.22	156.82	0.001	0.004	34.36	2 499.39	0.049	0.068
2000	2.56	180.67	0.002	0.005	33.94	2 469.37	0.048	0.067
2005	0.95	67.23	0.001	0.002	106.19	7 725.42	0.158	0.210
2010	1.85	130.64	0.001	0.004	68.75	5 001.21	0.094	0.136
2015	2.11	149.27	0.001	0.004	48.20	3 506.73	0.068	0.095
2016	1.68	116.63	0.001	0.003	47.32	3 442.59	0.064	0.094
2017	1.97	138.78	0.001	0.004	44.99	3 281.18	0.065	0.089
2018	2.32	163.68	0.001	0.005	36.89	2 690.19	0.053	0.073
2019	1.99	140.17	0.001	0.004	23.16	1 689.13	0.039	0.046
2020	1.59	110.14	0.001	0.003	10.58	769.74	0.016	0.021
2021	1.50	104.19	0.001	0.003	16.31	1 186.75	0.025	0.032
2022	2.26	156.35	0.001	0.004	18.22	1 325.12	0.026	0.036
2023	2.31	160.23	0.001	0.004	19.21	1 397.14	0.030	0.038

**Road transport (CRT 1.A.3.b)** - Short distance passenger transport is an important part of road transport. It is the most exploited type of transport in the Slovak Republic due to a high density and quality of road network and interconnection of all municipalities. In recent years, road transport has expanded significantly in the transport of goods and persons. In 2023, the transport network included 544 km of highways, 320 km of motorways and 3 335 km of the category 1<sup>st</sup> class roads. Total road network represented 18 143 km of roads in the Slovak Republic<sup>5</sup> in 2023. Road transport is the most important and key category with the highest share of emissions and continually increasing trend in fuels consumption within transport. This increase was however interrupted by the COVID pandemic and Slovakia observed a temporary major decrease of fuel consumption and GHG emissions in 2020. After that it is again observed rise in fuel consumption and emissions in 2021 and 2022, with a decrease in 2023. Total aggregated emissions from road transport reached 7 617.99 Gg of CO<sub>2</sub> eq. in 2023. The decrease in emissions compared to 2022 is 0.60%, and increase compared to the base year is 66.12%. The major share of emissions belongs to heavy duty vehicles and passenger cars ([Table 3.13](#)). Total blended CO<sub>2</sub> emissions were 8 035.51 Gg in 2023. These blended emissions include also emissions from lube oil from two-stroke petrol passenger cars. After separation of biomass content, the final CO<sub>2</sub> balance for fossil part of fuels was 7 538.72 Gg. Biomass content in fuels increased in 2018 compared to the previous year mainly to introduction of E10 petrol and subsequently decrease because of COVID-19, these emissions represent 496.79 Gg of bio-CO<sub>2</sub> in 2023. The most of the emissions come from the city traffic ([Table 3.17](#)).

<sup>5</sup> [Slovak Road Database 2023](#)

**Table 3.16: Overview of total GHG emissions according to the type of vehicles in 2023**

CATEGORY OF ROAD VEHICLE	EMISSIONS			CATEGORY OF ROAD VEHICLE	EMISSIONS		
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	t/year	kg/year			t/year	kg/year	
Passenger Cars	4 788 758	121 320	122 777	Heavy Duty Trucks	1 923 085	28 617	115 834
Petrol Mini	2 942	182	28	Petrol >3,5 t	24	6	0
Petrol Small	884 003	52 051	10 167	Diesel Rigid <=7,5 t	125 804	2 932	5 521
Petrol Medium	657 151	29 451	7 379	Diesel Rigid 7,5 - 12 t	166 483	2 385	5 359
Petrol Large-SUV-Executive	101 676	3 604	809	Diesel Rigid 12 - 14 t	39 029	493	2 146
2-Stroke	41	14	0	Diesel Rigid 14 - 20 t	55 722	1 484	2 534
Petrol Hybrid Mini	34	3	0	Diesel Rigid 20 - 26 t	5 544	246	181
Petrol Hybrid Small	10 763	875	89	Diesel Rigid 26 - 28 t	560	13	25
Petrol Hybrid Medium	73 608	5 076	541	Diesel Rigid 28 - 32 t	637	28	30
Petrol Hybrid Large-SUV-Executive	40 010	1 643	174	Diesel Rigid >32 t	689	19	28
Petrol PHEV Small	1 739	141	14	Diesel Articulated 14 - 20 t	1 523 231	19 145	99 750
Petrol PHEV Medium	5 166	356	38	Diesel Articulated 20 - 28 t	1 617	174	58
Petrol PHEV Large-SUV-Executive	5 993	244	26	CNG Articulated < 40 t	220	52	12
Diesel Mini	237	2	13	LNG Articulated < 40 t	3 527	1 641	191
Diesel Small	35 763	307	1 360	Buses	311 115	16 662	13 512
Diesel Medium	2 196 067	12 483	83 018	Urban Buses Midi <=15 t	24 673	185	1 458
Diesel Large-SUV-Executive	603 632	2 760	16 775	Urban Buses Standard 15 - 18 t	15 810	73	712
Diesel PHEV Large-SUV-Executive	208	0	8	Urban Buses Articulated >18 t	463	2	16
LPG Bifuel Mini	32	2	1	Coaches Standard <=18 t	250 179	2 083	9 641
LPG Bifuel Small	85 772	6 132	1 062	Coaches Articulated >18 t	6 980	29	245
LPG Bifuel Medium	64 696	4 426	1 003	Diesel Hybrid	478	2	27
LPG Bifuel Large-SUV-Executive	15 299	976	241	Urban CNG Buses	12 531	14 288	1413
CNG Bifuel Mini	42	2	0	L-Category	24 285	11 157	433
CNG Bifuel Small	2 444	402	20	Mopeds 2-stroke <50 cm <sup>3</sup>	27	48	0
CNG Bifuel Medium	1 344	173	9	Mopeds 4-stroke <50 cm <sup>3</sup>	1 032	645	19
CNG Bifuel Large-SUV-Executive	95	16	1	Motorcycles 2-stroke >50 cm <sup>3</sup>	90	135	2
Light Commercial Vehicles	988 263	5 477	24 904	Motorcycles 4-stroke <250 cm <sup>3</sup>	2 602	2 256	90
Petrol N1-I	30 075	1 709	321	Motorcycles 4-stroke 250 - 750 cm <sup>3</sup>	9 127	5 143	147
Petrol N1-II	17 773	553	217	Motorcycles 4-stroke >750 cm <sup>3</sup>	11 395	5 143	147
Petrol N1-III	3 790	124	56	Quad & ATVs	3	2 929	174
Diesel N1-I	25 265	285	973	Micro-car	8	1	0
Diesel N1-II	253 488	1 259	6 424	<b>Total</b>	<b>8 035 507</b>	<b>183 234</b>	<b>277 460</b>
Diesel N1-III	657 872	1 548	16 913				



**Table 3.17: Results from COPERT model in distribution for driving mode (CO<sub>2</sub> emissions are from blended fuels with bio-component) in 2023**

TRAFFIC	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	t/year		
Urban	3 564 389	107.31	117.04
Rural	3 169 603	55.72	122.03
Highway	1 301 514	20.21	38.38
TOTAL	8 035 507	183.23	277.46

**Railways (CRT 1.A.3.c)** - Railways are the second largest source of emissions in transport, despite the decreasing character of this transport mode. Railways and rail transport are slowly modernised in Slovakia with the support of the EU funds. Improved quality and ecology of rail transport and the increase in passengers' number are the results of this modernisation. Modernisation of rail infrastructure results in an increase of operational speed to 160 km/h and increase of safety. According to the Annual Report of Slovak Railways<sup>6</sup> in 2023, the length of managed railways was 3 630 km of which the length of electric railways was 1 585 km. Total emissions from railways transport reached 89.65 Gg of CO<sub>2</sub> eq. in 2023 and they decreased by 2.01% compared to 2022 (**Table 3.18**) and decreased by 78.19% compared to the base year. The decrease of fuels consumption compared to the base year was caused by the improvements of technical parameters. Rising of passenger transport on railways, partly caused by governmental measure<sup>7</sup> led to emissions increase, while cargo is fluctuating without visible trend.

**Table 3.18: Overview of fuels consumption and GHG emissions in railways in particular years**

YEAR	TOTAL CONSUMPTION	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg/year		
1990	5 024.137	372.289	0.021	0.144
1995	2 693.369	199.579	0.011	0.077
2000	2 080.683	154.179	0.009	0.060
2005	1 411.206	104.570	0.006	0.040
2010	1 162.771	82.320	0.005	0.033
2015	1 220.277	84.332	0.005	0.035
2016	1 250.911	86.533	0.005	0.036
2017	1 222.536	84.352	0.005	0.035
2018	1 197.061	82.933	0.005	0.034
2019	1 174.056	81.024	0.005	0.034
2020	1 052.530	72.532	0.004	0.030
2021	1 186.310	82.150	0.005	0.034
2022	1 193.836	82.294	0.005	0.034
2023	1 171.636	81.291	0.005	0.031

**Domestic navigation (CRT 1.A.3.d)** - The major share of emissions from shipping in Slovakia are realized as transit on Danube River. Due to international character of this river, emissions are included in the subcategory 1.D.1.b - Memo Items/International Bunkers/International Navigations (**Chapter 3.8**). Based on the information from the State Navigation Administration (the SNA), there are several movements realized between the Bratislava, Komárno and Štúrovo ports on the Slovak territory (national transport). Usually ships do not stop their operation on the Slovak territory, but the transit continues to Austria or Hungary. However, the part of GHG emissions from the movements between the ports on

<sup>6</sup> [Annual Report of Slovak Railway 2023, p. 18](#)

<sup>7</sup> Since 2013, social measure was introduced – free railways for students and retired on lower categories of trains.

Slovak Territory is included in the national emissions inventory. Detailed information was based on statistics made by the SNA and the Slovak Shipping and Ports Company. The share of “national fuel consumption” is available since 2005. Inland shipping transport on small lakes for tourist purposes was not included in the 2020 report as those were not operating during the COVID-19 pandemic. In 2021 only a few restored their activity.

Total aggregated emissions from inland shipping excluding international navigations (on Danube River) reached 5.37 Gg of CO<sub>2</sub> eq. in 2023. After a decrease in 2018, an increase is observed from 2019 despite of COVID-19 pandemic and no tourist tours on lakes ([Table 3.19](#)).

**Table 3.19: Overview of fuels consumption and GHG emissions in domestic navigation in particular years**

YEAR	TOTAL CONSUMPTION	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg/year		
1990	0.303	0.022	0.000002	0.000001
1995	0.274	0.020	0.000002	0.000001
2000	0.328	0.024	0.000002	0.000001
2005	0.468	0.035	0.000003	0.000001
2010	4.488	0.327	0.000031	0.000009
2015	83.942	6.215	0.000587	0.000168
2016	64.239	4.757	0.000452	0.000128
2017	63.324	4.689	0.000445	0.000126
2018	34.530	2.556	0.000244	0.000069
2019	56.361	4.172	0.000397	0.000113
2020	72.251	5.350	0.000506	0.000145
2021	78.250	5.823	0.000548	0.000157
2022	71.451	5.293	0.000500	0.000143
2023	71.710	5.321	0.000502	0.000143

**Pipeline transport (CRT 1.A.3.e.i)** – Total fuels in 1.A.3.e.i expressed in energy units represented 372.60 TJ and total GHG emissions represented 20.95 Gg of CO<sub>2</sub> eq. in 2023. The share of this category on total transport emissions significantly decreased to 0.27% in 2023. This significant decrease is caused by war in Ukraine and lower transport of natural gas from east to west. The fuel consumption and GHG emissions are shown in [Table 3.13](#).

### Methodological Issues

**Domestic aviation (1.A.3.a)** – Domestic Aviation is not a key category. The airport traffic in Slovakia is determined only by the origin of airlines. It means, that there is no direct information about the number of domestic and international flights in statistics. Tier 1 approach for emission estimation in domestic aviation, both for aviation gasoline and jet kerosene was used for time series 1990 – 2004. Tier 1 approach is based on fuel sold on the airports. For this period, only total number of LTO cycles is known, therefore average disaggregation of activities between national and international aviation was judged. The expert judgment of share of national and international aviation activities for the period 1990 – 2004 was improved based on the known real numbers for time series 2005 – 2018 based on tier 3. Then the time series 1990 – 2004 was revised using constant share for national and international flights. Real share of national and international activities for the period 2005 – 2023 was taken from the EUROCONTROL ([Table 3.20](#)).

**Table 3.20: The share of fuel consumption in domestic aviation and international bunkers for the period 1990 – 2004**

FUELS	DOMESTIC AVIATION		INTERNATIONAL BUNKERS	
	PREVIOUS ESTIMATE	REVISED ESTIMATE	PREVIOUS ESTIMATE	REVISED ESTIMATE
AVIATION GASOLINE	90%	30%	10%	70%
JET KEROSENE	10%	5%	90%	95%

The implied emission factors applied in previous submissions for the years 1990 – 2004 were not in the IPCC range, therefore the new EFs for all GHG gases were calculated as average from the available EUROCONTROL data for years 2005 – 2018 and used from 2019 onwards for the years 1990 – 2004. These average EFs are EUROCONTROL based and were used since 2004 back to the base year to maintain consistency in the time-series. Activity data for the years 1990 – 1993 are not available and were estimated as expert judgment according to real LTO cycles in this period. For the period 1994 – 2004, activity data were directly provided by the airports on annual basis. Due to the time series consistency, the net calorific values from the EUROCONTROL data were used to convert obtained activity data.

From the year 2005 onwards, Slovakia decided to use directly the EUROCONTROL data. The decision was based on analysis of the national data and data obtained from the EUROCONTROL. Results showed that EUROCONTROL data are more consistent and accurate in line with the QA/QC rules. EUROCONTROL data used tier 3 applying the Advanced Emissions Model (AEM).

Following data were taken from the EUROCONTROL ([Tables 3.21](#) and [3.22](#)):

- fuel consumption of aviation gasoline for domestic flights;
- fuel consumption of aviation gasoline for international flights;
- fuel consumption of jet kerosene for domestic flights;
- fuel consumption of jet kerosene for international flights;
- CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O emissions for all subcategories;
- NCVs calculated from fuel consumption.

**Table 3.21: Average EFs for the GHG emissions used in domestic civil aviation according to tier 1 based on fuel consumption**

PARAMETER	EMISSIONS FACTORS	
	INTERNATIONAL FLIGHTS	NATIONAL FLIGHTS
Emissions	Jet kerosene	
	kg/TJ of fuel	
CO <sub>2</sub>	72 748	72 748
CH <sub>4</sub>	0.707	1.343
N <sub>2</sub> O	1.977	1.977
Emissions	Aviation gasoline	
CO <sub>2</sub>	6 959	6 959
CH <sub>4</sub>	0.541	0.572
N <sub>2</sub> O	1.953	1.953

**Table 3.22: Average NCVs for the GHG emissions used in domestic civil aviation according to tier 1 based on fuel consumption**

NCVs		
Aviation Gasoline	TJ/Gg	44.00
Jet Kerosene	TJ/Gg	43.30

**Road transport (1.A.3.b)** – COPERT model 5 (v.5.8) was used for estimation of road transport emissions. The model distinguishes vehicle categories and emission factors reflecting the recent development and research. These data are not available before 2000. The methodology is often referred to the name of program (methodology “COPERT”). The model is based on the fuel approach, what is used for the CO<sub>2</sub> emissions estimation (tier 2). The fuel consumption and other variables such as H/C and O/C ratio and carbon content in fuels is used in this approach. According to the previous ERT recommendation, the country specific H/C ratio and NCVs were used in model calculation. Slovakia is analysing composition of fuels sold by the majority of companies on the market, representing 3 different refineries on regular basis. Delivering actual and most recent data on fuels’ composition is crucial for correct country-specific EFs estimation. The H/C and O/C ratio of the fuels was analysed by the Research Institute for Crude Oil and Hydrocarbon Gases (VÚRUP) in 2022 ([Tables 3.23](#) and [3.24](#)). According to measured data and previous information provided by the Slovnaft refinery, the H/C ratio rose between 2015 and 2017 only by 0.26%. The NCVs of the fuels were obtained from the Statistical Office of the Slovak Republic and are shown in [Table 3.25](#) for the years 1990 – 2023.

**Table 3.23: Results of the H/C ratio analyses of fuel types and lube oil in 2023**

FUEL	PETROL	DIESEL OIL	LPG	CNG	BIO-ETHANOL	BIO-DIESEL	LUBE OIL
H/C Ratio	1.767	1.946	2.589	3.900	3.000	1.857	1.900

**Table 3.24: Results of the O/C analyses of fuel types and lube oil in 2023**

FUEL	PETROL	DIESEL OIL	LPG	CNG	BIO-ETHANOL	BIO-DIESEL	LUBE OIL
O/C Ratio	0.000	0.005	NA	NA	0.500	0.110	NA

NA=oxygen is not present

**Table 3.25: Net calorific values (NCVs) for the fuel type obtained by the ŠÚ SR for particular years**

YEAR	PETROL BLENDED	DIESEL OIL BLENDED	LPG	CNG	BIO-ETHANOL	ETBE	ESTERS
	TJ/Gg						
1990	43.206	42.511	NO	NO	NO	NO	NO
1995	43.388	42.076	46.000	NO	NO	NO	NO
2000	43.316	42.588	46.000	48.814	NO	NO	NO
2005	43.800	42.208	46.000	48.767	NO	NO	NO
2010	43.728	42.218	46.000	48.948	27.000	36.000	37.000
2011	43.780	42.206	46.000	48.923	27.000	36.000	37.000
2012	43.740	42.206	46.000	48.802	27.000	36.000	37.000
2013	43.952	42.043	46.000	48.753	27.000	36.000	37.800
2014	43.905	42.043	46.000	48.597	27.000	36.000	38.450
2015	43.909	42.143	46.000	48.760	27.000	36.000	39.265
2016	43.908	42.136	46.000	48.800	27.000	36.000	39.486
2017	43.899	42.127	46.000	48.800	27.000	36.200	39.699
2018	43.774	42.695	46.564	48.000	28.800	36.000	37.300
2019	43.934	42.600	46.000	48.800	27.000	36.000	39.867
2020	43.932	42.086	46.000	48.780	27.000	36.000	39.807
2021	43.928	42.087	46.000	48.070	27.000	36.000	39.646
2022	43.924	42.108	46.000	48.004	27.336	36.000	39.987
2023	43.919	42.072	46.000	48.004	27.060	36.000	37.471

Statistically recorded fuel consumption and fuel consumption calculated through COPERT 5 model are equal, except of fossil petrol. There is a statistically insignificant difference on the level up to 2%. This is caused by highly complicated calculation and due to drastically shorten the time needed for

calculation. The new version added new vehicle categories for the CH<sub>4</sub> and N<sub>2</sub>O emissions estimation, with the disaggregation into 5 basic categories and 379 subcategories. Further disaggregation was applied according to the operation of road vehicles in the urban, rural and highway driving mode. In COPERT 5, buses were divided into 2 subcategories (urban and coaches) and seven weight categories. Heavy-duty vehicles are divided into 2 basic categories (rigid and articulated). Rigid vehicles are further divided by weight into 8 and articulated into six subcategories. EMEP/EEA methodology used technical parameters of different vehicle types and country-specific characteristics, such as the composition of car fleet, the age, operation and fuels or climate conditions.

Model estimates emissions from the following input data:

- total fuel consumption,
- composition of vehicle fleet,
- driving mode,
- driving speed,
- emission factors,
- annual mileage.

Information about the vehicle fleet is based on database IS EVO (Information System for Vehicle Evidence) operated by the Police Presidium of the Slovak Republic.

The EFs values for CH<sub>4</sub> and N<sub>2</sub>O in COPERT 5 model are defined separately for the different types of fuels, types of vehicles, different technological level of vehicles, driving mode and season as these emissions are depended on ambient and vehicle temperature. In case of CH<sub>4</sub> emissions, the balance is based on the average speed and drive mode for certain vehicles' group. The emission factors for pollutants such as CO<sub>2</sub>, SO<sub>2</sub>, N<sub>2</sub>O, NH<sub>3</sub>, PM and partially also CH<sub>4</sub> can be obtained by the simple formula of driving mode and consumed fuel. Emission factors are then calculated automatically by the model based on the input parameters such as the average speed, the quality of fuels, the age of vehicles, the weight of vehicles and the volume of cylinders.

Accurate and actual data on distance-based values and parameter values are necessary to run the COPERT 5 model (**Table 3.26**). Therefore, new input data on mileages was requested from the Technical Inspection -PTI (odometers) and the IS EVO (from the Police Department). As the unique key for binding data from these two registries, VIN number (Vehicle Identification Number) was used. Using MS Access, the average annual mileages were calculated. Further data, needed for calculation were: the first registration of vehicle, vehicle type, engine volume, weight, emission category and data from odometer. At least that many years as are between two technical controls were needed.

The average annual mileages including consistency with fuel consumption were also used for identifying distribution of vehicles to their appropriate COPERT category. The Traffic Census of Slovakia conducted in every five years (2000, 2005, 2010 and 2015<sup>8</sup>) was the main source for activity data such as intensity on urban, rural and highways.

**Table 3.26: Overview of input data used in the COPERT 5 model in 2023**

CATEGORY OF ROAD VEHICLE	ACTIVITY DATA		CATEGORY OF ROAD VEHICLE	ACTIVITY DATA	
	No.	km/veh.		No.	km/veh.
Passenger Cars	2 464 256	12 244	Diesel N1-III	148 890	15 444
Petrol Mini	7014	4 868	Battery electric N1-I	14	27 284
Petrol Small	785 499	4 803	Battery electric N1-II	150	15 045
Petrol Medium	388 454	5 729	Battery electric N1-III	362	16524

<sup>8</sup> Data were published in 2016

CATEGORY OF ROAD VEHICLE	ACTIVITY DATA		CATEGORY OF ROAD VEHICLE	ACTIVITY DATA	
	No.	km/veh.		No.	km/veh.
Petrol Large-SUV-Executive	47 536	5 968	Heavy Duty Trucks	69 317	22 513
Petrol 2-Stroke	154	1 163	Petrol >3,5 t	110	471
Petrol Hybrid Mini	35	6 478	Diesel Rigid <=7,5 t	21 519	18 232
Petrol Hybrid Small	10 193	9 972	Diesel Rigid 7,5 - 12 t	12 277	24 043
Petrol Hybrid Medium	40 203	12 363	Diesel Rigid 12 - 14 t	3 328	16 465
Petrol Hybrid Large-SUV-Executive	11 351	13 975	Diesel Rigid 14 - 20 t	4466	15 430
Petrol PHEV Small	1396	15 405	Diesel Rigid 20 - 26 t	1123	7 567
Petrol PHEV Medium	3 490	15 409	Diesel Rigid 26 - 28 t	47	20 342
Petrol PHEV Large-SUV-Executive	1900	18 043	Diesel Rigid 28 - 32 t	190	13 706
Diesel Mini	389	5 503	Diesel Rigid >32 t	171	3 942
Diesel Small	24 549	9 991	Diesel Articulated 14 - 20 t	25989	55 321
Diesel Medium	884 091	15 175	Diesel Articulated 50 - 60 t	17	72 972
Diesel Large-SUV-Executive	197 635	13 483	CNG Articulated < 40 t	40	6 702
Diesel PHEV Large-SUV-Executive	68	23 406	LNG Articulated < 40 t	40	110 954
LPG Bifuel Mini	19	10 822	Buses	6 552	44 934
LPG Bifuel Small	24 844	21 625	Diesel Urban Buses Midi <=15 t	620	26 585
LPG Bifuel Medium	19 373	19 271	Diesel Urban Buses Standard 15 - 18 t	236	57 890
LPG Bifuel Large-SUV-Executive	4 939	19 372	Diesel Urban Buses Articulated >18 t	13	30 248
CNG Bifuel Mini	29	10 316	Diesel Coaches Standard <=18 t	5 340	30 672
CNG Bifuel Small	874	14 882	Diesel Coaches Articulated >18 t	37	107 099
CNG Bifuel Medium	484	18 949	Urban Buses Diesel Hybrid	16	52 400
CNG Bifuel Large-SUV-Executive	55	14 262	Urban CNG Buses	247	44 990
Battery electric Mini	452	7 500	Battery electric -	43	25 394
Battery electric Small	1 303	11 650	L-Category	173 054	959
Battery electric Medium	2 902	11 624	Petrol Mopeds 2-stroke <50 cm <sup>3</sup>	1 347	692
Battery electric Large-SUV-Executive	5 025	20 576	Petrol Mopeds 4-stroke <50 cm <sup>3</sup>	27309	672
Light Commercial Vehicles	276 982	11 929	Petrol Motorcycles 2-stroke >50 cm <sup>3</sup>	2819	691
Petrol N1-I	23 909	18 949	Petrol Motorcycles 4-stroke <250 cm <sup>3</sup>	49681	824
Petrol N1-II	9 239	14 262	Petrol Motorcycles 4-stroke 250 - 750 cm <sup>3</sup>	50418	1 308
Petrol N1-III	2 141	12 838	Petrol Motorcycles 4-stroke >750 cm <sup>3</sup>	41353	1 840
Diesel N1-I	16 910	12 209	Petrol Quad & ATVs	59	332
Diesel N1-II	75 367	14 405	Micro-car	68	1 257

Regarding non-CO<sub>2</sub> emissions, the values used for setting and calculating the emission factors and the corresponding emissions in the COPERT model were verified and discussed in the previous years. The results of a comparative assessment for CH<sub>4</sub> and N<sub>2</sub>O emissions showed, that the emissions inventory of Slovakia is comparable with other European countries and therefore the use of emission factors in the COPERT model are fully in agreement with the Middle European (Slovakia) national circumstances. The IEFs used in COPERT model are regularly updated and verified ([Table 3.27](#)) in a more advance versions of model. Methane IEFs are gradually decreasing for all vehicle categories, including light-duty vehicles owing to changes in the vehicle fleet. Newer vehicles are emitting fewer hydrocarbon pollutants, to which oxidation catalysts contribute. Methane behaves just like other hydrocarbons, so it declines,

resulting in a decline in total emissions and also in IEFs. The emissions of N<sub>2</sub>O are slowly increasing for light-duty vehicles (diesel) owing to NO<sub>x</sub> reduction devices (SCR and EGS/DPF system).

**Table 3.27: Overview of CH<sub>4</sub> and N<sub>2</sub>O IEFs for the road vehicle categories in 2023**

CATEGORY OF ROAD VEHICLE	EMISSION FACTORS		CATEGORY OF ROAD VEHICLE	EMISSION FACTORS	
	CH <sub>4</sub>	N <sub>2</sub> O		CH <sub>4</sub>	N <sub>2</sub> O
	mg/km			mg/km	
Passenger Cars	4.53	4.58	Diesel N1-III	0.61	6.64
Petrol Mini	9.32	1.41	Heavy Duty Trucks	7.83	31.70
Petrol Small	10.48	2.05	>3,5 t	109.90	6.00
Petrol Medium	9.08	2.27	Rigid <=7,5 t	8.07	15.21
Petrol Large-SUV-Executive	9.67	2.17	Rigid 7,5 - 12 t	7.54	16.95
2-Stroke	79.58	0.00	Rigid 12 - 14 t	7.07	30.75
Petrol Hybrid Mini	9.32	1.00	Rigid 14 - 20 t	17.17	29.33
Petrol Hybrid Small	9.32	0.95	Rigid 20 - 26 t	35.84	26.33
Petrol Hybrid Medium	9.32	0.99	Rigid 26 - 28 t	17.45	32.19
Petrol Hybrid Large-SUV-Executive	9.32	0.99	Rigid 28 - 32 t	38.39	40.98
Petrol PHEV Small	6.92	0.70	Rigid >32 t	23.35	34.56
Petrol PHEV Medium	6.92	0.73	Articulated 14 - 20 t	6.83	35.59
Petrol PHEV Large-SUV-Executive	6.92	0.74	Articulated 20 - 28 t	86.98	28.81
Diesel Mini	1.08	6.66	Articulated < 40 t	194.73	43.02
Diesel Small	1.42	6.29	Articulated < 40 t	369.73	43.02
Diesel Medium	0.94	6.27	Buses	46.36	37.60
Diesel Large-SUV-Executive	1.03	6.27	Urban Buses Midi <=15 t	5.17	40.78
Diesel PHEV Large-SUV-Executive	0.06	4.93	Urban Buses Standard 15 - 18 t	4.24	41.47
LPG Bifuel Mini	10.61	2.95	Urban Buses Articulated >18 t	4.17	41.50
LPG Bifuel Small	11.39	1.97	Coaches Standard <=18 t	7.35	34.03
LPG Bifuel Medium	12.23	2.77	Coaches Articulated >18 t	4.21	35.55
LPG Bifuel Large-SUV-Executive	12.57	3.11	Diesel Hybrid	2.85	31.85
CNG Bifuel Mini	7.21	0.98	Urban CNG Buses	1021.24	101.00
CNG Bifuel Small	27.87	1.40	L-Category	49.42	1.92
CNG Bifuel Medium	26.62	1.36	Mopeds 2-stroke <50 cm <sup>3</sup>	114.10	1.00
CNG Bifuel Large-SUV-Executive	26.05	1.35	Mopeds 4-stroke <50 cm <sup>3</sup>	34.86	1.00
Light Commercial Vehicles	1.40	6.34	Motorcycles 2-stroke >50 cm <sup>3</sup>	130.03	2.00
Petrol N1-I	11.15	2.09	Motorcycles 4-stroke <250 cm <sup>3</sup>	49.92	2.00
Petrol N1-II	8.08	3.17	Motorcycles 4-stroke 250 - 750 cm <sup>3</sup>	69.76	2.00
Petrol N1-III	7.98	3.60	Motorcycles 4-stroke >750 cm <sup>3</sup>	33.75	2.00
Diesel N1-I	1.85	6.29	Quad & ATVs	53.35	2.00
Diesel N1-II	1.29	6.58	Micro-car	5.12	0.89


Input parameters for CNG buses are known only since 2000. Before the year 2000, CNG consumption in transport was negligible. The consumption of CNG as fuel can be used neither for a diesel engine nor for a petrol engine without modifications. The CNG buses have completely different combustion and after-treatment technology despite using the same fuel as CNG passenger cars. Hence, their emissions performance may vary significantly. Therefore, CNG buses also need to fulfil specific emissions standards (Euro II, Euro III, etc.). Due to the low NO<sub>x</sub> and PM performance compared to diesel oil, an

additional emissions standard has been set for CNG vehicles, known as the standard for Enhanced Environmental Vehicles (EEV). The emission limits imposed for EEV are even below Euro V and usually EEVs are benefited from taxation waivers and free entrance to low emissions zones. New stoichiometry buses are able to fulfil the EEV requirements, while older buses were usually registered as Euro II, Euro III, Euro IV or Euro V.

The statistical consumptions of petrol, diesel oil and biofuels were received from data reported under the Fuel Quality Directive art. 7a by SHMU and cross-checked according to data received from the Ministry of Economy (MH SR). According to the latest QA/QC these consumptions are the most accurate (Chapter 3.2.8). Data about LPG distribution and sale were obtained from the Slovak Association of Petrochemical Industry (SAPPO). CNG consumption were obtained from Financial Administration of the Slovak Republic (FR SR). All documents are available in Slovak language and they are official. Share of diesel oil represents 68.38%, followed by petrol with 23.52% share, then LPG (1.57%), CNG (0.29%), biomass (6.05%) and other fossil fuels (fossil part of biofuels) (0.18%) in 2023 (Figure 3.8).

The blending of biomass in liquid fuels was considered and the bio-emissions are calculated since 2007 (first year of using blended fuels in transport in Slovakia). Fuel quality is provided by the MH SR in terms of implementing Directive No 2009/29/EC and the Directive No 2009/30/EC on the replacement of fossil fuels with bio-component. The share of biomass in liquid fuels in transport was calculated as bio-component percentage. In ETBE as bio-component is considered only in 47% by mass in calculation of total bio-components in fuel. From the biomass (biodiesel) is also subtracted the 5.34% fossil methanol part and all emissions from the bio-parts of biofuels are reported as biomass emissions, and the fossil part is reported in its associated fossil fuel (ETBE – petrol; FAME – diesel). Fossil part of FAME was calculated as national average according to data from the report under Fuel Quality Directive Art. 7(a) (Table 3.28).

Figure 3.8: Share of fuels on total consumption in road transport in 2023



Requirements for the quality of motor fuels containing bio-component must be at the level of the specifications listed in the STN EN 228:2004 and STN EN 590:2004, respectively. The quality of blending in bio-liquid fuels must meet the requirements specified in the STN EN 14 214, STN EN 15 376.



**Table 3.28: Estimated activity data and share of biomass for the time series 2007 – 2023**

YEAR	PETROL		DIESEL OIL	
	BIOMASS SHARE (ENERGY)	BIOMASS CONSUMPTION	BIOMASS SHARE (ENERGY)	BIOMASS CONSUMPTION
	%	TJ	%	TJ
2007	2.30%	652.26	4.09%	2 677.29
2008	1.23%	358.17	4.77%	2 795.75
2009	2.58%	706.72	5.14%	3 090.30
2010	2.95%	779.13	5.28%	3 577.88
2011	2.97%	715.87	6.05%	3 741.68
2012	2.94%	710.56	5.79%	3 846.12
2013	3.21%	726.60	6.43%	4 107.36
2014	3.88%	859.33	5.65%	3 766.08
2015	3.33%	747.87	5.74%	4 342.97
2016	3.10%	725.62	6.68%	5 158.95
2017	4.06%	943.49	6.92%	5 464.18
2018	4.52%	1 018.32	6.97%	5 697.80
2019	4.46%	1 042.07	6.45%	5 371.36
2020	6.20%	1 390.40	7.27%	5 401.90
2021	6.20%	1 419.47	6.96%	5 617.89
2022	6.04%	1 645.29	6.95%	5 779.55
2023	5.96%	1 644.39	6.46%	5 231.16

**Table 3.29: National fossil carbon content in biofuels in 2023**

FEEDSTOCK	VOLUME	C FOSSIL PART	CARBON CONTENT	g FOSSIL CO <sub>2</sub> /g FAME
	m <sup>3</sup>	%	%	
Rapeseed	76 757.10	5.30%	75.50%	0.147
Palm oil	188.72	5.50%	71.80%	0.145
Sunflower seed	10 291.44	5.30%	77.20%	0.150
Used cooking oil*	52 278.82	5.40%	74.40%	0.147
<b>NATIONAL AVERAGE</b>	-	<b>5.34%</b>	<b>75.21%</b>	<b>0.147</b>

\*for used cooking oil are no data of carbon content available, thus data for lard were used

The CO<sub>2</sub> emissions from urea-based catalysts were estimated using COPERT 5 model for categories “heavy duty trucks Euro V and EURO VI” and “passenger cars diesel Euro 6 a, b, c, d-temp and d”. These vehicles occurred in Slovakia since 2010 and therefore, time series 2010 – 2023 were reported in this submission. As the number of vehicles with the SCR technology is equal to heavy duty vehicle in Euro VI category, the default value in COPERT model was used. In line with the UNFCCC Reporting Guidelines (these emissions are not energy-related), these emissions are allocated in the IPPU sector category 2.D.3 ([Chapter 4.5](#)).

**Railways (CRT 1.A.3.c)** – GHG emissions from railways were estimated from diesel oil consumed by the operation of diesel traction and using the simple tier 1 according to the IPCC 2006 GL. According to the key category analysis, this source is not key category in 2024 submission. The IPCC default emission factors were used, except for CO<sub>2</sub> were country-specific emission factor was used ([Table 3.14](#)). According to the previous UNFCCC recommendation, the country specific NCVs were used in calculations for time series and therefore the fuel consumptions (and subsequently GHG emissions). The NCVs of blended diesel oil and esters are shown in [Table 3.27](#).

The consumption of diesel oil for the motor traction in the Slovak Republic is obtained from the Railways Company, a. s. (ZSSK) annually. It is assumed that the consumption of diesel oil in motor traction of railways transport is equal to the diesel oil sold for the railways. The mobile sources of pollution in the

---

railways transport include vehicles of motor traction of ZSSK. This motor traction is divided into 2 basic groups of vehicles: motor locomotives (Traction 70) and motor wagons (Traction 80). The motor traction has been operated by four depots in the organizational structure of ZSSK since 2002 (Bratislava, Zvolen, Žilina and Košice).

In terms of implementing Directive No 2009/29/EC and Directive No. 2009/30/EC on replacement of fossil fuels with biofuels emissions from biomass are calculated and reported since 2007. The share of biomass in diesel oil was calculated as bio-component percentage, by weight of the total weight of the fuel ([Table 3.29](#)).

**Domestic navigation (CRT 1.A.3.d)** – Domestic navigation includes emissions from national shipping between ports on Danube River on Slovak territory and domestic shipping on lakes and dams for touristic purposes. According to the key category analysis, this source is not key category in 2025 submission.

Shipping between Slovak ports on Danube River: The Slovak Shipping and Ports Company is providing detailed information on diesel oil consumption on the Danube River. The consumption is allocated between national and international companies. The total fuels sold to international companies is reported in the Memo Items (1.D.1.b) and total fuels sold to national companies (Slovak Water Management Enterprise) is reported in the Domestic Navigation (1.A.3.d). This activity represents movements of ships between Slovak ports (Bratislava, Komárno and Štúrovo). This approach was introduced in 2005 and the reallocation of fuels led to the reallocation between subcategories 1.A.3.d and 1.D.1.b.

Shipping on lakes: The State Navigation Administration was officially requested to check availability of information about the shipping activity in the Slovak Republic except the Danube River movements. Only total number of ships and boats operated outside of the Danube River is registered, but without information about their activity or fuel consumption. Based on expert research three other relevant shipping routes occur in Slovakia, however in limited extent:

- River – basin of the Váh (Piešťany, Trenčín, Liptovská Mara dam);
- The tributary River of the Váh (Oravská Priehrada dam);
- River – basin of the Bodrog (Zemplínska Šírava dam).

While the public and tourist shipping activities in the Slovak Republic are not very frequent and have expanded only in the recent years (due increase of tourism), it was necessary to propose an appropriate methodological approach for emissions estimation. Chosen activity data were:

- The number of trips per year - is limited by the daily schedule of trips mostly in summer months (May-October);
- The duration of trips (in hours) - can differ according to the type of trips (mostly short or long tours);
- The technical parameters of the most populated ships – the country specific technical parameters of vessels can be found on the webpage. The engines are mostly with 100 kilowatts power, which is a common type of engine used in non-road mechanisms, or in agricultural machinery (type Zetor). The engines run mostly on diesel oil;
- The average consumption of diesel oil in litres per hour - based on technical description of the engines it is 12 litres of diesel oil per hour of work. The consumption of diesel oil in tons was calculated using average density of diesel oil (0.83 kg/dm<sup>3</sup>).
- During the pandemic year 2020 there was no traffic on lakes observed, thus no petrol and biofuels consumption was observed. Therefore notation keys “NO” were used.

The GHG emissions are calculated multiplying fuel consumption by diesel motor boats with emission factor. The country specific NCVs, obtained from the ŠÚ SR, were used to convert the fuels consumption in energy units. The NCV for diesel oil is shown in [Table 3.26](#). The emission factors are taken from the

IPCC 2006 GL and GHG emissions were recalculated for the whole time series. The default emission factors used in categories 1.A.3.d and 1.D.1.b are identical (*Table 3.30*). Activity data for domestic navigation are shown in *Tables 3.30* and *3.32*.

**Table 3.30: The emission factors used in GHG inventory for navigation in 2023**

PARAMETER	EMISSIONS FACTORS	
	DOMESTIC NAVIGATION	INTERNATIONAL NAVIGATION
EMISSIONS	<i>kg/TJ of fuel</i>	
CO <sub>2</sub>	74 196.78	74 196.78
CH <sub>4</sub>	7	7
N <sub>2</sub> O	2	2

**Table 3.31: Total fuels consumption (petrol + diesel) in domestic navigation in particular years**

YEAR	FUEL CONSUMPTION	
	TJ	t
1990	0.30	7.14
1995	0.27	6.51
2000	0.33	7.70
2005	0.47	11.08
2007	4.52	94.85
2008	4.79	99.38
2009	4.40	90.73
2010	4.49	104.49
2011	11.27	265.31
2012	14.96	352.35
2013	46.01	1 092.89
2014	59.11	1 403.26
2015	83.94	1 990.22
2016	64.24	1 524.12
2017	63.32	1 506.80
2018	34.53	819.41
2019	56.36	1 337.89
2020	72.25	1 716.75
2021	78.25	1 859.18
2022	71.45	1 690.20
2023	71.71	1 699.49

**Table 3.32: Diesel oil sold by shipping companies and allocation to the categories 1.A.3.d and 1.D.1.b**

YEAR	SHIPPING COMPANIES	SALE OF DIESEL OIL		
		NATIONAL	INTERNATIONAL	TOTAL
		1.A.3.d	1.D.1.b	1.A.3.d + 1.D.1.b
<i>t/year</i>				
2005	Slovak Shipping and Ports (Danube)	1.3	128.7	130
	International shipping companies	-	84	84
	Total	1.3	212.7	214
2010	Slovak Shipping and Ports (Danube)	91.8	9 087.20	9 179.00
	International shipping companies	0	1 363.00	1 363.00
	Total	91.8	10 450.20	10 542.00

YEAR	SHIPPING COMPANIES	SALE OF DIESEL OIL		
		NATIONAL	INTERNATIONAL	TOTAL
		1.A.3.d	1.D.1.b	1.A.3.d + 1.D.1.b
<i>t/year</i>				
2015	Slovak Shipping and Ports (Danube)	1 981.80	5 945.40	7 927.20
	Slovak Water Management Enterprise	NO	-	NO
	Other companies	0.5	47.5	48
	International shipping companies	-	1 016.00	1 016.00
	Total	1 982.30	7 008.90	8 991.20
2016	Slovak Shipping and Ports (Danube)	1 515.10	4 545.40	6 060.50
	Slovak Water Management Enterprise	-	NO	NO
	Other companies	2	189	191
	International shipping companies	-	1 272.00	1 272.00
	Total	1 517.00	6 006.50	7 523.50
2017	Slovak Shipping and Ports (Danube)	1 492.90	4 478.70	5 971.60
	Slovak Water Management Enterprise	-	NO	NO
	Other companies	2.4	236.6	239
	Morsevo (Komárno)	NO	1 034.00	1 034.00
	International shipping companies	-	168.5	168.5
	Total	1 495.30	5 917.80	7 413.10
2018	Slovak Shipping and Ports (Danube)	3 239.00	809.75	2 429.25
	Slovak Water Management Enterprise	-	NO	NO
	Other companies	232	2.32	229.68
	Morsevo (Komárno)	824	NO	824
	International shipping companies	-	NO	NO
	Total	4 295.00	812.07	3 482.93
2019	Slovak Shipping and Ports (Danube)	1 327.00	3 981.00	5 308.00
	Slovak Water Management Enterprise	NO	-	NO
	Other companies	3.26	322.74	326
	International shipping companies	-	760	760
	Morsevo (Komárno)	NO	NO	NO
	Total	1 330.26	5 063.74	6 394.00
2020	Slovak Shipping and Ports (Danube)	1 555.75	4 667.25	6 223.00
	Slovak Water Management Enterprise	NO	-	NO
	Other companies	161	NO	161
	International shipping companies	-	94	94
	Morsevo (Komárno)	NO	NO	NO
	Total	1 716.75	4 761.25	6 478.00
2021	Slovak Shipping and Ports (Danube)	1 764.25	5 292.75	7 057.00
	Slovak Water Management Enterprise	0	0	0

YEAR	SHIPPING COMPANIES	SALE OF DIESEL OIL			
		NATIONAL	INTERNATIONAL	TOTAL	
		1.A.3.d	1.D.1.b	1.A.3.d + 1.D.1.b	
<i>t/year</i>					
	Other companies	95	0	95	
	International shipping companies	0	165	165	
	TaM Terminal (Komárno)	NO	NO	NO	
	Total	1 859.25	5 457.75	7 317.00	
	2022	Slovak Shipping and Ports (Danube)	1 569.25	4 707.75	6 277.00
		Slovak Water Management Enterprise	0	0	0
Other companies		120	0	120	
International shipping companies		0	855	855	
TaM Terminal (Komárno)		0	0	0	
Total		1 689.25	5 562.75	7 252.00	
2023	Slovak Shipping and Ports (Danube)	1 614.50	4 843.50	6 458.00	
	Slovak Water Management Enterprise	0	0	0	
	Other companies	85	0	85	
	International shipping companies	0	902	902	
	TaM Terminal (Komárno) <sup>9</sup>	0	0	0	
	Total	1 699.50	5 745.50	7 445.00	

Slovakia reconstructed the time series for petrol fuel consumption as appropriate till 2008. Slovakia used expert judgement with the combination of statistical yearly income of the company, which operates the ships, and the yearly number of tourists in the region to estimate petrol consumption. Outcomes of this calculation are presented in [Table 3.33](#). During the data investigation it was found out that the company started the operation of these ships only in the year 2008 and after the COVID pandemic there is no information about further operation.

**Table 3.33:** *Outcomes of the petrol consumption reconstruction and emission estimation for the years 2008 – 2023*

YEAR	FOSSIL PETROL				BIO-PETROL			
	Energy	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Energy	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	<i>TJ</i>	<i>t</i>			<i>TJ</i>	<i>t</i>		
2008	0.0339	2.3486	0.0017	0.0001	0.0003	0.0218	0.00002	0.000001
2009	0.0389	2.6972	0.0019	0.0001	0.0008	0.0524	0.00004	0.000002
2010	0.0566	3.9244	0.0028	0.0001	0.0013	0.0880	0.00006	0.000003
2011	0.0508	3.5175	0.0025	0.0001	0.0012	0.0859	0.00006	0.000002
2012	0.0629	4.3602	0.0031	0.0001	0.0016	0.1107	0.00008	0.000003
2013	0.0549	3.8077	0.0027	0.0001	0.0015	0.1060	0.00008	0.000003
2014	0.0928	6.4306	0.0046	0.0002	0.0041	0.2810	0.00020	0.000008
2015	0.0428	2.9678	0.0021	0.0001	0.0017	0.1150	0.00008	0.000003
2016	0.0573	3.9742	0.0029	0.0001	0.0021	0.1428	0.00010	0.000004
2017	0.0573	3.9736	0.0029	0.0001	0.0021	0.1427	0.00010	0.000004

<sup>9</sup> Previously Morsevo

YEAR	FOSSIL PETROL				BIO-PETROL			
	Energy	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Energy	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	<i>TJ</i>	<i>t</i>			<i>TJ</i>	<i>t</i>		
2018	0.0639	4.4253	0.0032	0.0001	0.0027	0.1882	0.00014	0.000005
2019	0.0636	4.4602	0.0032	0.0001	0.0029	0.1892	0.00014	0.000005
2020	NO	NO	NO	NO	NO	NO	NO	NO
2021	NO	NO	NO	NO	NO	NO	NO	NO
2022	NO	NO	NO	NO	NO	NO	NO	NO
2023	NO	NO	NO	NO	NO	NO	NO	NO

**Pipeline transport (CRT 1.A.3.e.i)** - The consumption of natural gas used for energy to drive turbines in pipeline system were obtained from the NEIS database. Tier 2 approach and the country specific emission factor was used for CO<sub>2</sub> emissions estimation in pipeline. The emission factor for NG combustion is 56.18 t (CO<sub>2</sub>)/TJ in 2023.

### Uncertainties and Time-series Consistency

According to the previous recommendations, Slovakia is using hybrid combination of Approaches 1 and 2 in this submission for calculation of total uncertainty of the inventory (**Annex 3** of this Document). Uncertainty analyses performed by the Approach 1 in transport was carried out using Table 3.2 for uncertainty calculation and country specific uncertainties for activity data and emission factors were inserted into calculation table.

The Slovak Republic provided and published also Approach 2 for uncertainty analyses according to the Chapter 3 of the IPCC 2006 GL for the complete Energy (including transport) and the IPPU sectors for the year 2015. The methodology and results were described in previous SVK NIR 2017 and 2018. The latest Monte Carlo simulation was performed for the 2015 emissions inventory. Due to capacity reasons and according to the QA/QC plan in this sector, new calculation of Monte Carlo uncertainty (Approach 2) in the Energy sector and categories (including Transport) will be performed in the next submissions. For more information, please see the **Chapter 1.2** of this Document. Results of the Monte Carlo simulations were almost identical since this exercise was performed (since 2011).

Increasing quality of the emissions inventory from transport depends closely on the reduction and removal of the following uncertainties:

- The uncertainties joint with the COPERT methodology;
- The uncertainties joint with the collection, preparation and application of the input data.

The quality of calculated results by the COPERT 5 has been influenced significantly by the uncertainty of the following statistic information:

- Statistic information about consumption of the fuels;
- Allocation of total number of vehicles among all the categories according to the methodology;
- The average annual mileage;
- The average speed in the traffic mode;
- The average temperatures;
- The beta-factor.

COPERT 5 requires the determination of CH<sub>4</sub> emission factors and the calculation of CH<sub>4</sub> emissions accumulated, respectively, in order to determine:

- Data on the numbers of road vehicles in the Slovak Republic in current year, divided into categories prescribed by the methodology;

- 
- Data on average monthly temperatures in current year;
  - The average speed of vehicle categories in city, road and highway driving modes;
  - The annual mileage – will take place between categories of vehicles, divided into urban, rural and highway traffic.

**Domestic aviation (CRT 1.A.3.a)** – Trend in aviation transport for the years after 2008 is decreasing. The period 2004 – 2008 was influenced by the boom of low-cost airlines and advantage of Bratislava airport with the lower charges in comparison with the big international airports in the neighbouring countries. After this period, aviation transport decreased back on the 2003 level and the trend is very stable. The aviation regarding the national circumstances is not very important transport mode in Slovakia. The airports Bratislava, Košice and Poprad are the busiest airports. Other airports have only local character for hobby and sport flights.

**Road transport (CRT 1.A.3.b)** – Using of COPERT version 5 for whole time series (since 1990) is limited by availability of input data. Development in model structure and complexity does not allow to use the more advance versions before 2000. Trend in the CO<sub>2</sub> and N<sub>2</sub>O emissions from road transport corresponds with the consumption of the liquid fuels. Emission factors are annually updated based on national data. The variability is caused by changes in inputs for vehicle fleet, fuel consumption and emission factors. Until 2008, trend of petrol consumption has fluctuated and after 2008, the trend is stable due to the improvement in fuel consumption and implementation of renewable directive. In 2015 and 2016 the consumption increased and afterwards stabilized again. The trend of diesel oil consumption was increasing since 1990, but it is more stable in the recent years with temporary decrease in 2020. This was caused by the variation of fuel price in transit, the development of construction, commercial, industrial activities, economic development and, of course, by the trend of increasing numbers of new cars within the commercial market of the Slovak Republic, which significantly determines the development of the emissions from transport. In addition, the decrease of N<sub>2</sub>O is caused by significantly lower N<sub>2</sub>O EF for LPG passenger cars in category EURO 3 and newer. Cars in these category from year 2016 prevail in vehicle fleet. Significant decrease of CNG consumption is caused by change of vehicle fleet and decrease of CNG consumption in the biggest public transport providers (Public Transport Companies in Bratislava and Košice cities and Zvolen Bus-intercity Company).<sup>10</sup> CNG and older diesel oil buses are slowly replaced by electric and EURO 6 diesel buses. Decrease of methane emissions in the category 1.A.3.b.i (passenger diesel cars) is caused by significantly lower CH<sub>4</sub> EF for passenger cars in category EURO 3 and newer.

The elimination of negative influences of road transport continues with the increase of LPG, CNG and electric vehicles (mostly passenger cars and buses).

**Railways (CRT 1.A.3.c)** – Methodology, activity data and used emission factors for diesel oil are consistent for the whole time series. The blending of biomass in liquid fuels used in railways transport was considered since 2007.

**Domestic navigation (CRT 1.A.3.d)** – Emissions from domestic navigation represent emissions from shipping on lakes for the period 1990 – 2022 and emissions from shipping on lakes and movements between national ports on Danube River for the years 1990 – 2022. In 2022 there were no movements on lakes. The time series consistency was improved in previous submissions. Based on the expert judgement from the Slovak Shipping and Ports Company, before the year 2005, only negligible fuels were sold for national shipping on the Danube River. The variability in consumption is because of

---

<sup>10</sup> Companies do not have English equivalent names

neighbourhood of bigger ports in Vienna and Budapest and different prices and taxation of fuels used in shipping activities.

Pipeline transport (CRT 1.A.3.e.i) – Methodology, activity data and used emission factors for natural gas are consistent in the time series and energy-related categories (natural gas used in energy combustion).

### Category-specific QA/QC and Verification Process

Category specific QA/QC is based on the general QA/QC plan described in the **Chapter 1.2** of this Document. The emissions inventory in transport categories were prepared by the sectoral expert. Variety of input data sources and databases led to inconsistencies in transport fuel consumption occurrence in the last years. Therefore, in agreement with our Improvement Plan in Transport, the extensive analyses of the available statistical information in liquid fuels in transport began in the 2017. Results are summarized in the next paragraphs.

Source specific comparison of fuel statistics - QA/QC procedures for the transport follow basic rules and activities of QA/QC as defined in the IPCC 2006 GL. The QC checks were done during the CRT and NID compilation, general QC questionnaire was filled in and is archived. Also according to agreement with CDV (Centrum dopravného výskumu) from 2023, there is a QA/QC cross-check between Slovak and Czech Transport sector emission estimation.

Due to frequent questions for data consistency between the IEA statistics and the national inventory, the data sources were investigated. Comparison of activity data and their sources is also crucial for evaluation of consistency in reporting. Petrol, diesel oil and biofuels consumption are key activity data in transport, thus the comparison was focused on these statistical data across several sources.

Datasets for this analysis are the years 2014 – 2021:

- Statistical Office of the Slovak Republic (ŠÚ SR) inserts data also from the Administration of State Material Reserves of the Slovak Republic (ŠHR SR);
- Ministry of Economy of the Slovak Republic (MH SR);
- Finance Administration of the Slovak Republic (FR SR);
- Ministry of Environment of the Slovak Republic (MŽP SR).

Each source has specific forms or questionnaires, CN codes and different reporting rules, methodologies and dates of publication or collection. Different institutions further process these data. The ŠÚ SR used import/export and production data, the FR SR used data from taxes on sales of products of crude oil and from taxes on sales of biofuels.<sup>11,12</sup>

**Table 3.35: Crude oil and crude oil products data flow and utilisation (final user is the SHMÚ)**

ORIGIN OF DATA	PRIMARY USER	SECONDARY USER
Import-export data (ŠÚ SR - Depart. of Foreign Trade)	Statistical Office of Slovak Republic (Depart. of Energy Statistics)	EUROSTAT
Data regarding production and sales (companies)		Slovak Hydrometeorological Institute
Data from taxes on sales of biofuels		Ministry of Economy

<sup>11</sup> Council Directive (EU) 2015/652 laying down calculation methods and reporting requirements pursuant to Directive 98/70/EC of the European Parliament and of the Council relating to the quality of petrol and diesel fuels

<sup>12</sup> Act 309/2009 Coll. on the Promotion of renewable energy sources and high-efficiency cogeneration and on amendments to certain acts as amended, <http://www.minzp.sk/en/areas/renewable-energy-sources/biofuels-bioliquids/>




ORIGIN OF DATA	PRIMARY USER	SECONDARY USER
Data from taxes on sales of products of crude oil	Financial administration of Slovak Republic	SK - BIO <sup>13</sup>
Confirmation (certificate) of the sustainability of biofuels	Slovak Hydrometeorological Institute (according to Art. 7a of Directive 98/70/EC)	European Environmental Agency
Data on production and sales (companies)	Slovak State Material Reserves	International Energy Agency (data on crude oil and crude oil products) EUROSTAT (natural gas)
Data of fuel sales on gas stations (NEIS)	Ministry of Environment (according to Art. 8 of Directive 98/70/EC)	European Environmental Agency

As it is shown in **Table 3.34** and on **Figure 3.9**, discrepancies occurred between major data sources-providers. During discussions with the main authorities, several information was collected by the sectoral experts, which were further analysed:

- Each authority reports different data in different forms for different institutions or requirements (**Table 3.34**);
- The conversion factors (e.g. density) differ throughout all data suppliers not only between authorities and companies, but also for each delivered supply has own characteristics;
- Dates of collection for tax reports and reports to the ŠÚ SR differ.

**Figure 3.9: Results of fuels consumption comparison according to different sources (thousand m<sup>3</sup>)**



The main outcome of this analysis is harmonisation of fuels consumption in country on the most possible level and lowering the differences in reporting by different subjects in 2023. Full consistency of data on national level is not possible. This is due to different legislation that each authority is required to fulfil (e.g. statistical reporting to EU institutions, tax collection, etc.).<sup>14</sup>

**Domestic aviation (CRT 1.A.3.a)** – Since 2011, the agreement of the European Commission (EC) and the EUROCONTROL is in place. Based on this agreement, annual comparison of aviation fuel consumption and emissions data with AEM model calculations is prepared. The comparison of the EUROCONTROL and the UNFCCC aviation data is provided on the level of individual EU Member State (EU MS). The information and data evaluated are part of the QA/QC activities in aviation. The EC works


<sup>13</sup> [SK-BIO](#) is the national register for biofuels and bioliquids

<sup>14</sup> Regulation (EC) 1099/2008 of the European Parliament and of the Council, Act No. 268/2017, which amend Act No. 98/2004 Coll. on the Excise Duty on mineral oil as amended, which amends Act No. 309/2009 Coll. on the Promotion of renewable energy sources and high-efficiency cogeneration and on amendments to certain acts as amended (only § 14a), <https://www.financnasprava.sk/en/businesses/taxes-businesses/excise-duties-businesses#TaxRatesMineralOil>

towards making data from the EUROCONTROL available to the EU MS on a regular basis, for quality check, however this information is not possible to make public available. Consistent time series (**Figure 3.10**) is maintained by using calculated average EFs from EUROCONTROL. The methodology is explained in the **Chapter 3.2.8**.

The verification process is also based on cross-checking of input data from the Slovak airports and the comparison with the sectoral statistical indicators (ŠÚ SR). The background documents are archived by the sectoral experts in the central archiving system at the SHMÚ. The quality manager of the NS SR has responsibility for the verification, approval and archiving.

**Figure 3.10:** Demonstration of time-series consistency between pre-EUROCONTROL methodology and EUROCONTROL methodology



**Road transport (CRT 1.A.3.b)** – QC activities ensuring the quality standards for the preparation of the emissions inventory in the road transport are based on the cooperation of several experts and institutions. The activity data and input parameters provided from the different data sources are collected and then checked for the basic quality criteria (consistency, transparency, etc.) and archived by the sectoral experts. The process of verification is based on cross-checking of input data from the ŠÚ SR and the comparison with the fuel balance from the COPERT. The background documents are archived by the sectoral experts and in central archiving system of SNE at SHMÚ.

**Other/Urea based catalysts (CRT 1.A.3.b.v allocated in 2.D.3)** – The COPERT 5 model was used for these emissions estimation and information of category specific QA/QC and verification are described in section road transport.

**Railways (CRT 1.A.3.c)** – Verification process is based on cross-checking of the input data on fuel consumption from the Railways Company, a. s. and the ŠÚ SR. The preliminary results of emissions inventory are sent to other subjects (MŽP SR) for valuation and QA activities. The QC verification process includes the comparison of statistical and calculated data on fuel consumption.

**Domestic navigation (CRT 1.A.3.d)** – Verification of activity data on fuels sold for shipping activities was performed by the sectoral expert and compared with the statistical information from requested institutions and companies as mentioned in this chapter above. New survey among small companies and municipalities operating touristic boats and ships on lakes and dams in Slovakia was made during the year 2020. These data were used to estimate the emissions from domestic shipping in 2019.

**Pipeline transport (CRT 1.A.3.e.i)** – Information of category specific QA/QC and verification are described in section for fugitive emissions 1.B.

---

## Category-specific Recalculations

This chapter describes the recalculations of emissions from fugitive emissions and its subcategories with respect to the previous submission. The list of recalculations is provided in this chapter.

**Revision of methodology** – COPERT model update was conducted and following changes were done:

- Corrected N<sub>2</sub>O emissions of petrol hybrid and petrol PHEV Euro 5 and 6 Passenger cars,
- Removal of CO<sub>2</sub> correction from the model,
- Updated emission factors of Euro 6 CNG passenger cars,
- Updated emission factors of Euro VI diesel buses,
- Updated emission factors of Euro VI diesel hybrid buses,
- Hot CH<sub>4</sub> emission factors of LPG cars corrected,
- Introduction of Euro VI CNG & LNG HDVs to the model

More information can be found on the COPERT model webpage<sup>15</sup>. Changes were done between version 5.5.1 and 5.8.1. These methodological adjustments resulted in changes in CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions during the period 2013 – 2022 (**Table 3.35**).

**Table 3.35: Differences in emissions between previous submission and current submission caused by recalculations**

YEAR	CO <sub>2</sub>	CH <sub>4</sub>
	Gg	
2013	0.01	0.00
2014	0.01	0.00
2015	0.02	0.00
2016	0.01	0.00
2017	0.02	-0.02
2018	0.01	-0.01
2019	0.01	-0.02
2020	0.01	-0.02
2021	1.10	-0.02
2022	0.02	-0.03

## Category-specific Improvements and Implementation of Recommendations

In the preliminary findings under the UNFCCC, there were no recommendations for the transport sector.

### 3.2.9. Other Sectors (CRT 1.A.4)

The source category 1.A.4 Other Sectors includes stationary combustion in agriculture, forestry, The source category 1.A.4 Other Sectors includes stationary combustion in agriculture, forestry, commercial and institutional and households.

**Commercial/Institutional (1.A.4.a)** – total volume of fuels in 1.A.4.a expressed in energy units represented 23 908.55 TJ in 2023. Total CO<sub>2</sub> emissions were 1 332.68 Gg, total CH<sub>4</sub> emissions were 0.313 Gg and total N<sub>2</sub>O emissions were 0.0072 Gg in 2023.

---

<sup>15</sup> <https://copert.emisia.com/copert/versions/>

---

**Residential (1.A.4.b)** – total volume of fuels in 1.A.4.b expressed in energy units represented 63 394.02 TJ in 2023. Total CO<sub>2</sub> emissions were 2 611.44 Gg, total CH<sub>4</sub> emissions were 6.0759 Gg and total N<sub>2</sub>O emissions were 0.07881 Gg in 2023.

**Agriculture, forestry and fisheries (1.A.4.c)** – total volume of fuels in 1.A.4.c expressed in energy units represented 4 794.38 TJ in 2023. Total CO<sub>2</sub> emissions were 253.41 Gg, total CH<sub>4</sub> emissions were 0.1141 Gg and total N<sub>2</sub>O emissions were 0.06459 Gg in 2023. The fuels are allocated in solid, liquid, gaseous and biomass fuels categories. In recent years, the IEF of biomass has shown a significant year-to-year fluctuation, especially in the case of CH<sub>4</sub>. This fluctuation is caused by changes in the ratio of biogas and solid biomass in the fuel mix.

The large difference in the EF of biogas and solid biomass is therefore also translated into the total IEF of biomass.

All non-road mobile machinery is also reported in this category. Agricultural machinery (tractors, harvesters, etc.), forestry machinery, industry machinery (forklifts, excavators, etc.) and residential machinery (hedge cutters, garden shredders, etc.) are included in the category 1.A.4.c.ii. The data collected by questionnaires in households in the frame of the project “Quality Improvement of Air Emission Accounts and Extension of Provided Time-series” were used for estimation of emissions from residential machinery the first time in 2018 inventory. In addition, liquid fuels used in residential machinery (hobby, gardens, cleaning) were collected and reported in the 1.A.4.c.ii.

#### **Methodological Issues, Activity Data, Emission Factors and NCVs**

A description of general methodologies used for GHG emissions estimation from fuel combustion is given in the **Chapters 3.2.6** and **3.2.7**.

Activity data (emission factors and NCVs) are collected from several sources (in agreement with the other energy categories):

- Annual energy balance (publication Energy,<sup>16</sup> published by the ŠÚ SR, annually);
- Disaggregated data provided by the ŠÚ SR (restricted from public, provided only for the SNE);
- The NEIS Central database;
- Results from project, surveys and research.

The Residential category is the key emissions source and represents 10.7% share on the total GHG emissions in the year 2023. Category 1.A.4.b balanced mostly gaseous (natural gas), solid (coal) and biomass (wood) fuels. Whereas the gaseous fuels consumption is consistent and accurate due to statistics made directly by the natural gas suppliers on distribution network, solid fuels and biomass statistics were not fully covered by the ŠÚ SR. Direct regular statistics is missing. Due to these reasons, several inconsistencies between fuels consumption reported in this category were recorded and commented in the previous submissions. Therefore, in 2018, the Project Grant “Quality Improvement of Air Emission Accounts and Extension of Provided Time series” launched by the European Commission – EUROSTAT was successfully finished. Results were published online in several partial reports and on the international conferences. The Project Grant was carried out in cooperation with the Statistical Office of the Slovak Republic and concluded in December 2022. Outcomes and Final Report will be available after validation from the EUROSTAT in 2023. Cooperation with the Statistical Office of the Slovak Republic continued and resulted in to the second more complex statistical survey in households, with primary solid fuels heating. This activity, together with help and interest of other relevant national

---

<sup>16</sup> Energy 2022, Statistical Office of Slovak Republic (2023) ISBN: 978-80-8121-918-4

authorities, confirmed and improved previous estimation of solid fuels and biomass consumption in households.

In addition, in the frame of the project LIFE IP – Improvement of air quality supported by the European Union, the OEaB experts have prepared a report that describes the structured distribution of small sources of pollution (available only in Slovak language). The main task of the analysis presented in this annual report was to obtain information on the regional distribution of boiler types in Slovakia. In addition, precise estimates of the consumption of solid fuels at the regional level, especially biomass (firewood), were developed. The input data were obtained on the basis of an extensive third statistical survey carried out in 2022 in cooperation with the Statistical Office of the Slovak Republic. The statistical sample was chosen in order to streamline the results and thus allow for a more even distribution of small combustion sources at the regional level. The results presented in this Document will serve to identify and select regions where modernization of boilers is needed and will help direct the decision-making process of allocating funds in the form of subsidies.

In previous inventory, data on solid fuels and biomass (wood) energy consumption in households collected and evaluated in a frame of this Project Grant were used and updated. Statistical data and time series were corrected based on improved methodology and inputs were also provided to the ŠÚ SR for energy balance. According to the information provided by the ŠÚ SR, revision of households' energy statistics to the EUROSTAT was reported for the year 2018 and expected revision will be provided to EUROSTAT also for time series in this year. Revision was focused on solid fuels and biomass (non-fossil fuels) consumption since the year 2012. With this revision, consistency in the reporting data in households was improved. In 2022, third survey focused on households with individual heating with solid fuels took place. New results are available in 2024 inventory submission for biomass and solid fuels consumption.

Methodology introduced by new background data further corrected and improved the energy and emissions balance considering the effect of regional-climatological data. The principle of new methodological approach was supported by statistical survey and further estimation of “total energy demand for heating and hot water preparation” in households, calculated using data from questionnaires and climatological data in different regions. In principle, average value of “energy demand” is a parameter on heating demand (including preparation of hot water) for 1 m<sup>2</sup> of housing area for 1 year. Total housing area, energy effectivity of houses and climatological factors in regional scaling were taking into consideration for the calculation of total energy demand for heating in houses without central heating system.

**Table 3.36: Overview of the country or plant specific CO<sub>2</sub> EFs in t/TJ the category 1.A.4 in 2023**

1.A.4.a	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	69.78	Liquefied Petroleum Gases	17.22	63.14
		Gas/Diesel Oil	20.34	74.58
		Residual Fuel Oil	20.80	76.30
Solid	96.68	Lignite	26.89	98.60
		Brown coal briquettes	26.61	97.57
		Other Bituminous Coal	25.96	95.19
		Gas Coke	29.60	108.53
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	79.69	Wood/Wood waste	30.50	111.83
		Sludge gas	14.90	54.63
1.A.4.b	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	63.14	Liquefied Petroleum Gases	63.14	63.14
Solid	97.69	Other Bituminous Coal	25.96	95.19

		Lignite	26.89	98.60
		Brown coal briquettes	26.61	97.57
		Gas Coke	29.80	109.27
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	111.83	Wood/Wood waste	30.50	111.83
<b>1.A.4.c</b>	<b>WEIGHTED CO<sub>2</sub> EFs</b>	<b>FUEL TYPE</b>	<b>C EFs</b>	<b>CO<sub>2</sub> EFs</b>
Liquid	72.73	Liquefied petroleum gases	17.22	63.14
		Gas/Diesel oil	20.34	74.58
		Diesel oil	20.24	74.20
		Petrol	18.37	67.36
Solid	97.38	Lignite	26.89	98.60
		Gas coke	29.80	109.27
		Other bituminous coal	25.96	95.19
		Brown coal briquettes	26.61	97.57
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	77.44	Other biogas	14.90	54.63
		Wood/Wood waste	30.50	111.83
		Other primary Solid biomass	27.30	100.10

### Uncertainties and Time-series Consistency

Description of uncertainty is similar to the [Chapter 3.2.6](#) of this Document.

Time series is consistent in all aspects (methodological approach, country specific EFs and oxidation factor used, fuel characteristics, etc.) to the detailed level of disaggregation (on plant specific level).

### Category-specific Recalculations

Recalculations were made in sector 1.A.4.b based on improved data about the number of apartments connected to district heating system. This resulted in changes of fuel consumption for this sector. This affected the calculation of biomass consumption in households, leading to a reduction in the fuel (biomass) consumption for heating in residences. This recalculation affected biomass consumption in the years 2023 to 2021, the base year was not affected. The comparison of original data and recalculated is summarized in following table.

The comparison of original data and recalculated is summarized in following table.

**Table 3.37: Comparison of recalculated data in 2024 and 2025 submissions**

YEAR	SUBMISSION 2024				SUBMISSION 2025			
	ENERGY	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	ENERGY	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	TJ	Gg			TJ	Gg		
2021	28 811.3	3 222.1	8.6434	0.1152	22 068.6	2 468.0	6.6206	0.08827
2022	24 834.9	2 777.4	7.4505	0.0993	20 718.5	2 317.0	6.2156	0.08287

### Category-specific Improvements and Implementation of Recommendations

Improvements are implemented in line with the Improvement and Prioritization Plan for the year 2025. Further improvements in the category 1.A.1.4.a are not planned in the near future.

#### 3.2.10. Non-Specified (CRT 1.A.5)

Emissions reported in this category arising from the military aviation and from fuel combustion in stationary sources that are not specified elsewhere. Total volume of fuels in the 1.A.5 expressed in energy units represented 1 538.69 TJ in 2023.

Total CO<sub>2</sub> emissions were 65.31 Gg, total CH<sub>4</sub> emissions were 0.0868 Gg and total N<sub>2</sub>O emissions were 0.0005 Gg in 2023.

### Methodological Issues, Activity Data, Emission Factors and NCVs

A description of the general methodology, activity data, EFs and NCVs used for estimation of emissions from fuels combustion is given in the **Chapters 3.2.6** of this Document.

In 1.A.5.a, the main source of activity data is provided by the ŠÚ SR (disaggregated data – information on fuels consumption at the level of individual subjects). The sources allocated here are not included in the EU ETS. Total volume of fuels in the 1.A.5.a expressed in energy units represented 1 480.37 TJ in 2023. Total CO<sub>2</sub> emissions were 61.07 Gg, total CH<sub>4</sub> emissions were 0.0082 Gg and total N<sub>2</sub>O emissions were 0.0002 Gg in 2023.

The jet kerosene, petrol and diesel oil from military usage is reported in the 1.A.5.b. GHG emissions from military aviation, i.e. jet kerosene consumption, are estimated since 1990 and military petrol and diesel oil are estimated since 2016. Data for military petrol and military diesel oil before 2016 were statistically estimated by the sectoral experts using linear regression back to basic year 1990 based on years 2016 – 2019. The information is directly provided by the Ministry of Defence of the Slovak Republic. Also fuels used for military machinery do not have a biofuel part. The methodology is comparable with the methodology used for the emissions estimation of civil aviation, based on fuel consumption in military service multiplied by the default emission factor for jet kerosene. **Table 3.38** provides overview of the weighted average emission factors and fuels in the category 1.A.5 for 2023.

**Table 3.38: Overview of the country or plant specific CO<sub>2</sub> EFs in t/TJ in the category 1.A.5 in 2023**

1.A.5	WEIGHTED CO <sub>2</sub> EFs	FUEL TYPE	C EFs	CO <sub>2</sub> EFs
Liquid	70.15	Liquefied petroleum gases	17.22	63.14
		Residual fuel oil	20.80	76.30
		Diesel oil	20.24	74.20
		Jet kerosene	19.84	72.75
		Petrol	18.37	67.36
Solid	98.62	Gas coke	29.80	109.27
		Lignite	26.89	98.60
		Other bituminous coal	25.96	95.19
Gaseous	56.18	Natural gas	15.32	56.18
Biomass	55.01	Sludge gas	14.90	54.63
		Other biogas	14.90	54.63
		Other primary solid biomass	27.30	100.10
		Wood/Wood waste	30.50	111.83

### Uncertainties and Time-series Consistency

Description of uncertainty is similar to the **Chapter 3.2.6** of this Document. Time series is consistent in all aspects (methodological approach, country specific EFs and oxidation factor used, fuel characteristics, etc.) to the detailed level of disaggregation (on plant specific level).

### Category-specific Recalculations

No recalculations were implemented in this submission.

### Category-specific Improvements and Implementation of Recommendations

Improvements are implemented in line with the Improvement and Prioritization Plan for the year 2024, no specific improvement is planned for the next submission.

---

### 3.3. Comparison of the Sectoral Approach with the Reference Approach (CRT 1.AC)

The data gathered and processed by the Statistical Office of the Slovak Republic (the annual energy statistics balance) is the background for the reference approach. Therefore, the data provided in the reference approach is consistent with official energy balance data. The reference approach balance includes emissions from fuel combustion differentiated according to the gaseous, liquid, solid and biomass categories and different sectors.

The reference approach is based on the top-down methodology and is characteristic of minimum requirements on input data. The reference approach provides only aggregated estimates of emissions by fuel type distinguishing between primary and secondary fuels. The aggregated nature of the reference approach means that stationary combustion cannot be distinguished from the mobile combustion. The method is applied also as the quickest control and verification method. It is necessary to mention, that this approach does not include fugitive emissions, i.e. uncontrolled emissions from mining and post-mining activities, from transport and other use of fuels (technological use).

The methodology for reference approach estimation is consistent during time series across of the main types of fuels and followed the methodology provided in the IPCC 2006 GL.

The official frame contract was signed between the Statistical Office of the Slovak Republic and the Ministry of Environment to ensure direct cooperation with the National Inventory System (SHMÚ). Frame contract specifies major responsibilities in providing information about energy balance and any changes or recalculations directly to the ŠÚ SR. A close cooperation of the NS and the ŠÚ SR ensures consistency and transparency in reporting. The cooperation on the official level and the ongoing discussions on removing any discrepancy between the several statistical systems of energy data (NEIS, ŠÚ SR or EU ETS) is in place. A bottom-up methodology was used for the emissions balance in the sectoral approach. More information is provided in the [Chapter 3.2](#) of this Document.

Based on the actual data provided in the 2025 submission, time series consistency was improved leading to increase of transparency reported in this area ([Figure 3.11](#)). A difference between CO<sub>2</sub> emissions allocated in reference and in sectoral approaches is less than 2% for last eight years. In 2023, the difference in CO<sub>2</sub> emissions was -0.50% and difference in the total energy consumption was 0.02%.

The reference and sectoral approach were estimated on fully independent data sets, whereby obtained differences in CO<sub>2</sub> emissions are not significant. Based on the IPCC methodology, reference approach in apparent consumption of fuels was estimated after consideration of carbon stored in iron and steel and in chemical industry and refinery. Due to the different methodology used by the ŠÚ SR, not all fuels used as technological input in production are also reported in the statistical questionnaires in this way. This is a case of natural gas used in ammonia production (allocated in the IPPU sector, but in the statistical questionnaire allocated in the Energy sector), or coking coal used as reducing agent in steel production (allocated in the IPPU, but in the statistical questionnaire allocated in the Energy sector).

These reallocations were considered in the apparent consumption and the results are provided in [Tables 3.39 - 3.43](#). However, due to differences in methodological approaches used in the national inventory for the sectoral approach and used in the statistical energy balance, in some years the differences are higher than required according to the QA/QC process. After thorough analyses of these years (1996-1998, 2003, 2009, 2015), the results show the major inconsistencies in other and liquid fuels.

One of the reasons for the reference and sectoral approach discrepancies during time series is used source of activity data. The RA is based on national fuel delivery statistics, the bottom-up approach is based on fuel consumptions (EU ETS reports and disaggregated energy balance data). However, the




main reason is the effect of emission factors (and/or calorific values) in the reference approach of liquid fuels. This is enhanced by the fact that all volume of used crude oil which is processed in the Slovak Republic is imported. Practically all resulting CO<sub>2</sub> emissions from combustion of the liquid fuels reported in the reference approach is from the import, export and stock changes of crude oil. A small variation in the average net calorific value used (which is difficult to determine), has a large influence on the total CO<sub>2</sub> emissions. Similar situation is also in calorific values and emission factors of naphtha, lubricants and bitumen, which are used to estimate the fraction of carbon stored.

Further significant difference is visible in the case of waste. Based on our research, the main source of the difference is caused by data processing methodology of the ŠÚ SR on waste incinerated. An incorrect categorization of municipal and industrial waste in the energy balance provided by the ŠÚ SR was identified. Moreover, the estimation of composition (biogenic/fossil part) of waste in the SA is based on information provided directly by the operators. Several meetings are organized with the experts from the ŠÚ SR on this issue.

In 2023, the emissions decrease in most categories of Energy sector. For three years in a row (between 2020-2023), a continuous increase in fuel consumption in the transport sector can be observed. In 2023 the emission remain practically identical with previous year (inter-annual decrease of 43 kt CO<sub>2</sub>).

**Figure 3.11: Difference between the reference and sectoral approaches CO<sub>2</sub> emissions (in Gg) in 1990 – 2023**



**Table 3.39: The comparison of the RA and the SA in total fuels consumption and CO<sub>2</sub> emissions in 1990 – 2023**

YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	EMISSIONS DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
1990	753	660	641	-2.99	52 572	53 273	-1.32
1991	659	586	568	-3.00	46 616	46 952	-0.72
1992	625	540	539	-0.08	43 007	42 838	0.39
1993	587	498	495	-0.48	39 263	39 038	0.58
1994	562	469	460	-1.91	36 461	36 566	-0.29
1995	591	474	487	2.65	37 395	36 236	3.2
1996	600	476	493	3.62	37 776	35 883	5.27
1997	600	477	494	3.51	37 524	35 858	4.65
1998	583	475	477	0.31	36 135	35 293	2.39
1999	566	470	461	-1.84	34 848	34 625	0.64
2000	546	464	455	-2.00	33 699	33 938	-0.7
2001	577	494	494	-0.13	36 043	35 974	0.19

YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	EMISSIONS DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
2002	560	463	456	-1.52	34 107	33 760	1.03
2003	565	468	465	-0.74	35 643	34 683	2.77
2004	555	464	447	-3.74	34 551	34 154	1.16
2005	567	471	464	-1.64	34 861	34 662	0.57
2006	551	450	441	-1.93	33 974	33 921	0.16
2007	531	433	423	-2.36	32 233	32 395	-0.5
2008	533	443	431	-2.83	32 592	32 867	-0.84
2009	482	405	383	-5.52	29 175	30 484	-4.29
2010	514	412	413	0.21	31 331	30 824	1.64
2011	492	400	393	-1.68	30 235	30 348	-0.37
2012	466	377	370	-2.03	28 361	28 291	0.24
2013	468	375	366	-2.32	27 489	28 116	-2.23
2014	428	338	330	-2.48	25 431	25 934	-1.94
2015	433	351	337	-4.13	25 641	26 623	-3.69
2016	443	354	344	-2.67	26 292	26 793	-1.87
2017	473	369	372	0.74	28 208	27 760	1.61
2018	474	366	368	0.61	28 105	27 690	1.50
2019	441	354	352	-0.52	26 213	26 206	0.03
2020	423	332	332	0.26	24 199	24 031	0.70
2021	472	362	361	-0.30	26 771	26 716	0.21
2022	424	333	332	-0.15	24 655	24 521	0.54
2023	411	320	320	0.02	23 771	23 891	-0.50

**Table 3.40:** The comparison of the RA and the SA in liquid fuels consumption and CO<sub>2</sub> emissions

YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	EMISSIONS DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
1990	197	164	155	-5.71	11 628	12 252	-5.09
1995	145	103	109	6.3	8 084	7 662	5.51
2000	122	92	89	-2.53	6 769	6 769	0.00
2005	139	117	107	-8.34	8 333	8 651	-3.68
2010	144	117	114	-2.2	8 729	8 542	2.18
2015	129	120	108	-10.35	7 952	8 803	-9.67
2019	150	126	127	0.66	9 495	9 300	2.09
2020	150	118	119	1.46	8 861	8 622	2.77
2021	156	125	123	-1.76	9 165	9 219	-0.58
2022	160	131	130	-0.66	9 697	9 585	1.18
2023	155	129	129	-0.39	9 345	9 475	-1.38

**Table 3.41:** The comparison of the RA and the SA in solid fuels consumption and CO<sub>2</sub> emissions in particular years

YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
1990	342	282	286	1.25	29 866	28 958	3.14
1995	226	157	170	8.15	17 796	16 564	7.44

YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
2000	179	134	135	0.85	14 125	13 921	1.47
2005	178	124	123	-0.86	13 556	13 263	2.21
2010	159	100	99	-0.6	11 492	11 383	0.96
2015	137	81	80	-1.06	9 257	9 331	-0.8
2019	114	66	65	-1.96	7 541	7 687	-1.9
2020	97	54	54	-1.55	6 237	6 303	-1.05
2021	118	59	59	-0.34	7 350	7 350	-0.01
2022	100	53	53	-0.55	6 352	6 375	-0.36
2023	100	49	49	0.7	6 260	6 272	-0.2

**Table 3.42:** The comparison of the RA and the SA in gaseous fuels consumption and CO<sub>2</sub> emissions in particular years

YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
1990	214	211	200	-5.47	11 062	11 827	-6.47
1995	221	212	207	-2.41	11 472	11 814	-2.9
2000	244	237	230	-2.87	12 748	13 070	-2.46
2005	247	229	232	1.17	12 805	12 614	1.52
2010	210	193	198	2.78	10 974	10 640	3.14
2015	162	146	144	-1.08	8 043	8 131	-1.09
2019	171	156	155	-0.71	8 630	8 691	-0.7
2020	171	154	154	-0.11	8 575	8 583	-0.09
2021	191	172	173	0.65	9 692	9 629	0.65
2022	159	144	145	0.74	8 123	8 064	0.74
2023	151	137	137	0.24	7 719	7 701	0.23

**Table 3.43:** The comparison of the RA and the SA in other fossil fuels consumption and CO<sub>2</sub> emissions in particular years


YEAR	RA	SA	APPARENT ENERGY CONSUMPTION	ENERGY CONSUMPTION DIFFERENCE	RA	SA	DIFFERENCE
	PJ			%	Gg of CO <sub>2</sub>		%
1990	0.18	2.55	0.18	-92.97	16	236	-93.18
1995	0.48	2.1	0.48	-77.35	43	197	-78.34
2000	0.64	1.91	0.64	-66.61	57	180	-68.12
2005	1.89	1.43	1.89	31.59	168	135	24.79
2010	1.53	2.86	1.53	-46.66	136	259	-47.43
2015	4.4	3.91	4.4	12.4	389	342	14.02
2019	5.42	5.66	5.42	-4.29	546	524	4.22
2020	5.87	5.72	5.87	2.61	526	522	0.83
2021	6.03	5.82	6.03	3.74	565	518	9.05
2022	5.26	5.67	5.26	-7.14	482	498	-3.26
2023	4.93	5.04	4.93	-2.21	448	442	1.40

### 3.4. Feedstocks and Non-energy Use of Fuels (CRT 1.AD)


Using the IPCC 2006 GL, the quantity of carbon excluded from the RA (carbon used for ammonia production, petrochemicals production, carbide production, hydrogen production, iron and steel production, ferroalloys production, aluminium production as well as non-energy using of lubricants) was estimated. Total carbon excluded from the RA was 1 630.04 Gg in 2023, which represented 5 976.80 Gg of CO<sub>2</sub>. The emissions from the carbon excluded are reported in respective categories in the IPPU sector.

The major share of carbon excluded represents the carbon from coking coal, both in fuel consumption and in amount of carbon (55.8% and 55.3%, respectively) The other significant sources of carbon excluded are using of natural gas (15.0% in fuel consumption and 12.8% in quantity of carbon) and using of naphtha (15.8% in fuel consumption and 17.6% in quantity of carbon). Details on the share in fuel units and carbon units are presented on **Figures 3.12** and **3.13**. The CO<sub>2</sub> emissions excluded from the RA are presented on **Figure 3.14** for the whole time series 1990 – 2023.


**Figure 3.12:** The share of different fuels consumption for feedstock and non-energy use in 2023



**Figure 3.13:** The share of carbon for feedstock and non-energy use in 2023



**Figure 3.14:** The CO<sub>2</sub> emissions (Gg) according to the fuels excluded from the RA in 1990 – 2023



Liquid fuels (petroleum coke, naphtha, and refinery feedstocks), solid fuels (coking coal, other bituminous coal) and gaseous fuels (natural gas) are used as feedstock in Slovakia. Lubricants and bitumen (liquid fuels) are used for non-energy purposes. The respective amounts of mentioned fuels are allocated in the IPPU sector and emissions are included there. The allocation of the fuels excluded from the RA and included in the IPPU sector is presented in [Tables 3.44](#) and [3.45](#).

**Table 3.44:** The allocation of fuels excluded from the RA in the IPPU sector

FUEL	USED AND REPORTED IN CATEGORIES
Natural gas	2.B.1 Ammonia production 2.B.8 Petrochemicals 2.B.10 Hydrogen production 2.C.1 Iron and steel production
Petroleum Coke	2.C.3 Aluminium production
Naphtha	2.B.8 Petrochemicals
Lubricants	2.D.1 Lubricants
Bitumen	2.D.3 Solvents use
Refinery feedstock	2.B.8 Petrochemicals
Coking coal	2.B.5 Carbide production 2.C.1 Iron and steel production 2.C.2 Ferroalloys production
Other bituminous coal	2.B.5 Carbide production 2.C.1 Iron and steel production 2.C.2 Ferroalloys production

The plant-specific (where available) and country-specific NCVs and EFs are used for estimation the volume of carbon excluded and respective CO<sub>2</sub> emissions excluded from the RA. Natural gas, petroleum coke, naphtha, lubricants, refinery feedstock, coking coal and other bituminous coal were balanced as feedstock and non-energy use of fuels. The quantities of the fuels and carbon used for non-energy purposes were provided directly by the plant operators or by the ŠÚ SR. The results are presented in [Table 3.45](#).

**Table 3.45:** Total volume of carbon in different fuels excluded from the RA in particular years

YEAR	Natural Gas	Petroleum Coke	Naphtha	Lubricants	Bitumen	Refinery Feedstock	Coking Coal	Other Bituminous Coal
	kt							
1990	250.61	NO	296.25	65.54	418.77	65.58	1 209.70	IE
1995	254.92	NO	362.98	65.54	199.63	76.18	1 231.99	IE
2000	274.56	37.94	395.73	65.54	83.40	65.80	937.52	IE

YEAR	Natural Gas	Petroleum Coke	Naphtha	Lubricants	Bitumen	Refinery Feedstock	Coking Coal	Other Bituminous Coal
	kt							
2005	329.10	66.86	347.70	39.49	126.88	67.55	1 025.05	37.72
2010	263.78	65.44	338.98	16.90	112.07	63.64	1 111.31	37.91
2011	345.90	58.88	333.75	25.27	130.46	69.99	919.05	38.59
2012	331.44	59.02	216.90	36.99	114.05	50.60	972.18	103.11
2013	382.35	58.29	229.11	44.37	82.46	48.34	1 137.30	71.98
2014	308.83	62.11	197.85	36.27	86.39	37.60	1 102.47	116.29
2015	370.41	59.68	198.40	36.64	129.79	55.39	1 058.04	37.64
2016	351.55	64.46	208.34	36.04	133.40	53.57	1 022.86	104.30
2017	373.22	62.38	222.59	38.83	101.68	55.26	987.03	164.52
2018	382.34	62.33	278.41	39.41	128.81	61.32	902.15	178.42
2019	338.96	59.50	264.18	32.22	51.23	54.02	880.33	30.88
2020	347.89	53.37	353.79	26.02	119.10	60.63	740.50	72.85
2021	361.81	60.88	383.68	29.46	126.95	67.35	973.54	84.38
2022	223.45	24.74	331.76	38.41	142.58	62.01	838.27	NO
2023	208.87	16.50	286.28	41.68	105.00	71.02	900.69	NO

IE - included in coking coal

## 3.5. Fugitive Emissions from Fuels (CRT 1.B)

### 3.5.1. Overview of Fugitive Emissions from Fuels

Fugitive emissions from the categories 1.B.1 - Solid Fuel and 1.B.2 - Oil and Natural Gas are important sources of methane emissions in the national GHGs inventory. Fugitive methane emissions from charcoal production and coke production are included in the category 1.B.1.b – Solid Fuel Transformation.

In 2023, total aggregated fugitive emissions in the category 1.B represented 645.53 Gg of CO<sub>2</sub> eq. Overview of the total GHG emissions reported in the category 1.B is provided in [Table 3.1](#) and tier used is provided in [Table 3.2](#). Methane emissions from abandoned underground mines (category 1.B.1.a.1.iii) are reported in the inventory since 2015. [Tables 3.46](#) and [3.47](#) summarize emissions according to the most significant categories within 1.B in particular years. GHG emissions from the activities occurring in the category 1.B.2.a.5 – Distribution of Oil Products are not estimated because of the 2006 IPCC Guidelines and also 2019 Refinement to the 2006 IPCC Guidelines do not include methodologies to estimate them, therefore the notation key “NE” is used here.

The trend is steadily decreasing as an outcome of introduction of new technologies, methodologies and closing the coal mines. Fugitive emissions from the transport and distribution of fossil fuels (oil and natural gas) are significant because Slovakia is an important transit country for oil and natural gas from East-European countries to the European Union. Raw materials are transported through high-pressure pipelines and distribution network and they are pumped by pipeline compressors (1.A.3.e.i). Trend in fugitive emissions from the transport and distribution of oil and natural gas in the Slovak Republic was stabilized and since 2000 slightly decreased. The increase in the past was caused by the expansion of the distribution system for natural gas and growth of its consumption. Since 2000, fugitive emissions from oil have decreased due to the decrease in production and distribution.

**Table 3.46: GHG emissions by categories within the 1.B.1 - Solid Fuels in particular years**

YEAR	1.B.1.a Coal Mining and Handling					1.B.1.b Solid Fuel Transformation		
	1.B.1.a.i.1		1.B.1.a.i.2	1.B.1.a.i.3		Bio-CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	CO <sub>2</sub>	CH <sub>4</sub>	CH <sub>4</sub>	CO <sub>2</sub>	CH <sub>4</sub>			
	Gg							
1990	19.03	25.14	2.09	0.73	1.05	NO	0.11	NO
1995	21.54	7.46	2.27	1.37	0.62	4.71	0.21	0.0002
2000	21.51	7.09	2.20	1.11	0.48	7.85	0.28	0.0004
2005	20.78	3.50	1.51	6.56	1.58	75.36	2.02	0.0038
2010	19.74	3.18	1.43	7.89	1.84	4.74	0.20	0.0002
2015	19.51	2.63	1.17	7.06	1.38	6.28	0.24	0.0003
2016	18.62	2.43	1.11	18.70	3.56	2.27	0.13	0.0001
2017	21.40	1.90	1.11	24.95	3.50	6.28	0.23	0.0003
2018	18.64	1.31	0.91	33.61	4.00	6.44	0.24	0.0003
2019	17.89	1.64	0.86	23.76	3.33	6.28	0.23	0.0003
2020	12.26	1.18	0.59	20.49	2.96	5.97	0.21	0.0003
2021	13.43	1.26	0.65	23.58	3.35	6.28	0.24	0.0003
2022	10.72	0.65	0.52	40.79	4.46	6.28	0.24	0.0003
2023	9.76	0.46	0.46	38.54	3.63	6.28	0.24	0.0003

**Table 3.47: GHG emissions by categories within the 1.B.2 - Oil and NG and other emissions from energy production in particular years**

YEAR	1.B.2.a OIL			1.B.2.b Natural gas		1.B.2.c Venting and Flaring			
						1.B.2.c.i.2 Gas		1.B.2.c.ii.1 Oil	1.B.2.c.ii.2 Gas
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	CO <sub>2</sub>	CH <sub>4</sub>	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	N <sub>2</sub> O
	Gg							kg	
1990	39.69	0.47	0.0005	17.18	49.56	0.23	23.55	49.00	74.10
1995	33.58	0.44	0.0004	15.85	41.59	0.23	20.62	49.75	57.40
2000	34.49	0.39	0.0005	12.83	32.23	0.21	16.50	39.53	28.90
2005	34.15	0.32	0.0005	13.26	25.92	0.23	14.83	20.77	24.50
2010	32.49	0.26	0.0005	11.34	17.56	0.20	10.51	8.77	15.30
2015	35.27	0.26	0.0005	9.83	12.89	0.17	1.87	6.42	14.30
2016	33.95	0.25	0.0005	10.59	13.03	0.19	1.98	5.60	14.80
2017	32.77	0.24	0.0005	11.13	13.25	0.20	1.73	3.87	14.70
2018	32.16	0.23	0.0005	10.39	12.45	0.19	1.44	3.44	14.20
2019	30.09	0.21	0.0004	11.63	12.24	0.21	1.65	2.91	12.40
2020	37.76	0.25	0.0006	9.69	12.48	0.18	2.00	1.40	10.90
2021	32.43	0.23	0.0005	7.27	13.49	0.13	1.37	3.05	10.90
2022	31.69	0.22	0.0005	4.92	12.37	0.08	0.79	1.43	9.29
2023	30.65	0.21	0.0005	2.57	12.47	0.03	2.67	0.96	8.38

### 3.5.2. Uncertainties and Time-series Consistency

The Approach 1 of uncertainty analysis was performed according to the IPCC 2006 GL. Approach 2 uncertainty estimation was not performed due to lack of input data. Availability of inputs is the most facing issue in these categories. The amount of methane from underground mining is naturally fluctuating. The direct measurements of the CH<sub>4</sub> emissions from the ventilated air are with the ±20% of accuracy depending on the measurement's installation. The repeatability of the measurements increases the accuracy up to ±5%. For the continual measurements during 2 weeks, the uncertainty is in the range of ±10-15%.

---

The emissions inventory of fugitive methane emissions from fuels were revised in the previous years, the chosen emission factors for underground coal mining and handling correspond to the national circumstances in the Slovak mining industry. The important reason for this opinion is an occurrence of brown coal underground mines with mainly non-gaseous system in deep shafts. In addition, new emission factors (IPCC 2006 GL, Table 4.2.4) were used for the entire time series. The methodology in these categories is consistent during time series and across the main types of fuels.

### 3.5.3. Category-specific QA/QC and Verification Process

The verification process in the category 1.B.1 is based on cross-checking of input data from the mining companies and the comparison with the sectoral statistical indicators from the Ministry of Economy of the Slovak Republic and the ŠÚ SR. More information can be find in the [Chapter 3.5.6 \(Figure 3.15\)](#).

The verification process in the category 1.B.2 is based on cross-checking the input data from the supplier companies Nafta, a. s. (oil), Transpetrol, a. s. (oil), Eustream, a. s. (natural gas) and the SPP - Distribution, a. s. (natural gas) with the statistics from the Ministry of Economy of the Slovak Republic and the Statistical Office of the Slovak Republic (ŠÚ SR).

For the inventory preparation and verification of currently used methodology, the fugitive emissions from NG were estimated also with the use of data provided directly by (bottom-up approach):

- Eustream, a. s.; as the company responsible for the transmission and storage of the NG and venting (categories 1.B.2.b.4 and 1.B.2.c.1.ii);
- Slovenský plynárenský priemysel – distribúcia a.s (SPP-Distribution, a. s.); as the in-country distributor of natural gas (NG) reported in the category distribution of NG (1.B.2.b.5);
- Nafta, a. s.; as the exclusive company responsible for oil and NG production in Slovakia.

In this submission, further information on the status of implementation of recommendation is providing. Slovakia after cross-check of activity data, changed reported data from statistical data to plant based data. Each company, except of Transpetrol, a. s., is providing the activity data (production, processing and transport) and also with directly measured emissions of CH<sub>4</sub>. Slovakia used these data to recalculate time-series for most of the fugitive emissions categories.

In this submission a new approach for the category NG distribution category (1.B.2.b.5) was implemented. The SPP-Distribution, a. s., as one of largest contributors to fugitive emissions, provided fugitive emissions from distribution of natural gas. Previously a tier 1 methodology was used for this category based on the amount of distributed natural gas. In this submission a model approach was introduced. As this approach did not disturbed the time-series it was switched to this emission estimation approach. Detailed methodology to this calculation is in the moment available in Slovak language and as it is mentioned in [Chapter 3.5.5](#) am annex with a translation of this methodology is planned.

The background documents are archived by the sectoral experts and in the central archiving system of the SNE at the SHMÚ.

### 3.5.4. Category-specific Recalculations

This chapter describes the recalculations of emissions from fugitive emissions and its subcategories with respect to the previous submission.

[Revision of activity data](#) – Revision of statistical data for charcoal production in 2022.

[Reconstruction of post-meter](#) activity data time-series for the period 1990–2005 was conducted.

[Revision of methodology](#) – Slovakia changed its approach to calculate emissions from coal mining. Change in emissions factors from IEA CIAB to 2019 IPCC Refinements emission factors based on the depth of the mines resulted in recalculation of emissions.



CO<sub>2</sub> emissions from abandoned mines are calculated as ratio of these emissions during mining, thus also these emission were affected. Also the methane emission from abandoned mines were revised and corrected (more details in [Chapter 3.5.6](#)).

Missing emissions from N<sub>2</sub>O in the category 1.B.2.c due to error in calculation, were added.

CO<sub>2</sub> recalculations in 1.B - [Table 3.48](#) shows the recalculations of CO<sub>2</sub> emissions for categories of 1.B. Recalculations of CO<sub>2</sub> emissions in 1990 – 2022 are due to implementation of new methodologies, activity data and updated emission factors.

**Table 3.48: Differences in CO<sub>2</sub> emissions between previous submission and current submission caused by recalculations**

YEAR	1.B	1.B.1.a	1.B.2.b
	Gg		
1990	0.05	0.00	0.05
1991	0.05	0.00	0.05
1992	1.16	1.11	0.05
1993	-3.68	0.98	0.05
1994	-3.72	0.94	0.05
1995	-3.74	0.92	0.05
1996	-3.77	0.89	0.05
1997	-3.80	0.86	0.05
1998	-3.85	0.81	0.05
1999	-5.40	0.83	0.05
2000	-7.05	0.75	0.05
2001	-8.59	0.79	0.05
2002	-63.51	0.81	0.05
2003	-67.46	3.15	0.05
2004	-65.92	6.25	0.05
2005	-70.36	4.96	0.05
2006	-112.82	17.49	0.00
2007	-123.26	8.62	0.00
2008	-126.73	6.72	0.00
2009	-119.34	6.26	0.00
2010	1.23	5.97	0.00
2011	-2.06	4.58	0.00
2012	-2.63	4.12	0.00
2013	-2.26	4.02	0.00
2014	-0.78	5.81	0.00
2015	-1.14	5.14	0.00
2016	14.18	16.45	0.00
2017	15.34	21.62	0.00
2018	22.86	29.30	0.00
2019	13.79	20.07	0.00
2020	11.04	17.00	0.00
2021	13.70	19.98	0.00
2022	31.02	37.30	0.00

CH<sub>4</sub> recalculations in 1.B - **Table 3.49** shows the recalculations of CH<sub>4</sub> emissions for categories of 1.B – Fugitive emissions. Recalculations of CH<sub>4</sub> emissions in 1990 – 2022 are due to implementation of new methodologies, activity data and updated emission factors.

**Table 3.49: Differences in CH<sub>4</sub> emissions between previous submission and current submission caused by recalculations**

YEAR	1.B	1.B.1.a	1.B.1.b	1.B.2.b
	Gg			
1990	5.89	0.00	0.00	5.89
1991	5.93	0.00	0.00	5.93
1992	-13.95	-19.88	0.00	5.93
1993	-13.04	-18.98	0.00	5.94
1994	-14.14	-20.10	0.00	5.96
1995	-14.07	-19.98	0.00	5.91
1996	-14.36	-20.24	0.00	5.88
1997	-14.79	-20.66	0.00	5.87
1998	-15.22	-21.08	0.00	5.86
1999	-14.19	-20.05	0.00	5.85
2000	-13.68	-19.53	0.00	5.85
2001	-12.04	-17.84	0.00	5.80
2002	-11.67	-17.51	0.00	5.84
2003	-8.35	-14.21	0.00	5.86
2004	-7.27	-13.16	0.00	5.89
2005	-5.00	-10.83	0.00	5.82
2006	-6.74	-6.74	0.00	0.00
2007	-8.35	-8.35	0.00	0.00
2008	-10.58	-10.58	0.00	0.00
2009	-11.34	-11.34	0.00	0.00
2010	-10.35	-10.35	0.00	0.00
2011	-10.91	-10.91	0.00	0.00
2012	-10.74	-10.74	0.00	0.00
2013	-11.00	-11.00	0.00	0.00
2014	-10.33	-10.33	0.00	0.00
2015	-8.54	-8.54	0.00	0.00
2016	-6.29	-6.29	0.00	0.00
2017	-5.50	-5.50	0.00	0.00
2018	-3.93	-3.93	0.00	0.00
2019	-5.01	-5.01	0.00	0.00
2020	-3.42	-3.42	0.00	0.00
2021	-3.38	-3.38	0.00	0.00
2022	-0.56	-0.56	-0.003	0.00

N<sub>2</sub>O recalculations in 1.B - **Table 3.50** shows the recalculations of N<sub>2</sub>O emissions for categories of 1.B – Fugitive emissions. Recalculations of N<sub>2</sub>O emissions in 1990 – 2022 are due to calculation corrections.

**Table 3.50: Differences in N<sub>2</sub>O emissions between previous submission and current submission caused by recalculations**

YEAR	1.B	1.B.2.c
	kg	
1990	44.35	44.35
1991	31.36	31.36
1992	27.74	27.74

YEAR	1.B	1.B.2.c
	kg	
1993	25.38	25.38
1994	28.94	28.94
1995	34.35	34.35
1996	31.36	31.36
1997	28.94	28.94
1998	25.98	25.98
1999	21.33	21.33
2000	17.31	17.31
2001	19.57	19.57
2002	17.74	17.74
2003	23.42	23.42
2004	16.54	16.55
2005	14.65	14.65
2006	19.40	19.40
2007	12.82	12.82
2008	10.17	10.17
2009	10.30	10.30
2010	9.16	9.16
2011	9.16	9.16
2012	9.23	9.23
2013	9.54	9.54
2014	8.69	8.69
2015	8.56	8.56
2016	8.86	8.86
2017	8.81	8.81
2018	8.52	8.52
2019	7.44	7.44
2020	6.53	6.53
2021	6.52	6.52
2022	5.56	5.56

### 3.5.5. Category-specific Improvements and Implemented Recommendations

Slovakia is preparing annexes to the methodologies used for estimating emissions from natural gas transmission and distribution in future submissions.

### 3.5.6. Solid Fuels (CRT 1.B.1)

**Coal mining and handling (CRT 1.B.1.a)** – 790.00 kt of brown coal was mined from underground mines in the Slovak Republic in 2023, mostly for domestic consumption (energy industry and households). Total methane emissions from the underground coal mining were estimated to be 4.55 Gg (0.46 Gg of CH<sub>4</sub> from mining activities, 0.46 Gg of CH<sub>4</sub> from post-mining activity and 3.63 Gg from abandoned mines) in 2023. Total CO<sub>2</sub> emissions underground mines were estimated to be 48.30 Gg in 2023.

**Table 3.51: Overview of fugitive emissions from mining and post-mining activities in particular years**

YEAR	Brown coal produced	CH <sub>4</sub> emissions from mining	CH <sub>4</sub> recovery from mining	CH <sub>4</sub> emissions from post-mining	CH <sub>4</sub> emissions from abandoned mines	Total CH <sub>4</sub> emissions	CO <sub>2</sub> emissions from mines
	<i>kt</i>	<i>Gg</i>					
1990	3 456.00	25.143	NO	2.086	1.046	28.275	19.761
1995	3 759.10	7.460	NO	2.267	0.617	10.344	22.908
2000	3 649.30	7.090	NO	2.201	0.478	9.768	22.619
2005	2 511.20	3.500	NO	1.514	1.583	6.597	27.342
2010	2 377.53	3.183	NO	1.434	1.845	6.461	27.627
2015	1 939.33	2.634	NO	1.169	1.376	5.179	26.569
2016	1 847.13	2.432	NO	1.114	3.562	7.108	37.318
2017	1 834.00	1.898	NO	1.106	3.502	6.505	46.348
2018	1 502.00	1.314	NO	0.906	4.002	6.221	52.254
2019	1 431.00	1.643	NO	0.863	3.328	5.833	41.654
2020	980.00	1.179	NO	0.591	2.959	4.730	32.755
2021	1 074.00	1.262	NO	0.648	3.353	5.263	37.007
2022	868.51	0.649	NO	0.524	4.461	5.633	51.511
2023	762.00	0.459	NO	0.459	3.631	4.550	48.297

**Solid fuel transformation (CRT 1.B.1.b)** – total CO<sub>2</sub> eq. emissions from this category were 6.62 kt in 2023. Fugitive methane and N<sub>2</sub>O emissions from charcoal production and coke production in the Slovak Republic is reported in this category. Charcoal production is reported in the FAO database since 1993. The production of wood charcoal is included in this category and CH<sub>4</sub> emissions were estimated for the years 1993 – 2023. Total volume of wood charcoal produced in Slovakia was 4 kt in 2023. Total CH<sub>4</sub> emissions were 0.16 Gg in 2023. According to the new 2019 RF methodology it is possible to estimate also bio-CO<sub>2</sub> and N<sub>2</sub>O emission, as well as CH<sub>4</sub> emissions from coke production. CO<sub>2</sub> emissions from coke production are already included in the carbon balance in 1.A.1.c. Total coke production was 1 473 kt in 2023 and producing 0.07 kt of CH<sub>4</sub> emissions in 2023.

### Methodological Issues

**Coal mining and handling (CRT 1.B.1.a)** – Total emissions from fugitive sources in coal mining industry can be calculated by the following formula:

$$\text{CH}_4 = \text{underground mining emissions} + \text{post-mining activity emissions} - \text{recovery or flared methane with cogeneration} + \text{emissions from abandoned mines}$$

The amount of mined brown coal (in the raw form) is the primary activity data. According to the 2019 Refinements to the 2006 IPCC GL (2019 RF), tier 2 and the country specific EFs were used:

- IPCC 2006 Guidelines for National Greenhouse Gas Inventories (IPCC 2006 GL) and also 2019 Refinements to the 2006 IPCC GL, Volume 2: Energy Chapter 4: Fugitive Emissions: 4.1 fugitive emissions from mining, processing, storage and transport of coal. According to the Refinements, the emission factor is dependent on depth differentiation for coal mining and 0.9 m<sup>3</sup> CH<sub>4</sub>/t for post-mining activities.
- Measurements of EF CH<sub>4</sub> as specified by the mines operator - HBP, a. s. The emission factors estimated by the HBP, a.s. mine operator on the base of measured concentration values of the methane in the air ventilation was assigned for one single mine according to the suggestion of the operator. These emission factors are underestimated.

The emission factors for mining and post-mining activities were derived from IPCC 2019 RF Volume 2, Chapter 4 (Fugitives), section 4.1.3.2, pages 4.18 onwards (for mining without drainage with known gas amount). Overview of emission factors is presented in **Table 3.52**.

**Table 3.52: Coal production, characteristics of mines and the emission factors for mining and post-mining in single mines in the Slovak Republic in 2023**

MINE	COAL PRODUCTION	DEPTH OF MINE	EF CH <sub>4</sub>			
			1. 2019 RF		3. HPB, a.s.	
			Mining	Post-mining	Mining	Post-mining
t/year	m					
Mine Nováky	762 000	200	0.603	0.603	0.92	0.39
Mine Nováky 6 <sup>th</sup> logging place	NO	200	0.603	0.603	4.17	0.46
Mine Cigeľ	NO	500	2.680	0.603	0	0
Mine Cigeľ 7 <sup>th</sup> logging place	NO	500	2.680	0.603	4.17	0.46
Mine Handlová	NO	500-1500	2.680	0.603	0	0
Mine Handlová east shaft	NO	500-1500	2.680	0.603	4.17	0.46
Mine Dolina	NO	600	2.680	0.603	0.02	0.01
Mine Čáry	NO	400	1.675	0.603	0.02	0.01

Five localities of underground mines operated by two companies are in the Slovak Republic. Data of coal production from the underground mines were obtained from official sources (ŠÚ SR) and directly from the companies: Hornonitrianske bane Prievidza (HBP) and previously also from Baňa Dolina Veľký Krtíš (BD). According to the Regulation of the Slovak Office of Mines No 21/1988 Coll., mines are differentiated based on gas release as follows:

- HBP, a.s. Prievidza:
  - Mine Cigeľ – non-gaseous (closed in July 2017)
  - Mine Cigeľ 7<sup>th</sup> logging place - gaseous,
  - Mine Handlová – gaseous,
  - Mine Nováky – gaseous,
  - Mine Čáry Holíč – gaseous;
- Baňa Dolina Veľký Krtíš – gaseous (closed).

CH<sub>4</sub> emissions from post-mining activities are presented in mined coal and released into the atmosphere during the manipulation and storage of coal. The measurements of these emissions are not carried out so the emissions are estimated with the default emission factors based on coal mined.

CO<sub>2</sub> emissions estimation from coal mining is based on expert judgement provided by the Department of Ventilation and Drainage of the HBP, a. s. company. Annual quantities of mining winds and average CO<sub>2</sub> concentration are measured as part of the safety protocols. This is an approximation with respect to the accuracy of the measuring instruments for measuring the mining air. The last open mine in 2023 was mine Nováky, which was to the end of the year closed.

**Table 3.53: Overview of CO<sub>2</sub> emission factors for single mines in the Slovak Republic in 2023**

MINE	COAL PRODUCTION	EF	EMISSIONS CO <sub>2</sub>
	t/year	t CO <sub>2</sub> /t	t/year
Mine Nováky	762 000	0.012802	9 755
<b>TOTAL</b>	<b>7</b>	<b>0.012802</b>	<b>9 755</b>

**Abandoned mines (CRT 1.B.1.a1.iii)** – fugitive emissions from this category are calculated based on tier 1 methodology from the 2019 Refinements to the 2006 IPCC Guidelines. Except of the 6<sup>th</sup> logging unit of mine Nováky, all mines were closed in the time interval of 2001-2025. As there is no evidence of sealing these mines, conservative approach is used and all mines are still considered gassing. Summary of closed mines and emissions is in [Table 3.54](#). As the mines were closed in different years in the time band, emission factor for the first year of the time band was used for the first year of the closure.

**Table 3.54: Summary calculation of abandoned mines in 2023**

Abandoned mines	Interval of mine closure				
	1901 – 1925	1926 – 1950	1951 – 1975	1976 – 2000	2001 – 2025
Number of mines closed per time band	NO	NO	NO	1	6
Fraction of gassy mines	NO	NO	NO	1.0	1.0
<b>TOTAL CH<sub>4</sub> (kt)</b>	NO	NO	NO	0.269	3.362

**Solid Fuel Transformation (CRT 1.B.1.b)** – fugitive emission from solid fuel transformation have been calculated by the IPCC tier 1 default approach with using 2019 Refinements to the 2006 IPCC Guidelines. This category includes fugitive emissions from charcoal, biochar and coke production. The GHG emissions from charcoal and coke combustion are included in the Energy sector, where the activity data represents the quantity of production excluding export.

Production of charcoal and coke in Slovakia were obtained from the official FAO statistic for charcoal and the Statistical office of the Slovak Republic for coke (data used in 1.A.1.c category). A higher production of charcoal was recognised in years 2002 – 2009. This issue was also consulted with the Ministry of Agriculture of the Slovak Republic (responsible for FAOSTAT) but it was not possible to reconstruct the reasons of this trend. CO<sub>2</sub> emissions occur only in charcoal production and are considered as biomass origin, thus should be reported as memo items (reported in the documentation box in CRT). CO<sub>2</sub> emissions from coke production are based on the carbon content are balanced and reported in the Energy sector under the EU ETS. There is no biochar production in Slovakia.

**Table 3.55: Charcoal and coke production and fugitive emissions in particular years**

YEAR	Charcoal production	Coke production	Bio-CO <sub>2</sub> emissions	CH <sub>4</sub> emissions	N <sub>2</sub> O emissions
	Gg/year	Gg/year	Gg/year	Gg/year	Gg/year
1990	0.00	2 340.00	0.00	0.11	0.0000
1995	3.00	1 854.00	4.71	0.21	0.0002
2000	5.00	1 596.92	7.85	0.28	0.0004
2005	48.00	1 740.00	75.36	2.02	0.0038
2010	3.02	1 550.01	4.74	0.20	0.0002
2011	4.23	1 520.01	6.64	0.24	0.0003
2012	4.30	1 470.01	6.75	0.25	0.0003
2013	4.00	1 440.01	6.28	0.23	0.0003
2014	4.20	1 470.01	6.59	0.24	0.0003
2015	4.00	1 530.01	6.28	0.24	0.0003
2016	1.45	1 540.01	2.27	0.13	0.0001
2017	4.00	1 490.01	6.28	0.23	0.0003
2018	4.10	1 500.01	6.44	0.24	0.0003
2019	4.00	1 320.01	6.28	0.23	0.0003
2020	3.80	1 110.00	5.97	0.21	0.0003
2021	4.00	1 626.00	6.28	0.24	0.0003
2022	4.00	1 450.00	6.28	0.23	0.0003
2023	4.00	1 473.00	6.28	0.23	0.0003

## Source specific recalculations

Recalculations are described in the [Chapter 3.5.4](#).

### 3.5.7. Oil and Natural Gas and Other Emissions from Energy Production (CRT 1.B.2)


The production of oil and natural gas from domestic sources is negligible in the Slovak Republic and the major share of these stocks comes from import. Fugitive methane emissions from natural gas distribution and venting are key categories in level assessment. Total CH<sub>4</sub> emissions represented 429.85 Gg of CO<sub>2</sub> eq. (15.35 Gg of CH<sub>4</sub>) in 2023. Total CO<sub>2</sub> emissions were 33.25 Gg in 2023. Total N<sub>2</sub>O emissions were 0.46 t in 2023. The major share of emissions belongs to the NG post meter emissions (49.35%), NG distribution (26.64%) and oil refining and storage (9.03%). Production of natural gas has stabilised in 2023 and represented 2.13% from the total fugitive emissions from oil and NG activities.

Total fugitive GHG emissions from oil activities (1.B.2.a) were 36.73 Gg of CO<sub>2</sub> eq. (3.65 kt of CO<sub>2</sub> and 0.21 kt of CH<sub>4</sub>) in 2023. Total GHG emissions are decreasing continuously due to decrease in production and storage.

**Table 3.56:** Trend in fugitive emissions from oil activities in particular years

YEAR	1.B.2.a OIL									
	1.B.2.a.2 Production			1.B.2.a.3 Transport			1.B.2.a.4 Refining/Storage			
	Production	Emissions		Transport	Emissions		Refining/Storage	Emissions		
	kt	t CO <sub>2</sub>	t CH <sub>4</sub>	kt	t CO <sub>2</sub>	t CH <sub>4</sub>	kt	kt CO <sub>2</sub>	t CH <sub>4</sub>	t N <sub>2</sub> O
1990	73.14	3 290.43	212.83	13 581.00	6.65	73.34	6 221.14	36.39	186.63	0.54
1995	74.25	3 340.37	216.06	13 581.00	6.14	67.66	5 168.47	30.24	155.05	0.45
2000	59.00	2 654.41	171.69	9 300.00	4.56	50.22	5 442.00	31.84	163.26	0.47
2005	31.00	1 394.69	90.21	10 662.34	5.22	57.58	5 598.00	32.75	167.94	0.49
2010	13.08	588.60	38.07	10 075.33	4.94	54.41	5 453.00	31.90	163.59	0.47
2015	9.59	431.27	27.90	9 932.04	4.87	53.63	5 954.53	34.83	178.64	0.52
2016	8.36	376.03	24.32	9 171.32	4.49	49.53	5 738.02	33.57	172.14	0.50
2017	5.78	259.82	16.81	9 582.25	4.70	51.74	5 557.00	32.51	166.71	0.48
2018	5.14	231.25	14.96	9 460.16	4.64	51.08	5 457.49	31.93	163.72	0.47
2019	4.34	195.12	12.62	8 997.64	4.41	48.59	5 109.01	29.89	153.27	0.44
2020	2.09	94.03	6.08	9 974.83	4.89	53.86	6 437.93	37.66	193.14	0.56
2021	4.56	205.02	13.26	8 819.00	4.32	47.62	5 507.00	32.22	165.21	0.48
2022	2.14	96.14	6.22	9 595.06	4.70	51.81	5 400.00	31.59	162.00	0.47
2023	1.43	64.25	4.16	9 626.21	4.72	51.98	5 227.00	30.58	156.81	0.45

**Figure 3.15:** The share of individual activities in fugitive emissions of oil and natural gas in 2023



Total fugitive GHG emissions from natural gas activities (1.B.2.b) were 351.77 Gg of CO<sub>2</sub> eq. (2.57 Gg of CO<sub>2</sub> and 12.47 Gg of CH<sub>4</sub>) in 2023. Despite the expansion of the distribution system, the trend of the fugitive emissions from distribution of the natural gas in the Slovak Republic is decreasing in line with the decrease of natural gas transit.

**Table 3.57:** Trend in fugitive emissions from NG activities in particular years

YEAR	1.B.2.b NATURAL GAS								
	1.B.2.b.2 Production			1.B.2.b.3 Processing			1.B.2.b.4 Transmission and storage		
	Product.	Emissions		Process.	Emissions		Transfer	Emissions	
	mil m <sup>3</sup>	t CO <sub>2</sub>	t CH <sub>4</sub>	mil m <sup>3</sup>	t CO <sub>2</sub>	t CH <sub>4</sub>	mil m <sup>3</sup>	t CO <sub>2</sub>	t CH <sub>4</sub>
1990	444.00	1 753.80	2 548.56	444.00	4 195.80	333.00	73 600.00	11 040.00	35 328.00
1995	344.00	1 358.80	1 974.56	344.00	3 250.80	258.00	73 600.00	11 040.00	27 968.45
2000	173.00	683.35	993.02	173.00	1 634.85	129.75	68 600.00	10 290.00	19 208.84
2005	147.00	580.65	843.78	147.00	1 389.15	110.25	73 900.00	11 085.00	13 303.35
2010	94.03	361.99	526.04	94.03	866.03	68.73	65 302.00	9 795.30	5 225.75
2015	84.57	338.22	491.49	84.57	809.16	64.22	55 800.00	8 370.00	1 392.80
2016	87.89	350.33	509.09	87.89	838.14	66.52	60 600.00	9 090.00	1 450.61
2017	87.29	347.29	504.67	87.29	830.86	65.94	64 200.00	9 630.00	1 465.26
2018	84.15	334.76	486.46	84.15	800.88	63.56	59 700.00	8 955.00	866.75
2019	73.60	292.57	425.15	73.60	699.94	55.55	69 060.00	10 359.00	809.90
2020	65.26	257.87	374.73	65.26	616.93	48.96	56 980.00	8 547.00	1 059.67
2021	65.33	258.07	375.02	65.33	617.40	49.00	40 362.00	6 054.24	1 197.89
2022	55.61	219.66	319.21	55.61	525.52	41.71	25 772.86	3 865.93	1 110.91
2023	50.21	198.32	288.19	50.21	474.47	37.66	10 940.17	1 641.03	790.73

YEAR	1.B.2.b NATURAL GAS					
	1.B.2.b.5 Distribution			1.B.2.b.6 Other		
	Distribution	Emissions		Storage	Emissions	
	mil m <sup>3</sup>	t CO <sub>2</sub>	t CH <sub>4</sub>	mil m <sup>3</sup>	t CO <sub>2</sub>	t CH <sub>4</sub>
1990	6 666.00	133.32	4 132.92	1.00	0.04	0.29
1995	6 485.00	129.70	4 020.70	159.40	6.38	46.23



YEAR	1.B.2.b NATURAL GAS					
	1.B.2.b.5 Distribution			1.B.2.b.6 Other		
	Distribution	Emissions		Storage	Emissions	
	<i>mil m<sup>3</sup></i>	<i>t CO<sub>2</sub></i>	<i>t CH<sub>4</sub></i>	<i>mil m<sup>3</sup></i>	<i>t CO<sub>2</sub></i>	<i>t CH<sub>4</sub></i>
2000	7 136.00	142.72	4 424.32	524.30	20.97	152.05
2005	7 399.00	147.98	4 587.38	50.00	2.00	14.50
2010	6 098.00	121.96	3 780.76	3 435.21	137.41	996.21
2015	4 639.00	92.78	2 876.18	4 017.26	160.69	1 165.01
2016	4 716.00	94.32	2 923.92	3 969.67	158.79	1 151.20
2017	4 901.25	98.02	3 038.77	4 246.87	169.87	1 231.59
2018	4 777.99	95.56	2 962.35	3 724.15	148.97	1 080.00
2019	4 841.46	96.83	3 001.70	3 129.80	125.19	907.64
2020	5 003.88	100.08	3 102.40	2 783.82	111.35	807.31
2021	5 471.00	109.42	3 392.02	4 368.00	174.72	1 266.72
2022	4 463.69	89.27	2 767.49	3 997.00	159.88	1 159.13
2023	4 179.08	83.58	3 693.99	2 815.00	112.60	816.35

The IPCC 2019 Refinements also introduced new sources of fugitive emissions. These sources are poste-meter emissions from using CNG vehicles, appliances in households and services and fugitive emissions from industrial plants, where natural gas is combusted. The share of these emissions on the total fugitive emissions from oil and natural gas was 49.35% (191.71 CO<sub>2</sub> eq.). Overview of these emissions is summarized in the following table.

**Table 3.58: Trend in fugitive emissions from other activities in particular years**

YEAR	1.B.2.d Other (Poste-meter emissions)								
	CNG cars	CO <sub>2</sub>	CH <sub>4</sub>	Appliance	CO <sub>2</sub>	CH <sub>4</sub>	Industrial plants	CO <sub>2</sub>	CH <sub>4</sub>
	No.	tons		No.	tons		<i>mil. m<sup>3</sup></i>	tons	
1990	NO	NO	NO	1 472 938	48.61	5 891.75	3 319.38	10.95	1 327.75
1995	NO	NO	NO	1 477 188	48.75	5 908.75	3 528.17	11.64	1 411.27
2000	40	0.0001	0.12	1 461 904	48.24	5 847.62	3 692.88	12.19	1 477.15
2005	158	0.0004	0.47	1 455 729	48.04	5 822.92	3 097.17	10.22	1 238.87
2010	289	0.0007	0.87	1 496 033	49.37	5 984.13	2 453.04	8.10	981.22
2015	1 398	0.0032	4.19	1 510 532	49.85	6 042.13	2 136.90	7.05	854.76
2016	1 541	0.0035	4.62	1 514 666	49.98	6 058.66	2 165.34	7.15	866.13
2017	1 750	0.0040	5.25	1 514 262	49.97	6 057.05	2 214.59	7.31	885.84
2018	1 980	0.0046	5.94	1 519 409	50.14	6 077.64	2 279.11	7.52	911.64
2019	2 063	0.0047	6.19	1 522 827	50.25	6 091.31	2 365.51	7.81	946.20
2020	2 095	0.0048	6.29	1 527 512	50.41	6 110.05	2 416.83	7.98	966.73
2021	2 146	0.0049	6.44	1 529 546	50.48	6 118.18	2 707.13	8.93	1 082.85
2022	2 218	0.0051	6.65	1 532 244	50.56	6 128.98	2 087.94	6.89	835.18
2023	2 033	0.0047	6.10	1 511 931	49.89	6 047.72	1 977.29	6.53	790.91

Total fugitive GHG emissions from flaring and venting activities (1.B.2.c) were 74.71 Gg of CO<sub>2</sub> eq. (0.04 Gg of CO<sub>2</sub>, 2.67 kt of CH<sub>4</sub> and 9.34 kg of N<sub>2</sub>O) in 2023 ([Table 3.59](#)). Major emission decrease is caused by change in the Tier 1 methodology and emissions factors. According to the 2019 RF in most categories are now in the emissions factors also included emissions from venting and flaring of oil and natural gas. Separately are reported only directly measured emission (tier 3) and emissions with no new emission factors. In 2023 there was a high amount of vented gas caused by depressurization of one of

the lines due to an emergency gas leak. The amount of natural gas released during the emergency depressurization was 3 149 733 m<sup>3</sup> (at 20°C, 101.325 kPa).

**Table 3.59: Trend in fugitive emissions from venting and flaring activities in particular years**

YEAR	1.B.2.c.1 VENTING		1.B.2.c.2 Flaring	1.B.2.c.2 Flaring
	1.B.2.c.1.ii Gas		1.B.2.c.2.i Oil	1.B.2.c.2.ii Gas
	CO <sub>2</sub> (t)	CH <sub>4</sub> (t)	N <sub>2</sub> O (t)	N <sub>2</sub> O (t)
1990	228.16	23 552.00	0.049	0.074
1995	228.16	20 624.96	0.048	0.057
2000	212.66	16 495.62	0.040	0.029
2005	229.09	14 831.10	0.021	0.025
2010	202.44	10 508.53	0.009	0.015
2015	172.98	1 868.81	0.006	0.014
2016	187.86	1 984.26	0.006	0.015
2017	199.02	1 731.17	0.004	0.015
2018	185.07	1 442.56	0.003	0.014
2019	214.09	1 648.17	0.003	0.012
2020	176.64	2 003.84	0.001	0.011
2021	125.12	1 373.36	0.003	0.011
2022	79.90	790.46	0.001	0.009
2023	33.91	2 667.02	0.001	0.008

### Methodological Issues

The fugitive emissions from oil and natural gas in the Slovak Republic were calculated according to the IPCC 2019 Refinements to the IPCC 2006 GL using default tier 1 approach.

Emissions from NG transition and storage (fugitive and venting) were calculated using the OGMP 2.0 methodology (Oil and Gas Methane Partnership) on tier 4 approach, which is complementary with the IPCC tier 3 approach. Combination of direct measurements and modelling was used. The calculation were made by Eustream, a. s. and afterwards analysed and verified by the national expert. Throughout description of the methodology is available in Slovak language. This data provided the base for recalculation of the whole time-series of NG transmission. Trend analysis and calculation was used to back-recalculated emissions to the base year 1990. Since the year 2013, direct emissions measurements based on the data from the Eustream, a. s. company are reported. These data are in line with official reports of the company to the other national or international organisations.

Eustream, a. s. uses plant specific methodology for emissions estimation as fugitive emissions from compressors, accidents and planned repairs. Specific compressor stations and transmission system is described on the webpage of Eustream, a.s.<sup>17</sup> (according to the ERT recommendation E.6 based on the draft SVK ARR 2022 delivered on 28<sup>th</sup> February 2023). To monitor each of these possibilities infrared cameras are used.

Fugitive emissions from natural gas distribution in 2023 were estimated using the SIMONE model. SIMONE software is a dynamic modelling tool for gas flow in pipeline systems. Developed by SIMONE Research Group s.r.o., established in 1995, it combines academic research expertise with practical experience in gas transport. Widely adopted, SIMONE is used by over 350 applications in 78 gas

<sup>17</sup> Eustream, a. s. transmission system: <https://www.eustream.sk/en/transmission-system/grid-information-map-transmission-network/mapa-prepravneho-systemu/>

---

companies and 9 universities across Europe. The software is ISO 9001:2015 certified. The model is run by SPP-distribúcia a.s. The outputs of the SIMONE model were consistent with the established emissions time series, enabling Slovakia to transition seamlessly in 2023 to reporting fugitive emissions from natural gas distribution based on this model. Details of calculations are available in Slovak language.

#### **Source specific Recalculations**

Recalculations are described in the [Chapter 3.5.4](#).

## **3.6. International Bunker Fuels (CRT 1.D.1)**

International bunkers category includes emissions from the International Aviation (1.D.1.a) and International Navigation (1.D.1.b). These emissions are excluded from the national totals. This Document uses the GWP 100 based on IPCC Assessment report 5 for the year 2022. The difference between emission based on GWP 100 IPCC Assessment report 4 (AR4) and 5 (AR5) are shown in previous SVK NIR 2023.

### **3.6.1. International Aviation (CRT 1.D.1.a)**

Since 1990 to 2004, the Slovak Republic has been estimating the emissions from the international aviation based on the information provided by the airports about LTO cycles and fuel consumption. The expert approach was used for processing of activity data and disaggregation of fuels into national and international flights in the previous submissions. In this submission, the share was intended as constant value for the years 1990 – 2004 based on trend in years 2005 – 2021. Based on the national circumstances (size of country), the international aviation occurs more frequently than the national aviation. EUROCONTROL data was used in this submission for time series 2005 – 2023, data on the emissions, fuel consumption and division of domestic and international flights.

The GHG emissions estimation was performed based on the fuels sold at the Slovak airports (Bratislava, Košice, Poprad, Sliač, and Žilina) in the period 1990 – 2004. In 2023, the emissions in the international civil aviation represented 152.80 Gg of CO<sub>2</sub> eq. The decrease of emissions after 2008 is explained by the recession of economy and cancelling of many regular flights operated by the foreign companies at Bratislava airport. In recent years, the international aviation begins its rise back to pre-2008 emissions as Bratislava and Košice are a base for low-cost companies (WizzAir, Ryanair, Flydubai, and Eurowings) as well as Austrian Airlines. The major decrease of emissions in 2020 is caused by the COVID pandemic and cancelation of many regular flights. Methodology for emissions estimation in this category is consistent with the methodology used in the domestic aviation and is described in the [Chapter 3.2.8](#) of this Document.

The Slovak Republic has used a tier 1 based on fuel sold, both for aviation gasoline and jet kerosene for 1990 – 2004. In the previous submissions, there were used expert judgment on the sharing of domestic and international flights. According to previous recommendations, the share between domestic and international aviation for the years 1990 – 2004 was estimated by using the trend for the years 2005 – 2021 from the available EUROCONTROL data. The changes are shown in [Chapter 3.2.8](#). The emission factors of all gases were changed for jet kerosene and aviation gasoline and information is provided in the [Chapter 3.2.8](#) of this Document.

New EUROCONTROL data published in 2022 were used for emissions' estimation of aviation transport for time series 2005 – 2020. The decision follows an analysis of the national data and data obtained from EUROCONTROL and approved by the Ministry of Transport and Construction of the Slovak Republic. Aggregated national fuel and emissions balance was calculated using a tier 3 applying the Advanced Emissions Model (AEM) by EUROCONTROL.

Considering comparison between the EUROCONTROL results and national data on fuel consumption, emissions and implied emission factors, the following data were considered (taken from EUROCONTROL results) more accurate and reliable for 2025 inventory preparation:

- calorific values for fuels;
- fuel consumption of aviation gasoline for domestic flights;
- fuel consumption of aviation gasoline for international flights;
- jet kerosene for domestic flights;
- jet kerosene for international flights;
- CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions for all subcategories.

The overview of the international aviation fuels consumption according to the type (aviation gasoline and jet kerosene) is presented in [Table 3.60](#). For the period 1994 – 2004, data were obtained directly from the airports' statistics on annual basis. For the period 1990 – 1993, data were based on expert judgment according to the real LTO cycles in this period. To ensure consistency over time series, NCVs of fuels were used from EUROCONTROL data. Total consumption of jet kerosene was 2 082.83 TJ and total consumption of aviation gasoline was 2.06 TJ in international flights in 2023.

**Table 3.60: Fuels consumption and GHG emissions in international flights in particular years**

YEAR	AVIATION GASOLINE				JET KEROSENE			
	CONSUMPTION	EMISSIONS			CONSUMPTION	EMISSIONS		
	TJ	t CO <sub>2</sub>	t CH <sub>4</sub>	t N <sub>2</sub> O	TJ	t CO <sub>2</sub>	t CH <sub>4</sub>	t N <sub>2</sub> O
1990	7.82	552.964	0.004	0.016	914.43	66 523.27	0.632	1.808
1995	5.18	365.913	0.002	0.010	652.78	47 488.40	0.451	1.290
2000	5.96	421.562	0.003	0.012	644.94	46 918.05	0.446	1.275
2005	1.93	136.798	0.001	0.004	1 914.83	139 300.37	1.350	3.785
2010	2.09	147.709	0.001	0.004	1 814.71	132 016.84	1.269	3.588
2015	2.19	154.854	0.001	0.004	1 982.76	144 242.52	1.334	3.920
2016	3.64	253.476	0.002	0.007	2 113.08	153 722.75	1.493	4.177
2017	1.80	127.088	0.001	0.003	2 260.82	164 889.54	1.581	4.481
2018	1.87	131.574	0.001	0.004	2 527.74	184 357.20	1.777	5.009
2019	1.56	109.722	0.001	0.003	2 543.38	185 497.54	1.795	5.041
2020	1.16	80.152	0.001	0.002	750.40	54 590.51	0.574	1.483
2021	1.94	134.390	0.001	0.004	894.18	65 050.29	0.666	1.768
2022	1.72	119.177	0.001	0.003	1 793.50	130 474.11	1.269	3.546
2023	2.06	142.688	0.001	0.004	2 082.83	151 522.40	1.59	4.12

### Source specific Recalculations

In the NID 2025 there were no category specific recalculations made.

### 3.6.2. International Navigation (CRT 1.D.1.b)

GHG emissions inventory in international navigation transport includes CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from shipping activities in the Danube River. The consumption of diesel oil is determined indirectly by available statistical data on shipping activities of transit in the Slovak part of the Danube River during the year and the technical parameters of the Danube traction vessels. Total aggregated emissions from inland shipping included in international navigation reached 18.17 Gg of CO<sub>2</sub> eq. in 2023. The decrease is significant in comparison with the base year but the inter-annual fluctuations are visible also in recent years. The Slovak Republic used tier 1 approach based on the IPCC 2006 GL. The emissions of greenhouse gases are calculated from the consumed fuel by diesel motor boats multiplied by emission factor. The country specific NCVs were used to convert the quantity of fuel consumption in energy units. The NCVs for diesel fuel blended are shown in the [Chapter 3.2.8](#) of this Document. The emission factors

were taken from the IPCC 2006 GL and GHG emissions were recalculated for time series. Emission factors used in category 1.A.3.d and 1.D.1.b are identical and shown in [Table 3.61](#).

**Table 3.61:** *The default emission factors in kg/TJ used in navigation for time series*

PARAMETER	EMISSIONS FACTORS	
	DOMESTIC NAVIGATION	INTERNATIONAL NAVIGATION
EMISSIONS	<i>kg/TJ</i>	
CO <sub>2</sub>	74 196.78	74 196.78
CH <sub>4</sub>	7	7
N <sub>2</sub> O	2	2


The Slovak Shipping and Ports Company provided detailed information on diesel oil consumption on the Danube River. The consumption is allocated between national and international companies. It was assumed that total fuel sold to international companies is reported in the memo items category (1.D.1.b) and total fuel sold to national companies (Slovak Water Management Enterprise) is reported in the domestic navigation (1.A.3.d). This activity represents movements of ships between Slovak ports (Bratislava, Devín and Komárno cities). This approach was introduced in 2005 and the reallocation of fuels led to the reallocation between categories 1.A.3.d and 1.D.1.b. The GHG emissions from diesel oil sold to international transport in the important Slovak ports Bratislava and Komárno were balanced is shown in [Table 3.62](#).

**Table 3.62:** *GHG emissions balance of diesel oil sold for shipping companies in particular years*

YEAR	CONSUMPTION		EMISSIONS			
	<i>t/year</i>	<i>TJ</i>	<i>t of CO<sub>2</sub></i>	<i>t of CH<sub>4</sub></i>	<i>t of N<sub>2</sub>O</i>	<i>t of CO<sub>2</sub> eq.</i>
1990	20 500.00	871.48	64 576.6	6.10	1.74	65 209.25
1995	18 066.00	760.14	56 326.7	5.32	1.52	56 878.60
2000	NO	NO	NO	NO	NO	NO
2005	212.70	8.98	665.2	0.06	0.02	671.76
2010	10 450.21	441.19	32 692.0	3.09	0.88	33 012.26
2015	7 008.90	295.38	21 887.4	2.07	0.59	22 101.80
2016	6 006.47	253.08	18 753.9	1.77	0.51	18 937.59
2017	5 917.84	249.30	18 473.2	1.75	0.50	18 654.19
2018	3 482.93	146.67	10 868.4	1.03	0.29	10 974.93
2019	5 063.74	213.24	15 793.7	1.49	0.42	15 948.49
2020	4 761.25	200.38	14 838.70	1.40	0.40	14 984.18
2021	5 457.75	229.70	17 093.03	1.61	0.46	17 259.80
2022	5 562.75	235.17	17 421.88	1.65	0.47	17 592.61
2023	5 745.50	242.52	17 994.23	1.70	0.49	18 170.30

The sources of activity data for the period 1994 – 2023 are the Slovak Shipping and Ports Company in Bratislava, the State Shipping Administration and other international shipping companies operated in Slovakia in accordance with the annually provided statistical information in water transport. The activity data for the period 1990 – 1993 are not statistically documented so the expert judgment was performed on the base of the shipping traffic on the Danube River. The emissions for the year 2000 were estimated to be negligible, because of increasing prices of diesel oil in the Slovak Republic and decreasing prices of fuels in the neighbouring countries (market discrepancies).

**Figure 3.16:** Overview of diesel / oil consumption (TJ) for shipping transport in 1990 – 2023



**Source specific recalculations**

In the NID 2025, there were no category specific recalculations made.

<b>CHAPTER 4. IPPU (CRT 2)</b> .....	<b>128</b>
4.1. Overview of the Sector .....	128
4.2. Overall Trends in Industrial Processes.....	130
4.3. Uncertainty Analyses .....	132
4.4. Sector-specific QA/QC and Verification Processes.....	133
4.5. Sector-specific Recalculations .....	136
4.6. Sector-specific Improvements and Implementation of Recommendations .....	137
4.7. Mineral Products (CRT 2.A).....	137
4.8. Chemical Industry (CRT 2.B) .....	148
4.9. Metal Production (CRT 2.C) .....	161
4.10. Non-energy Products from Fuels and Solvent Use (CRT 2.D).....	174
4.11. Electronic Industry (CRT 2.E) .....	181
4.12. Product Uses as Substitutes for ODS (CRT 2.F).....	181
4.13. Other Product Manufacture (CRT 2.G).....	204
4.14. Other Production (CRT 2.H) .....	209
<b>ANNEX 4.1. CO<sub>2</sub> REFERENCE APPROACH AND COMPARISON WITH THE SECTORAL APPROACH, AND RELEVANT INFORMATION ON THE NATIONAL ENERGY BALANCE .....</b>	<b>210</b>
<b>ANNEX 4.2. METHODOLOGY OF ACQUISITION AND DATA PROCESSING ON F-GASES CONSUMPTION IN THE CATEGORIES 2.F, 2.G.1 AND 2.G.2 .....</b>	<b>213</b>
<b>ANNEX 4.3. BALANCE OF UREA: IMPORT-EXPORT-PRODUCTION-USE BALANCE .....</b>	<b>222</b>

## CHAPTER 4. IPPU (CRT 2)

This Chapter was prepared using GWP<sub>100</sub> taken from the [5<sup>th</sup> Assessment Report of the IPCC](#) by the sectoral experts and institutions involved in the National Inventory System of the Slovak Republic:

INSTITUTE	CHAPTER	SECTORAL EXPERT
Faculty of Chemical and Food Technology, Slovak Technical University	All chapters	Vladimir Danielik
Faculty of Chemical and Food Technology, Slovak Technical University	2.D – NMVOC inventory	Vladimir Danielik
Slovak Hydrometeorological Institute, Department of Emissions and Biofuels	2.D.3 – Urea Based Catalysts	Ján Horváth


### 4.1. Overview of the Sector

The Industrial processes and product use sector includes all GHG emissions generated from the technological processes producing raw materials and products. Within the preparation of the GHG emissions balance in the Slovak Republic, consistent methodology is put on the analysis of individual technological processes and disaggregation between the fuel combustion emissions (in heat and energy production) and emissions from the technological processes and industrial production. In this submission, 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories was fully implemented. Most important emission sources (installations) are balanced separately, and details are explained in [Annexes 4.1-4.2](#) to this Chapter. Emission and oxidation factors were determined, as well as other parameters entering the balancing equations and the results are compared with the verified emissions provided in the EU ETS reports.

In 2023, total aggregated GHG net emissions from the sector of industrial processes and product use were 7 292.59 Gg of CO<sub>2</sub> eq. and they decreased compared with the previous year by approximately 3%. Compared to the base year 1990 the emissions are lower by 23%. CO<sub>2</sub> is the most important gas with the share of 92.1%, followed by F-gases (6.2%) and N<sub>2</sub>O emissions (1.4%) shares. The most important emission sources are categories of metal production (51.6%), mineral products (27.3%), chemical industry (13.2%) and substituents for ODS (6.0%). Other product manufacture and non-energy products categories shares 1.0% and 0.8%, respectively ([Figure 4.1](#)). The most important source of N<sub>2</sub>O emissions are categories Nitric Acid Production and N<sub>2</sub>O from Product Use, which share almost the total amount of N<sub>2</sub>O emissions with the ratio near to 1:1.



**Figure 4.1:** The share on emissions of individual categories in the IPPU sector in 2023



The IPPU sector covers emissions from the technological processes in mineral products industry (CRT 2.A), in chemical industry (CRT 2.B), in metal production (CRT 2.C), in non-energy products from fuels and solvent use (CRT 2.D), in electronics industry (CRT 2.E), in product uses as substitutes for ODS (CRT 2.F) and in other product manufacture (CRT 2.G). The emissions inventory of technological processes includes direct greenhouse gas emissions (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, halocarbons and SF<sub>6</sub>) and indirect greenhouse gas emissions (NO<sub>x</sub>, CO, NMVOCs). List of GHG gases reported in individual categories in 2023 is presented in [Table 4.1](#).

**Table 4.1:** GHG gases reported in the IPPU sector according to the CRT categories in 2023

CATEGORY (CODE AND NAME)	METHODOLOGY/TIER	GHG GASES REPORTED
2.A.1 Cement Production	T2	CO <sub>2</sub>
2.A.2 Lime Production	T2	CO <sub>2</sub>
2.A.3 Glass Production	T3	CO <sub>2</sub>
2.A.4.a Ceramics	T3	CO <sub>2</sub>
2.A.4.b Other Uses of Soda Ash	NO	NO
2.A.4.c Non Metallurgical Magnesia Production	T3	CO <sub>2</sub>
2.A.4.d Other - Limestone for Desulphurization	T3	CO <sub>2</sub>
2.A.5 Other	NO	NO
2.B.1 Ammonia Production	T3	CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O
2.B.2 Nitric Acid Production	T3	N <sub>2</sub> O
2.B.3 Adipic Acid Production	NO	NO
2.B.4 Caprolactam, Glyoxal and Glyoxylic Acid Production	NO	NO
2.B.5 Carbide Production	T2	CO <sub>2</sub>
2.B.6 Titanium Dioxide Production	NO	NO
2.B.7 Soda Ash Production	NO	NO
2.B.8.a Methanol	NO	NO
2.B.8.b Ethylene	T2	CO <sub>2</sub>
2.B.8.c Ethylene Dichloride and Vinyl Chloride Monomer	T2	CO <sub>2</sub>
2.B.8.d Ethylene Oxide	NO	NO
2.B.8.e Acrylonitrile	NO	NO
2.B.8.f Carbon Black	NO	NO
2.B.9 Fluorochemical Production	NO	NO
2.B.10 Other - Hydrogen Production	NO	NO
2.C.1 Iron and Steel Production	T2, T3, T1	CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O

CATEGORY (CODE AND NAME)	METHODOLOGY/TIER	GHG GASES REPORTED
2.C.2 Ferroalloys Production	T3, T2	CO <sub>2</sub> , CH <sub>4</sub>
2.C.3 Aluminium Production	T3, T2, T1	CO <sub>2</sub> , PFCs
2.C.4 Magnesium Production	NO	NO
2.C.5 Lead Production	T1	CO <sub>2</sub>
2.C.6 Zinc Production – not occurring since 2015	T1	CO <sub>2</sub>
2.C.7 Other	NO	NO
2.D.1 Lubricant Use	T1	CO <sub>2</sub>
2.D.2 Paraffin Wax Use	T1	CO <sub>2</sub>
2.D.3 Solvent Use	T2	CO <sub>2</sub>
2.D.4 Other	NO	NO
2.E.1 Integrated Circuit or Semiconductor	NO	NO
2.E.2 TFT Flat Panel Display	NO	NO
2.E.3 Photovoltaics	NO	NO
2.E.4 Heat Transfer Fluid	NO	NO
2.E.5 Other	NO	NO
2.F.1 Refrigeration and Air Conditioning	T2	HFCs: 23, 32, 125, 134a, 143a, 152a, 227ea PFCs: 116
2.F.2 Foam Blowing Agents	T2	HFCs: 134a, 245fa, 365mfc, 227ea
2.F.3 Fire Protection	T1a	HFCs: 134a, 227ea, 236fa
2.F.4 Aerosols	T1a	HFCs: 134a, 227ea
2.F.5 Solvents	NO	NO
2.F.6 Other Applications	NO	NO
2.G.1 Electrical Equipment	T3	SF <sub>6</sub>
2.G.2 SF <sub>6</sub> and PFCs from Other Product Uses	NO	NO
2.G.3 N <sub>2</sub> O from Product Uses	T1	N <sub>2</sub> O
2.G.4 Other	NO	NO
2.H.1 Pulp and Paper Industry	NO	NO
2.H.2 Food and Beverages Industry	NO	NO
2.H.3 Other	NO	NO

## 4.2. Overall Trends in Industrial Processes

Overall trends from numbers provided by the Statistical Office of the Slovak Republic were updated. Energy intensity of industrial processes in the Slovak Republic decreased significantly in comparison with the base year 1990. A decrease in the final energy consumption by 12% was accompanied by an increase in the energy productivity. However, the energy productivity of the IPPU sector in Slovakia is still relatively lower in comparison with the EU average. This has been caused by the historical structure of industrial production. The internal structure of the Slovak industry underwent further changes after accession to the EU. The importance of mining and distribution of electricity, gas and water on production of value added has been significantly reduced and nowadays it is comparable with other developed countries.

The most important indicator is decrease in fuels, electricity and heat consumption in industry in 2023 in comparison with 2005. On the other hand, the increase of renewable energy sources in industry is dominant in recent years. The overview of emission trends in gases and categories is provided in [Tables 4.2](#) and [4.3](#) and [Figures 4.2](#) and [4.3](#).


**Table 4.2: GHG emissions according to the individual gases in the IPPU sector in particular years**

YEAR	CO <sub>2</sub> Emissions	CH <sub>4</sub> Emissions	N <sub>2</sub> O Emissions	HFC, PFC and SF <sub>6</sub>
	Gg of CO <sub>2</sub> eq.			
1990	8 111.12	14.13	1 088.43	213.98
1995	7 825.13	12.78	1 077.71	113.00
2000	7 124.79	14.29	923.06	129.47
2005	8 072.99	17.32	1 173.30	321.46
2010	7 519.11	18.20	842.99	617.72
2011	7 561.32	16.85	426.16	622.50
2012	7 542.59	17.85	337.68	652.59
2013	7 365.60	18.48	225.32	661.00
2014	7 623.71	19.42	201.36	659.02
2015	7 750.13	18.41	186.31	736.12
2016	8 029.87	19.10	171.11	669.12
2017	8 264.89	19.80	156.96	734.24
2018	8 323.96	19.43	157.47	701.44
2019	7 490.44	15.42	140.31	712.10
2020	6 988.93	13.99	126.73	677.60
2021	8 386.76	19.61	115.93	704.04
2022	6 915.91	15.02	111.05	502.15
2023	6 718.99	16.06	104.95	452.60


**Table 4.3: GHG emissions according to the categories in the IPPU sector in particular years**

YEAR	2.A	2.B	2.C	2.D	2.E	2.F	2.G
	Gg of CO <sub>2</sub> eq.						
1990	2 714.02	1 833.81	4 814.71	50.49	NO	NO	14.64
1995	2 070.94	2 137.46	4 720.38	50.49	NO	12.38	36.96
2000	2 230.10	2 045.76	3 735.14	50.49	NO	99.48	30.66
2005	2 532.96	2 219.40	4 434.35	30.17	NO	277.49	90.70
2010	1 941.18	1 761.92	4 619.47	16.94	NO	569.22	89.29
2011	2 359.34	1 586.68	3 994.19	23.90	NO	576.43	86.28
2012	2 116.99	1 265.40	4 432.75	33.55	NO	602.07	99.94
2013	2 030.23	1 216.56	4 229.79	41.14	NO	620.99	131.68
2014	2 181.08	995.33	4 578.94	36.25	NO	626.14	85.77
2015	2 151.36	1 144.04	4 579.71	35.70	NO	704.84	75.32
2016	2 183.45	1 073.85	4 878.50	37.71	NO	647.95	67.74
2017	2 277.13	1 144.35	4 933.31	41.64	NO	710.19	69.26
2018	2 279.54	1 353.00	4 781.31	41.35	NO	675.62	71.48
2019	2 284.96	1 175.92	4 098.19	42.85	NO	688.69	67.68
2020	2 218.73	1 198.98	3 626.61	40.54	NO	646.65	75.74
2021	2 335.45	1 269.22	4 820.21	54.13	NO	672.41	74.92
2022	2 332.71	1 076.42	3 533.08	48.67	NO	480.89	72.38
2023	1 991.90	965.60	3 765.58	60.21	NO	437.89	71.41

**Figure 4.2:** Trend of emissions in the IPPU sector according to individual gases in 1990 – 2023



**Figure 4.3:** Trend of emissions in the IPPU sector according to the categories in 1990 – 2023



### 4.3. Uncertainty Analyses

According to the previous recommendations, Slovakia is using hybrid combination of Approaches 1 and 2 in this submission for calculation of total uncertainty of the inventory ([Annex 3](#) of this Document). Uncertainty analyses performed by the Approach 1 in the IPPU sector were carried out using Table 3.2 for uncertainty calculation and country specific uncertainties for activity data and emission factors were inserted into calculation table.

The Slovak Republic provided and published also Approach 2 for uncertainty analyses according to the Chapter 3 of the IPCC 2006 GL for the complete Energy and IPPU sectors for the year 2015. The methodology and results were described in previous SVK NID 2017 and 2018. Due to the implementation of 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, uncertainty analyses of the IPPU sector were made in 2024 submission using the Monte Carlo simulation. Due to capacity reasons and according to the QA/QC plan in this sector, new calculation of Monte Carlo uncertainty (Approach 2) in the IPPU sector and categories will be performed every five years (next is planned for the year 2027, submission 2029).

Aggregated uncertainty is computed from partial uncertainties. Every category is computed from disaggregated data. The data are split by emission factors or by technological processes. Computed uncertainties are aggregated consecutively to the total uncertainty. The results of every category are generated from 60 000 trials, with random number generator of random numbers for adequate PDF.

---

From theory and knowledge, it is known, that the direct computation of aggregated uncertainty is difficult in many cases. For this reason, a statistical approach has been chosen and the Monte Carlo method is used. It induces the construction of PDF for all input parameters. In some cases, the absence of direct measurement was solved by expert contributions. Mean value and confidence interval have the background usually in measured data or in empirical relations. On the other hand, uncertainty shapes of input parameters are based on following data: (i) uncertainty of data from the EU ETS reports are taken from the criteria presented in the EU ETS reports (uncertainty of scales, of laboratory analysis, etc.); (ii) uncertainty of data that are not covered by the EU ETS reports was assumed as default values from the IPCC 2006 GL; (iii) uncertainties of HFCs in 2.F category and SF<sub>6</sub> in 2.G category were estimated by the sectoral expert for IPPU based on input data provided by the Ministry of the Environment of the Slovak Republic.<sup>1</sup> The results for the IPPU sector and its subsectors following the mentioned assumptions can be seen in the SVK NID 2024.

## 4.4. Sector-specific QA/QC and Verification Processes

The sector-specific QA/QC plan is based on the general QA/QC rules and activities in specific categories. Information used in the process of preparation GHG emissions inventory of the IPPU sector was obtained from the different data sources:

- Statistical Office of the Slovak Republic, Department of Cross-Cutting Statistics (energy balance),
- National Emission Information System (database of all stationary emission sources),
- Emission Trading System (reports from operators and from verifiers),
- Slovak Association for Cooling and Air-conditioning Technology (SZCHKT),
- Questionnaires that were sent to the producers (in case of any doubt).

More information on general QA/QC activities within the National Inventory System of the Slovak Republic is included in the [Chapter 1](#) of this Document.

Input data from producers (providers) are collected by the SNE in cooperation with the sectoral experts and the Slovak University of Technology in Bratislava (Faculty of Chemical and Food Technology). Complete input data related to the production and the quality of products from the previous year is available at the beginning of October (year X+1). The sectoral experts in the cooperation with the Slovak University of Technology in Bratislava (Faculty of Chemical and Food Technology) check and compare obtained information from different data sources during the sectoral inventory preparation in October and November (year X+1). Following QC activities are provided during data collection step:

- comparison with the information provided by the Statistical Office of the Slovak Republic,
- comparison with the information provided by the different associations of producers (if existent),
- comparison with the available information in EU ETS reports (if existent).

Further QC activities during the sectoral inventory preparation:

- outliers checking with developed automated tool,
- comparison of IEFs with the IPCC default EFs and IEFs of neighbouring countries (where available).

---

<sup>1</sup> Based on the Annex III of the Implementing Regulation 749/2014/EU on structure, format and review information pursuant to Regulation (EU) 525/2013, Article 7 (1) (m) (ii)

---

Any discrepancies are directly discussed with subject or data providers. Draft of the sectoral GHG emissions inventory is prepared in the middle of November (year X+1). Quality assurance activities are performed on the draft of the sectoral inventory and the sectoral report. The draft is cross-checked with the independent expert not participating on inventory preparation step from the Slovak University of Technology (independent review) and other experts involved in the NS SR. The independent review is then finished at the end of November and forwarded to the uncertainty analyses. During the application of Monte Carlo model for the uncertainty analyses, the methodology, EFs and other parameters are verified again (mathematically). The final sectoral inventory is prepared at the end of December and it is approved by the NS SR coordinator during the January (year X+2). All original data and protocols are archived at the SHMÚ and in the computers and back-up server of national experts involved in the inventory process.

**Cement Production** - Activity data provided by the Slovak Association of Cement Producers and from the EU ETS reports were verified with the statistical information. Based on the information provided in the EU ETS reports it follows that CO<sub>2</sub> emission was 1 444.72 Gg. All sources reported in this category are included in the EU ETS. The emissions reported in the national inventory were nearly the same (lower by 0.10%). The difference is caused by rounding.

**Lime Production** - Activity data provided by the Slovak Association of Lime Producers and from the EU ETS reports were verified with the statistical information. Sugar plants are not included in official statistics. If any discrepancies occur, the small occasional producers are identified and included in the inventory. Other possible activity data source is the NEIS database. Data there were recorded according to the category of products. In 2023, there were 3 plants included in "others" (2 sugar plants, 1 other plant – production of secondary aluminium). When comparing CO<sub>2</sub> emissions reported in the EU ETS reports and inventory emissions of EU ETS plants, the difference is about 0.4% (higher emissions are in GHG inventory). The difference is caused by rounding of CaO and MgO contents in lime.

**Glass Production** - All sources reported in this category are included in the EU ETS and final emissions are the same as in the GHG inventory.

**Ceramics** - The EU ETS covers all operators reported in this category. CO<sub>2</sub> emissions reported in the EU ETS reports and in the GHG inventory are the same.

**Magnesia Production** - All sources reported in this category are included in the EU ETS. CO<sub>2</sub> emissions reported in the EU ETS reports were 263.64 Gg in 2023 and are nearly the same as in the GHG inventory (+0.004 Gg).

**Other Carbonates** - All sources reported in this category are included in the EU ETS, however, part of them is not calculated but measured. CO<sub>2</sub> emissions calculated in the EU ETS reports were 30.62 Gg in 2023. In the GHG inventory, CO<sub>2</sub> emissions were calculated to be 46.47 Gg, which is in accordance with the EU ETS reports when also measured emissions are considered.

**Ammonia Production** - All sources reported in this category are included in the EU ETS. As ammonia production is one of the largest CO<sub>2</sub> emissions sources and key category (in the **IPPU sector**), a significant attention was paid to validation of activity data and procedures used for the estimation of CO<sub>2</sub> emission in this sector. Basic information on ammonia production and natural gas consumption are provided directly from producer.

Due the subtracting of CO<sub>2</sub> used for urea production, additional QA/QC exercise was performed. Amount of 183.19 Gg of CO<sub>2</sub> was used for the urea production. The CO<sub>2</sub> emissions from the urea consumption were 83.20 Gg in Slovakia (DeNO<sub>x</sub> technologies and using as fertilizers). The difference between these two values (99.99 Gg) is caused by the exporting of urea, because the rest of urea was exported. Based on the data provided by producer approximately 38.94 kt of urea was used for the production of AdBlue (catalyst for vehicles); from which 1.78 kt was exported. This export represents the value of CO<sub>2</sub> as follows: 1.30 kt. Based on the data from the Statistical Office of the Slovak Republic,

---

the urea was exported also under the commodity codes: (i) 31021010, "Urea containing more than 45% by weight of nitrogen on the dry anhydrous product"; (ii) 31028000, "Mixtures of urea and ammonium nitrate in aqueous or ammoniacal solution". Because urea contains 46% of nitrogen, it is clear, that the commodity code 31021010 represents pure urea and export-import difference can be easily calculated from the export and import data. Calculated in this way, the difference between import and export of urea was 5.38 kt of nitrogen in favour of *export*, which represents 8.54 Gg of "exported" CO<sub>2</sub>. Balance of the urea exported/imported under the commodity code 31028000 is much more difficult to estimate. The content of urea in products reported under commodity code 3102800 can vary. According to the announcement of the Ministry of Finance 555/2002 Z. z., the fertilizers with the different content of urea can be used. The nitrogen originating from the urea can be in the range (11-51) %. Because of import data are reported as kilograms of nitrogen, the amount of urea imported to Slovakia was calculated using this range. It follows, that the amount of urea import into Slovakia under the commodity code 31028000 was in the range (1.05-4.87) kt (4.39 kt of nitrogen). According to the data provided by the Slovak fertilizer producer, the fertilizer DAM-390 represents more than 98% export of this commodity. It is the mixture of ammonium nitrate and urea containing 29-30% of N, from which 15.5% of N origins from urea and the rest is from AN. To ensure conservative principle, it can be assumed that 50% of nitrogen origins from urea. Thus, the exported urea under this commodity code represents value 131.69 kt. It results from the balance that the difference between import and export of urea under commodity code 31028000 was (126.82 – 130.64) kt in favour of export, which represents (92.58 – 95.37) Gg of "exported" CO<sub>2</sub>. Balancing of CO<sub>2</sub> from the export/import of urea gives the range (102.42 – 105.21) Gg of "totally exported" CO<sub>2</sub> from Slovakia. Comparing with the value of "missing" CO<sub>2</sub> from the balance of production and use (99.99 Gg) it can be concluded that subtracting of CO<sub>2</sub> used for urea production was made in a correct way. The production/use/import/export balance of urea for the time series 2010 – 2023 is presented in the **Annex 4.3**. Data before 2010 are not available.

**Nitric Acid Production** - Activity data are from the EU ETS reports and information provided by the operators in questionnaires (requested by the SNE) were compared with the measurement's protocols on N<sub>2</sub>O concentration in output gases. All sources reported in this category are included in the EU ETS.

**Carbide Production** - The EU ETS report contains only CO<sub>2</sub> emissions from CaC<sub>2</sub> production no data about using of calcium carbide. Therefore, no comparison with EU ETS can be made, information provided in the separate questionnaires are used.

**Ethylene Production** - Activity data from the EU ETS reports and information provided by the operators in questionnaires (requested by the SNE) were compared. All sources reported in this category are included in the EU ETS.

**Ethylene Dichloride and Vinyl Chloride Monomer** - Activity data are from the EU ETS reports and information provided by the operators in questionnaires (requested by the SNE) are compared. All sources reported in this category are included in the EU ETS.

**Iron and Steel Production** - Specific QA/QC procedure was made for the integrated iron and steel company that represents the biggest source of CO<sub>2</sub> emissions in the **IPPU sector**. The comparison of two independent emission estimations was evaluated. The EU ETS reports contain information on CO<sub>2</sub> emissions. These results were compared with the results obtained by the carbon balance prepared and presented in the **Chapter 4.9.1** and in the **Annex 4.1** of this Document. The difference between CO<sub>2</sub> emissions calculated from these two sources is 0.06% in 2023.

**Ferrous Alloys Production** - Activity data are compared with the information from the ŠÚ SR (ferrous alloys production). Another source used for verification is the U.S. Geological Survey. Data for the period 1990 – 2011 were available and were compared with the results of the national GHG emissions inventory. The consistency of time series was verified.

**Aluminium Production** - Activity data and emissions were verified with the theoretical thermodynamic calculation provided at the Slovak University of Technology, Faculty of Chemical and Food Technology together with comparison with the EU ETS report. All sources of aluminium production in Slovakia are covered with the EU ETS.

**Lead Production** - This production is not covered by the EU ETS, therefore data was provided directly by the operators.

**Non-Energy Products from Fuels and Solvents Use** - This category is not covered by the EU ETS, the data were obtained from the special questionnaires of the ŠÚ SR. Due to the lack of appropriate statistical information and methodological advises in the IPCC 2019 GL, inputs were taken directly from the estimations of the NMVOC emissions reported under the CLRTAP submission (see **Chapter ES.5**). Total NMVOC emissions from solvent use, road paving with asphalt, asphalt roofing and asphalt blowing were estimated in the frame of the National Program for Emissions Reduction of Non-Methane Volatile Organic Compounds in the Slovak Republic.

QA/QC activities and verification process for F-gases is provided in the **Chapter 4.12.6** of this Document.

## 4.5. Sector-specific Recalculations

Recalculation in the CO<sub>2</sub> emissions from the urea consumption in cars was made in IPPU sector for years 2013 – 2022 in this submission. Software update of the COPERT model resulted in the corrections to several emission factors, and the addition of new vehicle categories. More details can be found in the energy sector in transport categories. The impact of the recalculation on the 2.D.3 category and whole IPPU sector is presented in the **Tables 4.4** and **4.5**, respectively.

**Table 4.4:** The comparison of CO<sub>2</sub> emissions estimates from urea consumption for the time series 2013 – 2022

YEAR	SUBMISSION 2024		SUBMISSION 2024	Changes in 2.D.3
	Gg CO <sub>2</sub>			
2013	6.052		6.090	0.6%
2014	6.421		6.504	1.3%
2015	6.073		6.315	4.0%
2016	8.549		8.767	2.5%
2017	8.981		10.669	18.8%
2018	9.539		10.571	10.8%
2019	8.807		16.696	89.6%
2020	8.258		18.951	129.5%
2021	9.695		30.009	209.5%
2022	10.077		17.979	78.4%

**Table 4.5:** The impact of the above-mentioned recalculation on the IPPU sector for the time series 2013 – 2022

YEAR	SUBMISSION 2024		SUBMISSION 2025		2023/2024	
	CO <sub>2</sub> emissions	Total emissions	CO <sub>2</sub> emissions	Total emissions	CO <sub>2</sub> emissions	Total emissions
	Gg CO <sub>2</sub> eq.				%	
2013	7 365.56	8 270.36	7 365.60	8 270.40	0.00%	0.00%
2014	7 623.63	8 503.43	7 623.71	8 503.51	0.00%	0.00%
2015	7 749.89	8 690.73	7 750.13	8 690.97	0.00%	0.00%
2016	8 029.65	8 888.99	8 029.87	8 889.20	0.00%	0.00%
2017	8 263.20	9 174.20	8 264.89	9 175.89	0.02%	0.02%
2018	8 322.93	9 201.27	8 323.96	9 202.30	0.01%	0.01%



YEAR	SUBMISSION 2024		SUBMISSION 2025		2023/2024	
	CO <sub>2</sub> emissions	Total emissions	CO <sub>2</sub> emissions	Total emissions	CO <sub>2</sub> emissions	Total emissions
	Gg CO <sub>2</sub> eq.				%	
2019	7 482.55	8 350.38	7 490.44	8 358.27	0.11%	0.09%
2020	6 978.23	7 796.55	6 988.93	7 807.24	0.15%	0.14%
2021	8 366.44	9 206.03	8 386.76	9 226.34	0.24%	0.22%
2022	6 908.01	7 536.24	6 915.91	7 544.14	0.11%	0.10%

## 4.6. Sector-specific Improvements and Implementation of Recommendations

2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories was fully implemented during the previous inventory preparation. No other improvements have been done in this submission; all issues raised by UNFCCC review were implemented in the previous 2023 submission.

The study of CO<sub>2</sub> captured during the use of lime for sugar production is ongoing and it is planned to incorporate it in the future submissions.

## 4.7. Mineral Products (CRT 2.A)

### 4.7.1. Source-category Description

The major share of CO<sub>2</sub> emissions comes from the production and transformation of mineral products. Total emissions were 1 991.90 Gg of CO<sub>2</sub> in 2023 (only CO<sub>2</sub> emissions are reported in this category), the decrease approximately 15% when compared to previous year 2022. Compared to 1990, the decrease in mineral production is approximately 27%. Major trend behind the decrease in mineral production is decrease in demand of products.


The major share of emissions in this category belongs to cement production (65.0%), lime production (23.8%) and dead burned magnesia production (8.4%). The ceramics production shared 0.3% and glass production 0.6%. The rest of emissions (1.9%) are reported in other category. Emissions in 2.A.4.b are not occurring.

**Table 4.6:** CO<sub>2</sub> emissions in the category 2.A by subcategories in particular years

YEAR	2.A.1 Cement Production	2.A.2 Lime Production	2.A.3 Glass Production	2.A.4.a Ceramics	2.A.4.c Magnesia Production	2.A.4.d Other
	Gg					
1990	1 464.50	794.92	7.88	14.79	431.94	NO
1995	1 154.63	593.23	18.01	11.04	294.03	NO
2000	1 190.45	556.73	22.82	10.36	409.82	39.92
2005	1 256.40	711.96	33.04	13.06	476.01	42.49
2010	859.92	651.88	13.15	12.75	376.35	27.13
2011	1 261.79	672.41	11.83	11.65	363.83	37.83
2012	1 095.93	632.00	11.46	12.93	318.04	46.65
2013	1 135.27	560.14	13.22	14.94	279.56	27.10
2014	1 266.76	570.80	12.26	12.99	278.33	39.94
2015	1 308.57	534.30	11.93	14.24	247.76	34.56
2016	1 340.95	521.62	14.83	17.65	220.19	68.21
2017	1 367.05	507.78	15.20	20.82	291.28	75.00
2018	1 346.68	522.65	16.02	21.29	304.39	68.51
2019	1 404.27	489.24	18.16	21.52	295.15	56.62
2020	1 443.15	430.65	18.39	16.45	263.63	46.47

YEAR	2.A.1 Cement Production	2.A.2 Lime Production	2.A.3 Glass Production	2.A.4.a Ceramics	2.A.4.c Magnesia Production	2.A.4.d Other
	Gg					
2021	1 452.93	539.96	19.70	16.07	257.70	49.10
2022	1 489.72	532.42	16.33	14.47	229.43	50.33
2023	1 294.23	475.06	12.52	4.99	167.62	37.49

Figure 4.4: The share of CO<sub>2</sub> emissions on individual categories in the 2.A in 2023



#### 4.7.2. Cement Production (CRT 2.A.1)

Cement production plants in the Slovak Republic (four plants), where cement clinker is produced, are included into the EU ETS. Therefore, input data are directly taken from the EU ETS reports and from the reports of verifiers. Presented parameters are weighted averages. Total CO<sub>2</sub> emissions from cement clinker production were 1 294.23 Gg in 2023 and lower by 13% than in previous year. In comparison with the base year 1990, the CO<sub>2</sub> emissions in this category decreased by 12%. The reason of the lowering trend is the decreasing need for construction purposes.

Table 4.7: Activity data and CO<sub>2</sub> emissions in the category 2.A.1 in particular years

YEAR	Cement Clink Production	CaO Content	MgO Content	Correction Factor	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	kt				Gg	t/t
1990	2 835.75	64.60%*	NE	1.0184	1 464.50	0.5164
1995	2 235.75	64.60%*	NE	1.0184	1 154.63	0.5164
2000	2 313.71	64.36%*	NE	1.0184	1 190.45	0.5145
2005	2 352.68	64.31%	1.79%	1.0184	1 256.40	0.5340
2010	1 653.59	66.07%	2.60%	0.9506	859.92	0.5200
2011	2 433.86	67.13%	1.50%	0.9541	1 261.79	0.5184
2012	2 126.12	65.25%	1.86%	0.9680	1 095.93	0.5155
2013	2 161.32	65.53%	2.52%	0.9693	1 135.27	0.5253
2014	2 415.34	66.00%	2.23%	0.9668	1 266.76	0.5245
2015	2 506.12	65.70%	2.58%	0.9600	1 308.57	0.5221
2016	2 599.39	64.84%	2.36%	0.9647	1 340.95	0.5159
2017	2 698.82	64.83%	2.50%	0.9447	1 367.05	0.5065
2018	2 695.74	64.84%	2.39%	0.9336	1 346.68	0.4996
2019	2 854.64	65.11%	2.33%	0.9168	1 404.27	0.4919
2020	2 944.94	65.31%	2.28%	0.9116	1 443.15	0.4900

YEAR	Cement Clink Production	CaO Content	MgO Content	Correction Factor	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	kt				Gg	t/t
2021	2 937.74	65.45%	2.52%	0.9137	1 452.93	0.4946
2022	3 041.27	65.33%	2.56%	0.9058	1 489.72	0.4898
2023	2 682.07	65.39%	2.43%	0.8939	1 294.23	0.4825

\* Aggregated CaO content = CaO Content + 1.092/0.785×MgO content

### Methodological Issues

Cement is produced by a high temperature reaction of calcium oxide (CaO) with silica (SiO<sub>2</sub>) and with alumina (Al<sub>2</sub>O<sub>3</sub>). A source of calcium oxide is limestone (CaCO<sub>3</sub>). As the cement clink is produced at the temperature of 1 450°C the reaction produces carbon dioxide. The other emissions originate from impurities in the raw material (SO<sub>2</sub>). Based on the information provided by the EU ETS verifiers, tier 2 method according to the IPCC 2006 GL has been applied since 2002 based on plant specific information. The calculations provided by the producers in the EU ETS reports balanced CO<sub>2</sub> emissions based on cement clinker production and CaO and MgO contents. The data required for calculation of CO<sub>2</sub> emissions are summarized in **Table 4.8** (C = confidential, but available for the sectoral experts).

**Table 4.8:** Input data used for the CO<sub>2</sub> emissions estimation in the category 2.A.1 in 2023

PLANT/OPERATOR	CEMENT CLINK	CaO CONTENT	MgO CONTENT	CKD	COMPOSITION FACTOR	CO <sub>2</sub>
	kt	%	%			Gg
Cemmac	C	65.90%	1.68%	1.0277	0.9643	182.58
VSH (Danucem)	C	64.05%	4.48%	1.0090	0.6476	195.96
Danucem – Portland	C	65.32%	2.28%	1.0170	0.9009	535.18
Danucem – white	C	68.59%	2.10%	1.0135	1.0000	60.91
Považská cementáreň	C	65.87%	1.35%	1.0000	1.0000	319.60
<b>TOTAL</b>	<b>2 682.07</b>	<b>65.39%</b>	<b>2.43%</b>	<b>1.0129</b>	<b>0.8825</b>	<b>1 294.23</b>

Based on availability of information, the plant specific emission factors were used since 2002. The annual estimation of overall EFs is expressed as weighted average and is based on the specific contents of CaO and MgO in cement clinker in each producer and varies over the years. The implied CO<sub>2</sub> emission factor was 0.4825 t CO<sub>2</sub>/t of cement clink in 2023 (correction factor is also included in this value). Correction factor consists of CKD (Cement Kiln Dust) and so-called composition factor that represents the amounts of non-carbonate origin of CaO and MgO (using of ground granulated blast-furnace slag). All these data are plant specific.

$$\text{Corr. factor} = \text{CKD} * \text{Composition Factor}$$

Composition Factor

$$= \frac{(0.785 * \%CaO_c + 1.092 * \%MgO_c) * m_c - (0.785 * \%CaO_s + 1.092 * \%MgO_s) * m_s}{(0.785 * \%CaO_c + 1.092 * \%MgO_c) * m_c}$$

where: %CaO<sub>c</sub> is the fraction of CaO in cement clinker produced; %MgO<sub>c</sub> is the fraction of MgO in cement clinker produced; m<sub>c</sub> is the mass of cement clinker produced; %CaO<sub>s</sub> is the fraction of CaO in slag entering; %MgO<sub>s</sub> is the fraction of MgO in slag entering; m<sub>s</sub> is the mass of slag entering. However, the factor is directly known from the EU ETS reports of the plants.

### Uncertainties and Time-series Consistency

According to the ERT recommendation I.1 of ARR 2022, in the period 1990 – 1999 the average aggregated CaO content in the cement clinker was assumed to be very close to the default IPCC value from the IPCC 1996 GL (64.6%). The using of this aggregated CaO content is based on the average value of the CaO content in 2000 – 2003 (64.36% in 2000; 63.90% in 2001; 64.50% in 2022 and 65.70% in 2003). The weighted average value is 64.62%, which is very close to that IPCC value. Therefore, the

value (64.6%) was also assumed as country specific. In 2003, one plant with the lowest CaO content was closed for reconstruction. It was reopened in 2004 and the cement clinker with higher content of CaO is produced there since that time. This is the reason of higher aggregated CaO content and IEF since 2002 and therefore the years since 2004 were not considered for calculation of the average value. Another plant was renovated and did not produce cement clink in 2010. It resulted in decrease of emissions in 2010 and thereafter-significant increase in 2011 after its reopening.

In the period 1990 – 2004, the contents of MgO are not explicitly known. It was included in the CaO content based on stoichiometry; therefore we call it as aggregated CaO content.

Ground granulated blast-furnace slag has been also used as raw material since 2009, which results in additional increase of CaO and MgO contents (non-carbonate origin) in the cement clinker. Therefore, we use the correction factor instead of CKD factor in the calculation. Correction factor is CKD multiplied by the so called “Composition Factor”. CKD and Composition factors are plant specific, and they are known since 2008. Because of conservative approach, the highest value of the CKD (1.0184; the value close to the default CKD) was used for time series before 2008. For this time series, Composition factor was assumed to be 1, no correction for slag was made.

There were totally five cement sites in 1990 – 1996 in Slovakia. In 1997, one of them finished production of cement clinker. In 2003 and 2010, one of the other four cement sites did not produce cement clinker. During the period 1990 – 2022, no changes in technologies were made in plants; only the changes in composition of the clinker and use of raw materials (slag) occurred.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.7.3. Lime Production (CRT 2.A.2)

From a chemical point of view, lime is calcium oxide (CaO). It is produced by a thermal decomposition of limestone at the temperatures of 1 040 – 1 300°C. Carbon dioxide is produced according to the same reaction scheme as shown above in the case of cement production. Only CO<sub>2</sub> emissions are reported in this category. Total CO<sub>2</sub> emissions from lime production decreased by 11% when compared with the previous year and were 475.06 Gg in 2023. The decrease in emissions by 40% is achieved when compared with the base year.

**Table 4.9: Activity data and CO<sub>2</sub> emissions in the category 2.A.2 in particular years**

YEAR	Lime Production	CO <sub>2</sub> Emissions	CaO Content		
	kt	Gg			
1990	1 076.00	794.92	91.20%		
1995	803.00	593.23	91.20%		
2000	753.59	556.73	91.20%		

YEAR	Lime Production	CO <sub>2</sub> Emissions	CaO Content	MgO Content	“HYPOTHETIC” CaO Content
	kt	Gg			
2001	815.96	602.80	90.56%	0.47%	91.20%
2005	913.08	711.96	89.55%	4.72%	96.12%
2010	822.36	651.88	86.95%	7.72%	97.70%
2011	856.05	672.41	85.94%	7.82%	96.82%

YEAR	Lime Production	CO <sub>2</sub> Emissions	CaO Content	MgO Content	“HYPOTHETIC” CaO Content
	kt	Gg			
2012	797.33	632.00	78.32%	13.96%	97.74%
2013	716.54	560.14	87.39%	6.40%	96.30%
2014	727.63	570.80	86.81%	7.26%	-
2015	680.20	534.30	87.34%	6.93%	-
2016	663.02	521.62	86.17%	7.49%	-
2017	640.06	507.78	87.47%	7.46%	-
2018	668.99	522.65	86.87%	6.95%	-
2019	634.58	489.24	87.28%	6.21%	-
2020	554.22	430.65	87.78%	6.49%	-
2021	696.12	539.96	74.33%	15.66%	-
2022	688.95	532.42	87.66%	6.56%	-
2023	630.68	475.06	86.84%	6.29%	-

“Hypothetic” CaO content = CaO Content + 1.092/0.785xMgO content

### Methodological Issues

**Table 4.9** shows “hypothetic” CaO content and includes stoichiometric data on the CaO and MgO contents. This approach was used due to no availability of distinguished data for the period 1900 – 2000. In that period, the same content of CaO in the lime is assumed (91.2%). This value is based on the data available for 2001 and 2002 and on all the data available in the period 1990 – 2000. The average content of CaO in lime was (91.2% ± 0.2%) in the period 1990 – 2002. “Hypothetic CaO content” is not presented in **Table 4.9** since 2014 because it is not necessary to report it for recent years. Tier 2 according to the IPCC 2006 GL was used for the whole time series with the combination of plant specific activity data and emission factors estimated for each plant. Calculation is based on data provided by the producers of lime in questionnaires and in the EU ETS reports (produced lime and CaO and MgO contents). Data required and used for CO<sub>2</sub> emissions estimation are summarized in **Table 4.10**.

Based on availability of information, the plant specific emission factors have been used since 2001. The annual estimation of national EFs varies over the years. Calculation of EFs is based on weighted average based on purity of lime in individual production unit. The implied CO<sub>2</sub> emission factor is 0.753 t CO<sub>2</sub>/t of lime in 2023 (correction factor is included in the IEF). Correction factor presented in **Table 4.10** represents LKD (Lime Kiln Dust) as introduced in the IPCC 2006 GL. Data necessary for determination of correction factor were provided by the plant operators. When LKD was not provided by operator, default value (1.02) was used. Total quantity of produced lime in Slovakia was 630.68 kt in 2023. Activity data used for inventory are summarized in **Table 4.10** Large and medium producers provided activity data in their EU ETS reports or reports from verifiers, small plants like sugar producers provided activity data based on questionnaires to the SNE.

**Table 4.10: Activity data necessary for the estimation of CO<sub>2</sub> emissions in the category 2.A.2 in 2023**

Plant	Lime Production	CaO Content	MgO Content	LKD	CO <sub>2</sub> Emissions
	kt				Gg
Calmit	C	92.00%	1.19%	1.0000	81.82
Dolvap Varín	C	89.19%	5.87%	1.0000	71.95
Carmeuse	C	84.45%	8.11%	1.0047	300.10
Others*	C	92.50%	2.00%	1.0200	21.19
TOTAL	630.68	86.84%	6.29%	1.0038	475.06

C = confidential, \*aggregated data from small plants not covered by the EU ETS as sugar producers

---

## Uncertainties and Time-series Consistency

Time series consistency is assured by using the “hypothetic” CaO content during the period 1990 – 2000 as explained in detail above. This content is compared with the data presented in 2001 and 2002. Dolomitic lime production started in one plant in 2003 and the CaO content is not comparable since this year. Because of the dolomitic lime production, the overall IEF has increased since that time, as well. Lime produced by sugar producers is included in inventory as “others”. The country specific LKD factor estimated in 2013 was used for the rest of the time series before 2013 because no other data on LKD were available. In 2014 and 2015, the country specific LKD factor was very close to the factor reported in 2015; therefore, no recalculation of the historical data was necessary.

In Slovakia, lime is produced by three lime producers that are included in the EU ETS system and four other producers (sugar plants, pulp and paper and the other plant – production of secondary aluminium) that are not included in the EU ETS. It can be assumed that CO<sub>2</sub>, which is evolved during the lime production in sugar plants, is back captured there. However, because of no detailed data about back capturing of CO<sub>2</sub> in the lime and due to the ensuring of conservatism, no capturing of CO<sub>2</sub> is reported in the inventory. The CO<sub>2</sub> emissions from lime production by the pulp and paper industry are not estimated because of the use of the Kraft chemical recovery process, which results in biogenic CO<sub>2</sub> emissions originating from biomass input.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.7.4. Glass Production (CRT 2.A.3)

Basic raw material for glass production is silica (SiO<sub>2</sub>). Limestone (CaCO<sub>3</sub>), dolomite (CaMg(CO<sub>3</sub>)<sub>2</sub>), soda ash (Na<sub>2</sub>CO<sub>3</sub>), potash (K<sub>2</sub>CO<sub>3</sub>), Pb<sub>3</sub>O<sub>4</sub>, Al<sub>2</sub>O<sub>3</sub>, and colouring agents are used in glass production process. NMVOC and CO are the most important emissions, but they are not reported in this category (notation key “IE” was used). These emissions are allocated in 1.A.2.f. Only CO<sub>2</sub> emissions were estimated in this category and were 12.52 kt in 2023.

#### Methodological Issues

CO<sub>2</sub> emissions from used carbonates were calculated by tier 3 method on the stoichiometry principle according to the IPCC 2006 GL. The calcination fraction was assumed one. Based on availability of information, the plant specific emission factors were used since 2004. The annual estimation of national EFs varies over the years. According to the [ERT recommendation I.3 of the ARR 2022](#), calculation of EFs is based on weighted average based of used carbonates and CO<sub>2</sub> emissions in individual production unit. Implied emission factor was 0.421 t/t of used carbonates mixture or 0.038 t/t of glass produced in 2023. This value is much lower than the default factor (using tier 1) used in the IPCC 2006 GL. It is caused by using alternative additions to raw materials as calumite (blast furnace granulated slag), colemanite (CaB<sub>3</sub>O<sub>4</sub>(OH)<sub>3</sub>·H<sub>2</sub>O) or clay as well as by using different amounts of recycled glass. However, it should be mentioned that due to the using of higher tier (tier 3), the amount of recycled glass is not necessary to follow. The main reason of such low IEF is the production of glass fibers at which the above mentioned non-carbonate raw materials are used (IEF = 0.015 t / t of glass fiber). For the inventory, the production of glass fibers is included in white glass production with the share ca 30%. The other glass producers report the IEF (0.09 – 0.11) t / t of glass, which is in accordance with tier 1 IEF.

Glass production based on direct information from producers was as follows: 325.54 kt of white glass in 2023. No leaded glass or green glass was produced in 2023. SrCO<sub>3</sub> and Li<sub>2</sub>CO<sub>3</sub> were not used for glass production. Total amounts of used carbonates were 12.52 kt in 2023 and time series is shown in [Table 4.11](#).

**Table 4.11: Total amounts of used carbonates and CO<sub>2</sub> emissions in particular years**

YEAR	CaCO <sub>3</sub>	K <sub>2</sub> CO <sub>3</sub>	Na <sub>2</sub> CO <sub>3</sub>	BaCO <sub>3</sub>	MgCO <sub>3</sub>	SrCO <sub>3</sub>	Li <sub>2</sub> CO <sub>3</sub>	Total	CO <sub>2</sub>
	kt								Gg
1990	17.91	a)	a)	a)	a)	a)	a)	17.91	7.880
1995	40.93	a)	a)	a)	a)	a)	a)	40.93	18.007
2000	51.87	a)	a)	a)	a)	a)	a)	51.87	22.821
2005	55.45	2.75	16.00	0.89	1.76	0.01	0.01	76.87	33.038
2010	15.89	0.48	13.62	1.52	0.01	NO	NO	31.52	13.145
2011	15.17	0.31	11.49	0.01	0.54	NO	NO	27.52	11.825
2012	14.75	0.03	11.45	0.01	0.39	NO	NO	26.63	11.456
2013	15.31	0.72	14.24	0.56	0.43	NO	NO	31.26	13.224
2014	14.22	0.64	13.29	0.48	0.34	NO	NO	28.97	12.262
2015	14.83	0.46	11.92	0.46	0.44	NO	NO	28.11	11.931
2016	17.64	0.57	15.55	0.70	0.53	NO	NO	34.99	14.828
2017	17.74	0.66	16.03	0.74	0.69	NO	NO	35.86	15.195
2018	17.70	0.76	17.99	0.78	0.67	NO	NO	37.90	16.020
2019	19.94	0.71	20.91	0.86	0.55	NO	NO	42.98	18.160
2020	20.19	0.65	21.27	0.74	0.59	NO	NO	43.44	18.389
2021	22.14	0.76	21.98	0.90	0.75	NO	NO	46.53	19.697
2022	19.11	0.72	17.93	0.88	0.12	NO	NO	38.76	16.335
2023	14.38	0.56	14.01	0.69	0.09	NO	NO	29.73	12.521

a) Carbonates are included in the form of calcium carbonate (based on stoichiometry).

### Uncertainties and Time-series Consistency

There were several operators producing glass (from 3 to 7) in Slovakia during the time series 1990 – 2023. Detailed statistics of used carbonates is available only since 2003 and therefore methodology used in GHG inventory is based on total carbonates (in the form of calcium carbonate) calculated based on stoichiometry. This calculation was provided by reverse method, it means, that the specific averages CO<sub>2</sub> EFs per 1 t of each type of glass was known for every producer (except for one plant, where the same EFs was used as for the similar type of glass production). Therefore, the CO<sub>2</sub> emissions are known and only one (“aggregated”) carbonate can be calculated from that data. The plant specific EFs are commercially confidential and they will be available during review process on request of the ERT. New production of lead glass started in 2000 and ended in 2002. Similarly, new production of lead glass started in 2003 and ended in 2006. Both productions were small. The increase in emissions since 2005 is caused by change of one big plant owner (resulting in increase of production and emissions). Other plants were closed in 2008 and 2012. Since 2008, colemanite and calumite slag are widely used in the biggest glass plant in order to replace carbonates, which resulted in significant emissions decrease. Since 2009, emissions from glass production have been almost stable.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.7.5. Other Process Uses of Carbonates – Ceramics (CRT 2.A.4.a)

Ceramics includes the production of bricks and roof tiles, vitrified clay pipes, refractory products, expanded clay products, wall and floor tiles, table and ornamental ware (household ceramics), sanitary ware, technical ceramics, and inorganic bonded abrasives. Process-related CO<sub>2</sub> emissions reported from ceramics result from the calcination of carbonates in the clay, as well as from addition of additives. CO<sub>2</sub> emissions from ceramics production were 4.99 Gg CO<sub>2</sub> in 2023 the decrease almost by 2/3 when compared with previous year 2022. The decrease is caused by closing of the one plant and significant decrease in production in the others.

##### Methodological Issues

CO<sub>2</sub> emissions from the used carbonates were calculated by tier 3 method according to the IPCC 2006 GL based on principle of the stoichiometry. The calcination fraction assumed to be one. Based on available information, plant specific emission factors were used since 1990. The annual estimation of country specific EF is expressed as weighted average and is based on the stoichiometry of carbonates and CO<sub>2</sub>. Implied emission factor calculated in 2023 was 0.48 t/t of used carbonates mixture. This approach was used for all years.

Fraction of carbonates in raw materials is determined analytically in each plant. Based on the analysis, amounts of used CaCO<sub>3</sub> and MgCO<sub>3</sub> are obtained. Total amounts of used carbonates were 10.42 kt in 2023 and time series is presented in [Table 4.12](#).

**Table 4.12:** Total used carbonates and CO<sub>2</sub> emissions the category 2.A.4.a in particular years

YEAR	CaCO <sub>3</sub>	MgCO <sub>3</sub>	Total Carbonates	CO <sub>2</sub> Emissions
	kt			Gg
1990	25.41	6.92	32.33	14.79
1995	17.19	6.66	23.85	11.04
2000	15.79	6.54	22.33	10.36
2005	21.80	6.64	28.44	13.06
2010	18.95	8.46	27.41	12.75
2011	16.61	8.32	24.93	11.65
2012	19.06	8.71	27.77	12.93
2013	22.76	9.43	32.19	14.94
2014	19.64	8.33	27.97	12.99
2015	21.83	8.88	30.71	14.24
2016	29.20	9.20	38.40	17.65
2017	34.82	10.53	45.35	20.82
2018	33.55	12.50	46.05	21.29
2019	35.65	11.18	46.83	21.52
2020	23.62	11.59	35.22	16.45
2021	17.30	16.20	33.50	16.07
2022	11.83	17.75	29.58	14.47
2023	5.52	4.90	10.42	4.99

##### Uncertainties and Time-series Consistency

The same tier approach is used for period 1990 – 2023. The presented data are obtained directly from producers. The missing data for some ceramics producers was interpolated or extrapolated for the periods 1990 – 1991 and 1993 – 1995 on the level of individual producer with the consideration of economic aspects of construction industry in Slovakia (it served as limiting conditions of interpolation or extrapolation methods) as it was described in previous submissions (SVK NID 2014). Several (14) plants were reported in this category during time series, recently only five of them report CO<sub>2</sub> emissions. The others were closed. New owner came into the market and bought three existing plants in 2007. The high increase in CO<sub>2</sub> emissions is caused by significant increase in production in those plants. However, in



---

2009, one plant was closed due to the economic reasons and decrease in production occurred in the other plants. In 2023, another plant was closed due to the economic reasons and the others significantly decreased the production.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### **4.7.6. Other Process Uses of Carbonates – Other Uses of Soda Ash (CRT 2.A.4.b)**

Soda ash is used in a variety of applications, including, glass production, soaps and detergents, flue gas desulphurization, chemicals, pulp and paper and other common consumer products. Using of soda ash (where is applicable in Slovakia) is reported in the category where it is consumed (see category 2.A.3 Glass Production). In Slovakia, soda ash is used in glass industry, only. No plants using soda ash for the other possible applications are present in Slovakia except of flue gas desulphurization. For flue gas desulphurization only calcium carbonate is used in Slovakia.

#### **4.7.7. Other Process Uses of Carbonates – Non-Metallurgical Magnesia Production (CRT 2.A.4.c)**

Magnesite clinker for refractory materials is produced in Slovakia. According to the IPCC 2006 GL production of dead burned magnesia for refractory materials is reported in this category. Carbon dioxide is produced by thermal decomposition of magnesite. This chemical reaction scheme of the thermal decomposition is  $\text{MgCO}_3 = \text{MgO} + \text{CO}_2$ . Total  $\text{CO}_2$  emissions from magnesite production were 167.62 Gg in 2023 and decreased by ca 27% when compared with the year 2022. The decrease is due to the decrease in the production of magnesite clinker. One of the plant was operating only 55% of time that was operating in 2022. The reduction in production was due to the energy crisis, as well as the war in Ukraine and the associated reduction in sales. When compared to 1990, the decrease is approximately 61%. It was caused by closing of one plant during the monitored time series and occasional closing of magnesite clinker production in others (in this case, the plant produced the refractory materials from stocked or bought magnesite clinker).

##### **Methodological Issues**

Magnesite raw materials used in the Slovak Republic contain small amounts of  $\text{CaCO}_3$  and  $\text{FeCO}_3$ . Emissions are calculated on the stoichiometric base ( $\text{CO}_2$  and respective carbonate). The amounts of magnesite raw materials and emissions of  $\text{CO}_2$  in the period of 1990 – 2023 are summarized in [Table 4.13](#).  $\text{CH}_4$  and  $\text{N}_2\text{O}$  emissions are not occurring and therefore notation key “NO” was used for time series.

$\text{CO}_2$  emission factors used for emissions estimation in this category are as follows: 0.44 t/t  $\text{CaCO}_3$ , 0.522 t/t  $\text{MgCO}_3$  and 0.38 t/t  $\text{FeCO}_3$ . Total consumption of magnesite raw materials in the Slovak Republic was 350.81 kt in 2023. The composition of raw materials is summarized in [Table 4.11](#). It should be noted that  $\text{CaCO}_3$  and  $\text{FeCO}_3$  contents are included in  $\text{MgCO}_3$  content on the basis of stoichiometry for the years before 1999, due to lack of input data.

**Table 4.13: Consumption and composition of magnesite raw materials and CO<sub>2</sub> emissions in the category 2.A.4.c in particular years**

YEAR	Raw Materials Used	MgCO <sub>3</sub> Content	CaCO <sub>3</sub> Content	FeCO <sub>3</sub> Content	CO <sub>2</sub> Emissions	EF
	kt				Gg	t/t
1990	887.74	0.9321	*	*	431.94	0.487
1995	604.32	0.9321	*	*	294.03	0.487
2000	850.57	0.8850	0.0324	0.0147	409.82	0.482
2005	988.58	0.8804	0.0382	0.0135	476.01	0.482
2010	820.32	0.8424	0.0400	0.0038	376.35	0.459
2011	724.27	0.9193	0.0444	0.0077	363.83	0.502
2012	634.97	0.9090	0.0436	0.0189	318.04	0.501
2013	603.38	0.8418	0.0489	0.0063	279.56	0.463
2014	590.33	0.8210	0.0452	0.0606	278.33	0.471
2015	550.04	0.8063	0.0299	0.0432	247.76	0.450
2016	462.81	0.8462	0.0383	0.0453	220.19	0.476
2017	622.44	0.8260	0.0475	0.0418	291.28	0.468
2018	657.28	0.8168	0.0477	0.0415	304.39	0.463
2019	634.89	0.8178	0.0498	0.0423	295.15	0.465
2020	560.73	0.8261	0.0533	0.0407	263.63	0.470
2021	549.84	0.8180	0.0560	0.0448	257.70	0.469
2022	476.47	0.8294	0.0666	0.0508	229.43	0.482
2023	350.81	0.8298	0.0639	0.0435	167.62	0.478

\*carbonates reported in MgCO<sub>3</sub> on the basis of stoichiometry

### Uncertainties and Time-series Consistency

There were six plants producing magnesite clinker in Slovakia in 1990 – 2023. One of them ended its production in 1991. New plant entered into market in 2004; in 2007, it finished its production. Another new plant entered into market also in 2004; in 2009, it finished its production. This second operator has had very limited production of clinker. Another one stopped its production of magnesite clinker for years 1992 – 1994. Two plants continuously produced magnesite clinker since 1990. These two plants have one owner.

The same tier approach is used for the whole period 1990 – 2020. Because of the lack of input data on the consumption of magnesite raw materials and their composition before 2008, the data on the production of magnesite clinker and its composition were used to reconstruct the time series before 2008. More details on this procedure were described in the [Annex 4.1](#) of the SVK NID 2016. However, only activity data on raw materials for the time series 1990 – 2007 were reconstructed (approximated data), while the CO<sub>2</sub> emissions are exactly calculated from the magnesite clinker production and its composition. Therefore, the comparison of the IEF changes is not possible between years.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.7.8. Other Process Uses of Carbonates – Other (CRT 2.A.4.d)

Carbon dioxide is produced at thermal and chemical decomposition of limestone or other carbonates. The maximum values of the CO<sub>2</sub> emission factors based on the stoichiometry are 440 kg CO<sub>2</sub> per ton of consumed CaCO<sub>3</sub> and 522 kg CO<sub>2</sub> per ton of consumed MgCO<sub>3</sub>, which are also recommended by the IPCC 2006 GL. The CO<sub>2</sub> emissions estimated in this category are based on limestone consumed in desulphurization process of coal.

##### Methodological Issues

Limestone used in Slovakia often contains a small amount of MgCO<sub>3</sub>. CO<sub>2</sub> emissions are estimated using the balance of carbonates according to the IPCC 2006 GL and the plant specific emission factors. The amount of consumed carbonates according to the different sources and CO<sub>2</sub> emissions in the period 1990 – 2023 are summarized in [Table 4.14](#).

Based on availability, the plant specific emission factors have been used since 2004. The annual estimation of EFs is expressed as weighted average and is based on the stoichiometry of limestone and dolomite in the mixtures in each producer. Therefore, the IEF varies over the years. Implied emission factor in 2023 was 0.441 t/t of used carbonates mixture.

Total amount of carbonates used at desulphurization was 84.92 kt in 2023, the activity data are summarized in [Table 4.14](#). The consumption increased significantly in 2016, the consumption of limestone reached the highest level since start of using of the desulphurization technology. The probable reason of the increased using of limestone is the new emission limits for SO<sub>2</sub> since January 1<sup>st</sup>, 2016. Consumption of limestone increased approximately two times in the power plant using brown coal. This trend continued also in 2018. In 2019, this trend was interrupted, the consumption of brown coal decreased. Total CO<sub>2</sub> emissions estimated in this category were 37.49 Gg in 2023. Such decrease in emissions and carbonates consumption in comparison with previous year (25%) is caused by closing a big coal power plant.

**Table 4.14: Total used carbonates and CO<sub>2</sub> emissions in the category 2.A.4.d in particular years**

YEAR	Desulphurization (CaCO <sub>3</sub> )	Desulphurization (MgCO <sub>3</sub> )	Total Carbonates	CO <sub>2</sub> Emissions
	kt			
1990	NO	NO	NO	NO
1995	NO	NO	NO	NO
2000	88.86	1.58	90.44	39.92
2005	94.52	1.73	96.25	42.49
2010	60.49	0.99	61.48	27.13
2011	84.46	1.28	85.74	37.83
2012	103.83	1.84	105.67	46.65
2013	59.84	1.48	61.32	27.10
2014	88.39	2.01	90.40	39.94
2015	76.95	1.35	78.30	34.56
2016	150.09	4.16	154.25	68.21
2017	166.50	3.34	169.84	75.00
2018	150.99	3.97	154.96	68.51
2019	125.39	2.78	128.17	56.62
2020	103.23	2.01	105.24	46.47
2021	109.13	2.09	111.21	49.10
2022	111.60	2.35	113.95	50.33
2023	83.45	1.48	84.92	37.49

## Uncertainties and Time-series Consistency

The same tier approach is used for period 1996 – 2023. Before 1996, no desulphurization technology was used in Slovakia. Data presented in [Table 4.14](#) were obtained directly from producers. The decrease in consumption of limestone for desulphurization in 2010 was caused by using of 15 654 t stock lime bought from lime producer for desulphurization. It represents (using back calculation to carbonates) approximately 25.55 kt of CaCO<sub>3</sub> and 0.17 kt of MgCO<sub>3</sub>. Emissions from that lime consumption were already allocated and reported in 2.A.2 in 2010. In 2012, no using of stock lime was reported and therefore emissions are higher than in previous years. In 2013 emissions decreased again (by 42%) due non-use of the desulphurization process in one plant. In 2014, the desulphurization process was again used in that plant. Since 1990, there have been seven plants with desulphurization technology. The significant increase in limestone consumption in 2016 is a result of the new emission limits for SO<sub>2</sub> since January 1<sup>st</sup>, 2016. Consumption of limestone increased approximately two times in the power plant using brown coal. The recent decrease is caused by closing coal power plant.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

## 4.8. Chemical Industry (CRT 2.B)


Production of ammonia is the major source of CO<sub>2</sub> emissions and nitric acid production is the major source of N<sub>2</sub>O emissions in this category. Total GHG emissions reported in this category were 965.60 Gg of CO<sub>2</sub> eq. in 2023. The decrease of emissions in the comparison with the previous year is approximately 10% and decrease by 47% in the comparison with the base year. The decrease is caused by significantly lower production of ammonia. The significant decrease in emissions was reported in nitric acid production where using of secondary YARA catalyst fully reflected this situation since 2011. In 2013, also the last producer of nitric acid started using of this secondary catalyst. Decrease of emissions in calcium carbide production is caused by decrease in production and change of raw material. Within category, major share (53.9%) in emissions belongs to ammonia production, 39.0% belongs to petrochemicals production, and 4.8% belongs to nitric acid production and 3.3% to carbide production. The hydrogen production (other) was reallocated into Energy sector according to the IPCC 2019 Refinement.

**Table 4.15:** Emissions in the category 2.B according to the subcategories in particular years

YEAR	2.B.1 Ammonia Production	2.B.2 Nitric Acid Production	2.B.5 Carbide Production	2.B.8 Petrochem. Production	2.B.10 Other
	<i>Gg of CO<sub>2</sub> eq.</i>				
1990	332.36	1 072.65	0.00	428.80	NO
1995	488.47	1 050.07	139.01	459.91	NO
2000	521.73	904.61	156.73	462.68	NO
2005	573.24	1 098.05	176.72	371.40	NO
2010	388.06	772.56	197.56	403.75	NO
2011	578.73	359.93	222.28	425.75	NO
2012	546.68	258.20	141.26	319.26	NO
2013	675.35	115.08	95.35	330.79	NO
2014	530.30	128.66	85.76	250.60	NO
2015	639.45	124.30	48.47	331.82	NO

YEAR	2.B.1 Ammonia Production	2.B.2 Nitric Acid Production	2.B.5 Carbide Production	2.B.8 Petrochem. Production	2.B.10 Other
	Gg of CO <sub>2</sub> eq.				
2016	564.58	107.89	63.16	338.22	NO
2017	633.80	93.35	59.35	357.84	NO
2018	791.47	93.85	68.26	399.41	NO
2019	689.15	80.58	62.05	344.14	NO
2020	703.96	67.54	45.83	381.66	NO
2021	769.53	56.64	47.57	395.49	NO
2022	638.54	52.76	35.49	349.63	NO
2023	510.59	46.73	31.82	376.47	NO

Figure 4.5: The share in CO<sub>2</sub> emissions of individual subcategories in 2.B in 2023



#### 4.8.1. Ammonia Production (CRT 2.B.1)

Ammonia is made from nitrogen and hydrogen by fine-tuned versions of the process developed by Haber and Bosch  $N_2 + 3H_2 = 2NH_3$ . In principle, the reaction between hydrogen and nitrogen is easy. However, to get a respectable yield of ammonia in a chemical plant a catalyst and extreme pressures up to 600 atmospheres and temperature of 400°C are needed. The results of ammonia production in Slovakia are summarized in [Table 4.16](#).

##### Methodological Issues

Tier 3 method according to the IPCC 2006 GL was applied in the emissions estimation of the category 2.B.1 and the plant specific emission factors were used for whole time series. The information on ammonia production and natural gas consumption for its production was provided directly by the operators. The measured values of natural gas consumption provided by the operator were used for CO<sub>2</sub> emissions estimation and calculated according to the relationship:

$$E(CO_2) = FR \cdot CF \cdot CCF \cdot OF \cdot \frac{44}{12} - R(CO_2)$$

where: FR is total consumption of natural gas for ammonia production in Nm<sup>3</sup>; CF is conversion factor in MJ/m<sup>3</sup> (35.471 in 2023); CCF is content of carbon in the fuel in t/TJ (15.337 in 2023) and OF is oxidation factor of the fuel (1). It should be noted, that parameters (NCV, EF) used for natural gas are plant specific. R(CO<sub>2</sub>) represents the amount of carbon dioxide that is recovered and used for urea production. In Slovakia, urea is produced and respective amounts of CO<sub>2</sub> are subtracted from the calculated emissions. Due the subtracting of CO<sub>2</sub> from urea production, the import/export of urea is yearly monitored. Emissions from the use of urea are reported in the Agriculture sector, category 3.H

Urea application and in 2.D.3 Other (using of urea in urea-based catalytic converters). The use of urea in catalytic converters for NO<sub>x</sub> emissions in cars is calculated by the COPERT 5 model ([Chapter 3](#)). The use of urea in industrial plants is annually monitored by questionnaires that are sent to the operators at which the decrease of NO<sub>x</sub> emissions occurred. QA/QC on the use of urea, its export/import comparison is described in the [Chapter 4.4](#) and [Annex 4.3](#).

The implied emission factor is 1.23 t CO<sub>2</sub> per 1 t of ammonia produced in 2023 after subtracting of CO<sub>2</sub> used for urea production. Without subtracting of CO<sub>2</sub> used for urea production the implied emission factor is 1.68 t CO<sub>2</sub> per 1 t of ammonia. The methane and N<sub>2</sub>O emission factors are IPCC default: 1 kg/TJ of natural gas (CH<sub>4</sub>) and 0.1 kg/TJ of natural gas (N<sub>2</sub>O). Results are provided in [Tables 4.16](#) and [4.17](#). Production of ammonia decreased by 11% in 2023 when compared with 2022 and it is a key category in level and trend assessment. The producer supplied the data on the total consumption of natural gas for the ammonia production in 2023 that are necessary for the calculation of emissions. The presented data are based on direct measurements in plant. In 2019, new, very modern, ammonia technological line started, which resulted in lower CO<sub>2</sub> emission.

**Table 4.16: Ammonia production and GHG emissions in particular years**

YEAR	Ammonia Production	CO <sub>2</sub> Emissions*	CH <sub>4</sub> Emissions	N <sub>2</sub> O Emissions	NG Consumption	
	kt	Gg	t		mil. m <sup>3</sup>	TJ
1990	360.00	616.97	10.83	1.08	322.54	10 827.83
1995	383.80	654.14	11.70	1.17	343.87	11 698.41
2000	403.00	683.85	12.36	1.24	361.07	12 359.46
2005	426.35	721.40	13.06	1.31	381.99	13 064.02
2010	233.56	484.65	8.75	0.88	254.31	8 753.49
2011	455.48	779.42	14.07	1.41	407.74	14 070.98
2012	377.30	717.42	12.92	1.29	373.90	12 922.60
2013	474.91	888.08	15.98	1.60	461.25	15 979.72
2014	346.27	660.68	11.86	1.19	340.71	11 856.72
2015	476.94	884.82	15.88	1.59	454.27	15 878.88
2016	403.96	787.01	14.10	1.41	401.92	14 103.50
2017	458.88	873.80	15.70	1.57	449.16	15 700.36
2018	516.74	1 028.79	18.47	1.85	529.40	18 474.44
2019	491.95	822.68	14.77	1.48	422.85	14 770.06
2020	545.23	883.52	15.86	1.59	452.87	15 856.94
2021	580.51	930.46	16.64	1.66	475.90	16 638.42
2022	462.12	750.41	13.37	1.34	379.70	13 366.19
2023	413.54	693.10	12.32	1.23	347.45	12 324.55

\* CO<sub>2</sub> emissions without consideration of urea production

**Table 4.17: Urea production, CO<sub>2</sub> used for the production and resulting CO<sub>2</sub> emissions in particular years**

YEAR	Urea Production	CO <sub>2</sub> Consumed	Net CO <sub>2</sub> Emissions*	IEF
	kt	Gg		t/t
1990	C	285.20	331.77	0.922
1995	C	166.31	487.83	1.271
2000	C	162.79	521.06	1.293
2005	C	148.87	572.52	1.343
2010	C	97.07	387.58	1.659
2011	C	201.46	577.96	1.269
2012	C	171.45	545.98	1.447
2013	C	213.60	674.48	1.420

YEAR	Urea Production	CO <sub>2</sub> Consumed	Net CO <sub>2</sub> Emissions*	IEF
	kt	Gg		t/t
2014	C	131.03	529.65	1.530
2015	C	246.24	638.58	1.339
2016	C	223.20	563.81	1.396
2017	C	240.86	632.94	1.379
2018	C	238.32	790.46	1.530
2019	C	134.34	688.35	1.400
2020	C	180.42	703.09	1.290
2021	C	161.83	768.62	1.324
2022	C	112.59	637.81	1.380
2023	C	183.19	509.92	1.233

\*CO<sub>2</sub> emissions with consideration of urea production, C = confidential (available in NS SR archive)

### Uncertainties and Time-series Consistency

Consistent tier 3 approach is used for the whole period 1990 – 2023. Higher emission factor in 2010 was caused by malfunctions in plant. The ammonia was not produced for 3.5 months in 2010. The emissions were higher as usual at the new start of the production. In 2011, the EF decreased to the values of the same level as before the malfunction. The reason of the increased production of ammonia is the new production line that was put in the operation during the year 2018. Since 2019, the new (modern) production line is fully operational. The investments in its construction amounted to 310 million €. Nowadays, the Agrofert Group in Šala has the most modern and the most ecological ammonia production technology not only in Slovakia, but also in Europe. It resulted in the decrease of the CO<sub>2</sub> emissions and IEF from the technological step (decrease by ca 15%).

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.8.2. Nitric Acid Production (CRT 2.B.2)

Globally, nitric acid production consumes about 20% of all produced ammonia. Nitric acid production in the Slovak Republic is an important source of N<sub>2</sub>O emissions and a key category in level and trend assessment. Total nitric acid production decreased by ca 9% in 2023 when compared to 2022. However, the N<sub>2</sub>O emissions decreased by 11% in 2023 when compared with 2022. Typical characteristic of the used technology (with secondary YARA catalyst) is that emissions are low but fluctuate in a certain degree. Thus, continuous monitoring of emissions is necessary.

#### Methodological Issues

Since 2005, N<sub>2</sub>O and NO<sub>x</sub> emissions are continuously monitored by the nitric acid producers with medium-pressure and high-pressure plant. Since 2013, the monitoring is running also in the third (the last) plant. Nitric acid is produced in three industrial plants situated in Slovakia owned by one provider since 2012. Nitric acid is produced by using two technologies: two medium-pressure plants and one high-pressure plant. The N<sub>2</sub>O emissions are directly measured during these processes. According to this, the emission factors were estimated annually, based on certified measurements in this plant.

- Atmospheric-pressure EFs: Production in atmospheric plant ended in 1999. The emission factor 4.5 kg N<sub>2</sub>O/1 t of HNO<sub>3</sub> was recommended for this type of technology in 2006 IPCC Guidelines.

In 2019 IPCC Refinement the emission factor was changed to 5.0 kg N<sub>2</sub>O/1 t of HNO<sub>3</sub>; which mean the increase by 11.1%. Because the technological line of the atmospheric plant was old we used for it the emissions factor 13.0 kg N<sub>2</sub>O/1 t of HNO<sub>3</sub> according the recommendation of the ERT review. We adopted the increase of emission factor by 11.1% also for this value. Thus, the recalculation of the historical data was done, that is presented in SVK NID 2024 ([Chapter 4.5, Table 4.4](#)).

- **Medium-pressure EFs:** Two medium pressure plants produce nitric acid in Slovakia. One of them directly measured N<sub>2</sub>O emissions in 2005 – 2010 (reg. No SNAS 230/S-189). Results are provided in [Table 4.18](#).

**Table 4.18: Measured EFs in medium pressure nitric acid plant in 2005 – 2010**

YEAR	2005	2006	2007	2008	2009	2010
	kg/t					
EF N <sub>2</sub> O	7.3	10.33	10.33	7.6	7.5	7.5

The malfunction in 2006 – 2007 resulted in higher N<sub>2</sub>O emission factors. The average value of the emission factor (7.5 kg/1 t of HNO<sub>3</sub>) calculated based on presented period (except 2006 and 2007 values) was used as EF in medium pressure plant for the period 1990 – 2004. The same value was also measured before the technological change, which took place in 2010. In September 2010, the producer started to use the technology with secondary YARA catalyst and use of continuous emission monitoring system. It resulted in significant decrease of N<sub>2</sub>O emissions.

The same EF was also used for the other medium-pressure plant in the Slovak Republic. The used medium-pressure technologies are very similar. The later medium-pressure plant changed owner at the end of the year 2012, and the plant was modernized in the same way as the other plant (secondary catalyst and continuously monitoring of N<sub>2</sub>O emissions).

- **High-pressure EFs:** The high-pressure plant started its production in 1999. The N<sub>2</sub>O emission factor in high-pressure plant was directly measured in 2006 and 2007 (9.02 kg N<sub>2</sub>O/1 t of HNO<sub>3</sub>). This value was then used for the whole time series for the high-pressure technology. It is very close to the IPCC default value (9 kg/t). In September 2010, the producer introduced the technology with secondary YARA catalyst and continuous emission monitoring system. It resulted in significant decrease of N<sub>2</sub>O emissions.

The detailed information on N<sub>2</sub>O emission factors from the nitric acid production in 2022 is presented in [Table 4.19](#). The overall EF = 0.370 kg N<sub>2</sub>O/t of HNO<sub>3</sub> in 2023 was estimated as weighted average. N<sub>2</sub>O emissions were 176.32 t in 2023. The detailed results are in [Tables 4.19](#) and [4.20](#).

**Table 4.19: Detailed information on measured N<sub>2</sub>O concentrations and EFs in 2023**

PLANT	N <sub>2</sub> O Concentration	Weighted Average EF
	ppm	kg/t
Medium Pressure Plant 1	133.34	0.481
Medium Pressure Plant 2	64.83	0.247
High Pressure Plant	87.09	0.324



**Table 4.20: Estimated N<sub>2</sub>O emissions and IEFs (N<sub>2</sub>O) in particular years**

YEAR	HNO <sub>3</sub> Production	EF N <sub>2</sub> O	N <sub>2</sub> O Atmospheric	N <sub>2</sub> O Medium Pressure	N <sub>2</sub> O High Pressure	TOTAL N <sub>2</sub> O Emissions
	kt	kg/t HNO <sub>3</sub>	t			
1990	400.54	10.106	2 170.86	1 876.88	NO	4 310.94
1995	398.80	9.936	2 020.78	1 941.77	NO	4 250.00
2000	407.22	8.383	NO	1 256.58	2 157.06	3 413.64
2005	497.68	8.326	NO	1 584.29	2 559.28	4 143.57
2010	510.97	5.706	NO	1 393.18	1 522.15	2 915.33
2011	593.75	2.288	NO	739.54	618.68	1 358.22
2012	550.51	1.770	NO	587.81	386.52	974.33
2013	611.65	0.710	NO	136.50	297.76	434.26
2014	580.09	0.837	NO	156.40	329.13	485.53
2015	634.31	0.740	NO	95.27	373.80	469.07
2016	568.55	0.716	NO	71.69	335.45	407.14
2017	646.23	0.545	NO	118.87	233.42	352.28
2018	575.32	0.616	NO	127.84	226.32	354.16
2019	571.27	0.532	NO	120.23	183.86	304.09
2020	580.24	0.439	NO	125.42	129.44	254.85
2021	636.32	0.336	NO	87.29	126.42	213.72
2022	523.76	0.380	NO	77.18	121.91	199.09
2023	476.13	0.370	NO	95.79	80.53	176.32

### Uncertainties and Time-series Consistency

There is only one owner, which has been operating several nitric acid production plants in Slovakia since 2012. Nitric acid is produced in two medium- and one high-pressure plants. Until 1999, also atmospheric-pressure plant had been operated in Slovakia. The plant specific emission factors are used for medium and high-pressure technologies since 1990.

The emission factors for medium-pressure plant are based on the measured data in 2005, 2008, 2009 and 2010. The average value (7.5 kg/1 t of HNO<sub>3</sub>) of EF is used for other years of time series except the years 2006 and 2007. According to the N<sub>2</sub>O emissions measured in 2006 and 2007, the EF was 10.332 kg/1 t of HNO<sub>3</sub> (malfunction in the plant). The emission factor for high-pressure plant was measured to be 9.02 kg/1 t of HNO<sub>3</sub> which is in good agreement with the IPCC default EF for this type of technology (9 kg/1 t of HNO<sub>3</sub>). The same value was used in 1990 – 2010, when the high-pressure production of nitric acid occurred.

In September 2010, technology was changed in medium- and high-pressure technologies by one producer. The secondary YARA catalyst was introduced, which resulted in significant decrease of N<sub>2</sub>O emissions since 2010. The second plant was using un-modified technology and EF equalled 7.5 kg/1 t of HNO<sub>3</sub>. At the end of 2012, the second medium-pressure plant was bought by the new owner (already owned the second plant). The plant was modernized in the same way as the other (secondary catalyst and continuously monitoring of N<sub>2</sub>O emissions) and emission factor was improved. In the end of 2020, there was another modernization of the one of the medium-pressure plants. During the modernization, the number of catalyst layers was increased that resulted in another decrease of N<sub>2</sub>O emissions. The decrease was fully evident in 2021.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth

year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.8.3. Adipic Acid Production (CRT 2.B.3)

Adipic acid is not produced in the Slovak Republic therefore notation key “NO” was used.

### 4.8.4. Caprolactam, Glyoxal and Glyoxylic Acid (CRT 2.B.4)

None of these products are produced in the Slovak Republic therefore notation key “NO” was used.

### 4.8.5. Carbide Production (CRT 2.B.5)

#### Silicon Carbide (CRT 2.B.5.a)

Silicon carbide is not produced in the Slovak Republic therefore notation key “NO” was used.

#### Calcium Carbide (CRT 2.B.5.b)

Calcium carbide (the correct chemical name of this compound is calcium acetylide) is produced by the reaction of CaO and coke in submerged arc furnace. The final CO<sub>2</sub> emissions balance is influenced by export of carbide, use of carbide in Slovakia and use of limestone. Total CO<sub>2</sub> emissions reached 31.82 Gg of CO<sub>2</sub> in 2023 and decreased by 10% in comparison with 2022. It corresponds to the increased export from Slovakia. Almost whole produced calcium carbide was exported and emissions from its use were very low. Since 2015, the calcinated anthracite is used instead of other bituminous coal. Methane emissions from calcium carbide production is not occurring, notation key NO is used.

#### Methodological Issues

Carbon balance of all input-output flows was used. The method is similar to tier 3 method according to the IPCC 2006 GL with the combination of country specific emission factors and NCVs. These EFs are updated annually. The CO<sub>2</sub> emissions are calculated from the coal consumption (reduction step), limestone use, and products use. Limestone has not been used since 2011. The CO<sub>2</sub> emissions from reduction step are calculated in the following way: CO<sub>2</sub> emissions = (Σ(consumption of coal x NCV x EF(C))-(carbide production x C content in carbide)) x 44/12.

Acetylene is produced in the plant not only for welding application. A part of produced acetylene can be used to produce the vinyl chloride monomer. When this occurs, the CO<sub>2</sub> emissions from this production are reported in 2.B.8.c (ethylene dichloride and vinyl chloride monomer (VCM)). The calcium carbide for acetylene production for welding application was calculated by conservative approach, as follows: Calcium carbide for welding = import + production – export – calcium carbide for VCM.

Results of CO<sub>2</sub> emissions from non-exported production are summarized in **Table 4.21** (C = confidential data are available in the SNE archive).

**Table 4.21: Estimated CO<sub>2</sub> emissions, carbide production and export in particular years**

YEAR	Carbide Prod.	Carbide Export-Import	Carbide for VCM Prod.	CaCO <sub>3</sub> Consum.	Coking Coal Consum.	Other Bituminous Coal Consum.	IEF CO <sub>2</sub>	CO <sub>2</sub>
	kt						t/t	Gg
1990	NO	NO	NO	NO	NO	NO	NO	NO
1995	84.30	C	C	131.63	66.61	7.14	1.65	139.01
2000	88.82	C	C	138.68	70.26	7.44	1.76	156.73
2005	97.03	C	C	151.50	76.73	8.15	1.82	176.72
2010	98.26	C	C	158.17	77.69	8.28	2.01	197.56
2011	107.40	C	C	172.89	84.89	9.07	2.07	222.28
2012	100.48	C	C	NO	79.44	8.46	1.41	141.26
2013	81.79	C	C	NO	60.93	6.16	1.17	95.35

YEAR	Carbide Prod.	Carbide Export-Import	Carbide for VCM Prod.	CaCO <sub>3</sub> Consum.	Coking Coal Consum.	Other Bituminous Coal Consum.	IEF CO <sub>2</sub>	CO <sub>2</sub>
	kt						t/t	Gg
2014	74.30	C	C	NO	57.99	4.34	1.15	85.76
2015	56.18	C	C	NO	41.05	3.55*	0.86	48.47
2016	67.95	C	C	NO	48.01	4.50*	0.93	63.16
2017	71.64	C	C	NO	47.82	5.08*	0.83	59.35
2018	70.15	C	C	NO	48.30	4.79*	0.97	68.26
2019	60.47	C	NO	NO	45.90	3.49*	1.03	62.05
2020	47.61	C	NO	NO	38.07	1.65*	0.96	45.83
2021	48.48	C	NO	NO	39.01	1.47	0.98	47.57
2022	36.12	C	NO	NO	27.97	1.83	0.98	35.49
2023	36.15	C	NO	NO	28.70	1.33	0.88	31.82

\* calcinated anthracite

Implied CO<sub>2</sub> emission factors of carbide production are updated annually based on the annual values of the NCV and EFs of used fuels and carbon content in the products (calcium carbide). Implied CO<sub>2</sub> emission factor in 2023 was 0.88 t CO<sub>2</sub>/t of produced CaC<sub>2</sub> (only from technological process, no acetylene production is included). When the acetylene production for welding application is included in formula, the IEF was the same due no acetylene production in 2023.

According to the direct information provided by the producers, a part of produced calcium carbide was exported from the Slovak Republic and another part of calcium carbide was used for acetylene and following vinyl chloride production (not in 2023). No calcium carbide was imported to Slovakia in 2023. The rest of produced calcium carbide was used for acetylene production for welding applications (conservative approach). No production of CaO occurred in 2023. The CaO was bought from the lime producers and approximately 40% of CaO was imported from neighbouring countries. Therefore, no CO<sub>2</sub> emissions from CaO preparation (limestone decomposition step) were allocated in this category. Since 2015, calcinated anthracite is used for the production of Søderberg anodes. The content of carbon in this type of material is declared min. 95%, for ensuring conservatism the assumption of 100% content of carbon is used for the calculation of emission estimates.

#### Uncertainties and time-series consistency

The production of calcium carbide in Slovakia started in 1992. Since that year, consistent methodology and tier method has been used for the whole time series for emissions estimation. Fluctuations and outliers in emission trend (1998, 2002, 2011 – 2020) and emission factors were caused by differences in exported volume of final calcium carbide and utilization of CaC<sub>2</sub> for acetylene and following VCM production ([Table 4.21](#)). The CaO production finished in 2011. In the present, the CaO is produced by the lime producers and bought for carbide production.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.8.6. Titanium Dioxide Production (CRT 2.B.6)

Titanium dioxide is not produced in the Slovak Republic and “NO” notation key was used.

#### 4.8.7. Soda Ash Production (CRT 2.B.7)

Soda ash is not produced in the Slovak Republic and “NO” notation key was used.

#### 4.8.8. Petrochemical and Carbon Black Production (CRT 2.B.8)

Methanol (CRT 2.B.8.a), ethylene oxide (CRT 2.B.8.d), acrylonitrile (CRT 2.B.8.e) and carbon black (CRT 2.B.8.f) are not produced in the Slovak Republic and “NO” notation keys were used.


##### Ethylene (CRT 2.B.8.b)

Ethylene is produced by steam cracking of naphtha in the Slovak Republic. Natural gas, refinery gas and low-pressurized methane are used as the other feedstock in the process. Propylene is a valuable co-product of the process. The other by-products and off-gases are transferred into refinery and they are reported in 1.A.1.b. Total CO<sub>2</sub> emissions from ethylene production were 373.21 Gg in 2023, which is higher by 8% compared with previous year. The increase is caused by increasing the production.

##### Methodological Issues

Carbon balance approach (tier 2), as described in the IPCC 2006 GL, was used. All feedstock (naphtha, natural gas, refinery gas and low-pressurized methane) and all products (ethylene, propylene, and other chemicals – by-products) are balanced ([Figure 4.6](#)). Methane emissions do not occur when using approach described in the IPCC 2006 GL.

**Figure 4.6:** Scheme of carbon material balance used in for 2.B.8.b



Input streams as naphtha and refinery gas originates in the refinery. During the reaction in the ethylene unit, a refinery gas with high content of methane is formed. This methane is separated from the refinery gas and creates an inner loop in the process. The rest of the refinery gas (after separating of methane) is going into refinery and it represents an input stream for emission estimates in the Energy sector (1.A.1.b category). In refinery, other chemicals as butadiene *etc.* are separated and off-gases are burned. The burning of off-gases is reported in the Energy sector (1.A.1.b category). The data “Carbon in other chemicals” presented in [Table 4.22](#) represents carbon outgoing from ethylene unit (due to many the other produced chemicals, total carbon content is reported). From this amount, the low pressurized methane is separated, while the rest is going into refinery. On the other hand, another stream of refinery gas is outgoing from refinery and it represents the input stream in the ethylene unit.

The total amount of carbon excluded from reference approach is the difference between carbon contained in input flows (naphtha, excess refinery gas, natural gas) and carbon in off-gases going to the refinery. Part of it is stored in products (ethylene and propylene) and the rest is evolved as CO<sub>2</sub> emissions. This approach (including the inner loop into the calculation) is chosen because of

comparability with the EU ETS report where the emission estimates are calculated based on the fuel combustion. The methodology used was also published in the paper by Eva Krtková et. al.<sup>2</sup>).

Ethylene is produced by one operator in Slovakia and therefore the NCVs and emission factors of all feedstock were provided directly (EU ETS reports). Total production of ethylene and propylene was provided by the plant operator. Detailed data are presented in **Table 4.22**.

**Table 4.22: Activity data and related CO<sub>2</sub> emissions from ethylene and propylene production in particular years**

YEAR	Naphtha	Natural Gas	Refinery Gas	Low-Pressurized CH <sub>4</sub>
	Inputs in TJ			
1990	14 867.6	3 074.8	4 366.1	0.0
1995	19 271.2	1 714.1	5 071.7	1 306.4
2000	21 625.6	1 419.9	4 380.5	2 357.3
2005	17 440.0	959.5	4 497.4	1 031.8
2010	17 004.0	1 610.6	4 237.1	1 244.2
2011	16 742.4	1 532.7	4 062.2	1 126.2
2012	10 900.0	1 487.9	2 928.5	612.1
2013	11 510.4	1 707.9	3 124.8	907.5
2014	11 264.0	1 319.6	2 522.0	584.2
2015	14 916.0	1 123.8	3 707.6	1 079.9
2016	10 472.0	1 150.2	3 584.5	1 250.4
2017	11 176.0	1 290.4	3 702.3	1 363.0
2018	13 948.0	1 355.5	4 105.8	1 718.6
2019	13 244.0	1 182.9	3 624.6	1 432.9
2020	17 732.0	909.9	4 081.9	2 004.0
2021	19 184.0	728.6	4 533.3	1 968.7
2022	16 588.0	535.6	4 156.6	1 700.7
2023	16 764.0	489.4	4 690.6	1 702.6

YEAR	Ethylene Production	Propylene Production	Carbon In Other Chem.	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	Outputs in kt			Gg	t/t
1990	216.5	98.6	27.3	416.80	1.925
1995	200.3	93.3	133.9	447.80	2.236
2000	207.4	92.9	175.5	449.28	2.166
2005	202.5	91.9	96.8	357.33	1.765
2010	197.0	93.0	91.8	391.16	1.986
2011	194.0	96.0	86.6	411.73	2.122
2012	128.0	68.0	50.2	306.42	2.394
2013	145.5	71.7	44.3	322.24	2.215
2014	102.8	55.2	90.1	243.55	2.369
2015	137.0	67.0	123.7	323.91	2.364
2016	146.0	71.0	23.7	328.16	2.248
2017	176.0	84.0	0.9	348.90	1.982
2018	198.0	98.0	25.6	391.74	1.978
2019	169.5	81.9	49.7	340.24	2.008

<sup>2</sup> Eva Krtková, Vladimír Danielík, Janka Szemesová, Klára Tarczay, Gábor Kis-Kovács and Vladimír Neužil, Non-Energy Use of Fuels in the Greenhouse Gas Emission Reporting, Atmosphere 2019, 10, 406; DOI: <https://www.mdpi.com/2073-4433/10/7>

YEAR	Ethylene Production	Propylene Production	Carbon In Other Chem.	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	Outputs in kt			Gg	t/t
2020	207.9	153.9	44.9	379.68	1.826
2021	213.6	165.5	59.1	391.28	1.832
2022	169.3	217.6	0.4	345.20	2.039
2023	190.0	144.3	49.1	373.21	1.964

### Uncertainties and Time-series Consistency

Consistent methodology based on tier 2 approach was used for the whole-time series since 1990. Fluctuations and outliers in emission trend were caused by the different amounts of other chemicals produced in process and by the different share of fuels (naphtha, NG, refinery gas, low-pressured methane). Fluctuations in IEF are caused by relating of the IEF to the production of ethylene only, while there is a varied share of the different products produced during the time series. Sensitivity of time series is caused by the limited number of operators produced in Slovakia and their actual activity. The corresponding volume of natural gas and other fuels presented in [Table 4.22](#) were subtracted from 1.A.2.c in the Energy sector.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### Ethylene Dichloride and Vinyl Chloride Monomer (CRT 2.B.8.c)

Ethylene dichloride (EDC) is produced by direct chlorination process in the Slovak Republic. Cracking of ethylene dichloride results in vinyl chloride monomer (VCM) and HCl. The HCl is consumed in the reaction with acetylene that results in another amount of vinyl chloride monomer. This amount of consumed acetylene is not reported in 2.B.5.b (calcium carbide production) to avoid double counting. Total CO<sub>2</sub> emissions from ethylene dichloride and vinyl chloride monomer production were estimated in this category for whole time series. The emissions were 3.26 Gg in 2023 and increased by ca 24% in comparison with the previous year 2022. It corresponds to the decrease in production.

### Methodological Issues

Tier 2 approach and carbon balance approach, as described in IPCC 2006 GL was used. The used approach is described on the following scheme ([Figure 4.7](#)).

Ethylene dichloride and vinyl chloride monomer is produced by one operator in Slovakia. All streams (inputs) shown on [Figure 4.7](#) were taken into account with respective emission factors and contents of carbon (plant specific data). These parameters were updated annually.

Total production of vinyl chloride monomer and the production of ethylene dichloride (a part of it that is a final product, not intermediate for VCM) were delivered directly by the plant operator. Information on streams inputs and outputs material balance are summarized in [Table 4.23](#).

Figure 4.7: Carbon material balance used in emissions estimation of the category 2.B.8.c




Table 4.23: Activity data and related CO<sub>2</sub> emissions from the EDC and VCM production in particular years

YEAR	Natural Gas Consumption	Ethylene Consumption	Acetylene Consumption	EDC Production*	VCM Production
	1 000 m <sup>3</sup>	kt			
1990	5 084	10.320	14.313	NO	55.536
1995	4 935	17.356	8.177	NO	56.159
2000	5 302	21.003	9.471	NO	66.963
2005	5 850	18.807	9.166	NO	61.568
2010	5 272	17.448	5.743	0.893	50.085
2011	5 872	19.294	5.772	1.150	53.928
2012	5 475	18.149	2.587	0.712	44.300
2013	3 548	11.915	3.462	0.666	33.059
2014	3 013	10.148	3.068	1.172	28.185
2015	3 174	10.816	3.486	-0.158	31.127
2016	4 694	11.762	6.357	1.571	39.484
2017	3 505	10.612	5.703	0.305	35.193
2018	4 030	8.970	2.810	0.502	26.295
2019	405	6.933	NO	0.348	12.957
2020	626	3.203	NO	-0.323	6.770
2021	1 754	7.132	NO	0.688	14.828
2022	1 539	7.415	NO	0.436	15.167
2023	1 240	5.521	NO	0.475	11.344

YEAR	Gas for Flaring	CHCl**	CH***	Proc. CO <sub>2</sub>	Combust. CO <sub>2</sub>	Flaring CO <sub>2</sub>	Total CO <sub>2</sub>	IEF (CO <sub>2</sub> )
	1 000 m <sup>3</sup>	kt						
1990	43.9	1.587	0.282	10.382	1.449	0.173	12.004	0.2161
1995	50.7	2.042	0.284	10.045	1.866	0.199	12.110	0.2156
2000	53.4	2.104	0.265	11.264	1.922	0.210	13.396	0.2000
2005	44.8	2.397	0.268	11.704	2.190	0.176	14.070	0.2285
2010	45.3	1.862	0.271	10.703	1.701	0.178	12.583	0.2512

YEAR	Gas for Flaring	CHCl**	CH***	Proc. CO <sub>2</sub>	Combust. CO <sub>2</sub>	Flaring CO <sub>2</sub>	Total CO <sub>2</sub>	IEF (CO <sub>2</sub> )
	1 000 m <sup>3</sup>	kt		Gg			t/t VMC	
2011	51.9	2.114	0.269	11.883	1.932	0.204	14.019	0.2600
2012	50.5	1.621	0.297	11.160	1.481	0.198	12.839	0.2898
2013	50.2	0.936	0.206	7.491	0.855	0.197	8.543	0.2584
2014	24.8	0.903	0.234	6.194	0.769	0.097	7.051	0.2502
2015	24.0	0.778	0.269	7.103	0.714	0.094	7.911	0.2541
2016	99.2	1.095	0.426	8.629	1.041	0.390	10.061	0.2548
2017	128.2	1.315	0.536	7.170	1.269	0.504	8.942	0.2541
2018	132.5	0.852	0.288	6.374	0.521	0.777	7.672	0.2918
2019	58.2	0.639	0.078	3.193	0.229	0.478	3.900	0.3010
2020	110.5	0.323	0.048	1.293	0.435	0.248	1.975	0.2918
2021	202.7	0.561	0.087	2.982	0.797	0.434	4.214	0.2842
2022	143.6	0.666	0.117	3.335	0.565	0.526	4.425	0.2918
2023	110.6	0.439	0.115	2.447	0.435	0.375	3.257	0.2871

\*production of EDC that is used as a product, not an intermediate to VCM; \*\*chlorinated hydrocarbons; \*\*\*other hydrocarbons

### Uncertainties and Time-series Consistency

Consistent methodology and tier method are used for the whole time series since the base year. Fluctuations and outliers in emissions and IEFs are caused by different amounts of vinyl chloride monomer produced by two methods (from ethylene and/or acetylene). Sensitivity of time series are caused also by the limited number of operators produced in Slovakia and their actual activity or production capacity. The respective amounts of natural gas were subtracted from 1.A.2.c of the Energy sector. It should be mentioned that the negative value of EDC production in 2015 and in 2020 means the using of stocked or bought amount of EDC. Not enough EDC was produced in the plant in those years for the purpose of VCM production.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.8.9. Hydrogen Production (CRT 2.B.10)

In this submission, 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories was fully implemented. In Vol. 3 – Introduction, Chapter 1.3.3, page 1.12 it is stated: *“Refineries manufacture petroleum products for fuel and for non-energy uses, and in doing so produce hydrogen and other gases, intermediate products and basic chemicals. The CO<sub>2</sub> emissions from fuel consumed by the refinery for this activity are reported as Energy Sector emissions. This principle is maintained in the Guidelines even when some fuel use in the refinery is to support manufacture of chemicals for sale (for example, propylene or aromatics). In the 2019 Refinement, this principle is reiterated within the new guidance presented for hydrogen production, which is a new IPPU source category; the emissions from hydrogen production within a refinery as an intermediate product are primarily to support Energy sector activities, with emissions to be reported in the Energy sector.”*

Until now, the hydrogen production in refinery was included in this category. Based on the above cited paragraph, the hydrogen production is allocated into Energy sector and “NO” notation key is used.



## 4.9. Metal Production (CRT 2.C)


This category produces emissions of CO<sub>2</sub>, CH<sub>4</sub> and PFCs emissions (Aluminium Production). Total emissions were 3 765.58 Gg of CO<sub>2</sub> eq. in 2023; the increase was 7% when compared with 2022 due to the significant increase of Iron and Steel production. Comparing with the base year, the emissions are lower by 22%. However, more efficient production results in significantly higher iron and steel production at the same emission production. According to the IPCC 2006 GL, also zinc and lead production are reported in 2.C.5 Lead Production and 2.C.6 Zinc Production.

**Table 4.24:** Emissions in the category Metal Production 2.C in particular years

YEAR	2.C.1 Iron and Steel	2.C.2 Ferroalloy Production	2.C.3 Aluminium Production	2.C.5 Lead Production	2.C.6 Zinc Production
	Gg of CO <sub>2</sub> eq.				
1990	4 182.73	296.74	335.24	NO	NO
1995	4 335.91	235.64	148.83	NO	NO
2000	3 359.30	182.72	193.11	NO	NO
2005	3 924.83	228.22	281.31	NO	NO
2010	4 107.71	220.01	291.75	NO	NO
2011	3 505.23	203.04	285.90	0.01	NO
2012	3 878.02	266.53	288.14	0.04	0.02
2013	3 781.27	166.19	282.26	0.05	0.01
2014	4 070.29	224.30	284.27	0.06	0.01
2015	4 045.76	241.03	292.86	0.06	NO
2016	4 353.72	238.14	286.59	0.06	NO
2017	4 346.99	295.50	290.76	0.06	NO
2018	4 206.59	283.05	291.66	0.01	NO
2019	3 569.05	240.13	288.99	0.01	NO
2020	3 159.09	215.56	251.93	0.03	NO
2021	4 294.18	253.13	272.87	0.03	NO
2022	3 339.31	76.35	117.33	0.08	NO
2023	3 750.71	7.90	6.89	0.08	NO

The major share of emissions (99.6%) belongs to the iron and steel production, 0.2% belongs to the ferroalloy production and 0.2% to the aluminium production. Other subcategories are not significant in emission share within the category 2.C. The share is influenced by significant decrease in ferroalloys production and closing production of aluminium in 2023.

**Figure 4.8:** The share in GHG emissions in the category 2.C by subcategories in 2023




### 4.9.1. Iron and Steel Production (CRT 2.C.1)

Total emissions in this category were 3 750.71 Gg CO<sub>2</sub> eq. in 2023, higher by 12% when compared with the year 2022. The reason of such increase is the significant increase in production due to the economic reasons. Comparing the base year, the decrease was 10%. Pig iron is produced by the reduction process of iron ore with coke in a blast furnace. The major emissions emitted from this process are CO<sub>2</sub> emissions. Limestone is added as an agent for slag formation. Pig iron contains about 4% of carbon and a part of this carbon is oxidized in the next step. This process is accompanied by the CO emissions release. The most of CO is burned to CO<sub>2</sub>. Iron ore was processed to pig iron. Category iron and steel production includes following processes: (i) steel production (2.C.1.a), (ii) pig iron production (2.C.1.b), (iii) sinter production (2.C.1.d) and (iv) steel production in electric arc furnaces (EAF) (2.C.1.f). Major sources of technological CO<sub>2</sub> emissions are pig iron and steel production in blast furnaces. Due to the difficult disaggregation between emissions originated from pig iron and from steel production, total CO<sub>2</sub> emissions from total production processes were allocated directly in steel production category. Therefore, the notation key “IE” was used in the other categories. The CO<sub>2</sub> emissions from the EAF steel production are reported separately in 2.C.1.f.

Implementation of 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories resulted in changes in the inventory: (i) methane emissions from sinter and coke production. Due to the technological scheme of the process ([Figure A4.1.1](#) in the [Annex 4.1](#) of this Document) we report the methane emission from coke production in IPPU sector instead of Energy sector; (ii) nitrous oxide emissions from the flaring of blast furnace gas and converter gas. The methane and nitrous oxide emissions were estimated back to the 1990.

**Figure 4.9:** Emission trend in the category 2.C.1 according to type of production in 1990 – 2023



#### Methodological Issues

Pig iron and steel are produced mainly in blast furnaces and by the EAF processes. Technological emissions from pig iron (2.C.1.b), steel (2.C.1.a) and emissions from coke electrodes used by the EAF steel production (2.C.1.f) are included in this category. Due to application of tier 2 method, methane emissions were not balanced until now.

The plant with blast furnaces is one complex with many energy-related installations (coke ovens, heating plant, manufacturing of steel products, etc.). After direct discussion with the plant operators, simplified scheme of the plant in order to carbon balance was proposed ([Figure A4.1.1](#) in the [Annex 4.1](#) of this Document).

All streams were calculated based on the plant specific conversion units and carbon EFs or based on carbon content of iron ore and steel. Carbon balance of iron and steel production is described in full details in the [Annex 4.1](#). The used method corresponds to tier 2 for CO<sub>2</sub> as described in the IPCC 2006 GL. Methane emissions from sinter and coke production are calculated using Tier 1 method and reported

in 2.C.1.a. No data for higher Tier are available. Nitrous oxide emissions from blast furnace gas and converter gas flaring are calculated using Tier 1 method, as well. No data for higher Tier are available. However, CRT Tables are not prepared for reporting of N<sub>2</sub>O emissions from iron and steel production, Therefore, they are reported separately in artificially created category 2.C.7.i Iron and Steel – N<sub>2</sub>O.

The CO<sub>2</sub> emissions were calculated by using following equation:

$$E(\text{steelBF}) = (\sum(\text{massofCininputstream}_i) - \sum(\text{massofCinoutputstream}_i)) \cdot \frac{44}{12}$$

$$E(\text{steelEAF}) = EF(\text{steelinEAF}) \cdot \text{massofSteelproducedinEAF}$$

$$\text{TotalEmissions} = \sum_i E(i)$$

The methane emissions were calculated using the equation:

$$\text{TotalEmissions} = \text{QuantityOfSinterProduced} \cdot EF(\text{sinter}) + \text{QuantityOfCokeProduced} \cdot EF(\text{sinter})$$

where  $EF(\text{sinter}) = 0.07$  kg/t and  $EF(\text{coke}) = 0.089$  kg/t.

The N<sub>2</sub>O emissions were calculated using the equation:

$$\text{TotalEmissions} = \text{BFGflared} \cdot EF(\text{BFG}) + \text{ConverterGasFlared} \cdot EF(\text{converter})$$

where  $EF(\text{BFG}) = 5.6 \times 10^{-7}$  t/TJ and  $EF(\text{converter}) = 4.0 \times 10^{-7}$  t/TJ.

EFs of CO<sub>2</sub> are estimated annually on plant level, what is equal to country specific level in this case. Inter-annual fluctuations in emission factors are caused by two basic technological situations:

- different volume of iron scrap is added to the charge in steel making process,
- different amounts of gas fuels are produced in blast furnaces.

The average content of carbon in iron ore was 3.552 kg/t, in pig iron it was 1.991 kg/t and 0.890 kg/t in steel (data supplied directly) in 2023. Emission factors and other parameters are summarized in [Tables 4.25-4.27](#). The CO<sub>2</sub> emissions from the EAF process are estimated based on carbon balance, where all material flows are considered. Methane and nitrous oxide emissions for the whole time series are summarized in [Tables 4.28-4.29](#).

Iron and steel is produced by several plants (U.S.Steel Košice, a. s., UNEX Prakovce, Metalurg, Slovakia Steel Mills and by Ironworks Železiarne Podbrezová, a. s.). The manufacturer of iron and steel in blast furnaces (integrated production of iron and steel) produced pig iron (part of which was sold and not processed to steel) and 4 074.96 kt of steel in 2023 (increase by 14% when compared with 2022). Total production of steel produced by the EAF technology was 244.94 kt in 2023. The plant UNEX Prakovce did not produce steel since 2013. New plant, Slovakia Steel Mills, started their production by the EAF technology in 2013. However, due to the sanctions to the Russian Federation, its production decreased and, in the end of 2014 the production was stopped. Since 2019, only one plant using EAF technology is in operation. Activity data on produced pig iron, what is sold to customers and not processed to steel are presented in 2.C.1.b. These data are presented without emissions because these emissions are balanced together with steel production and allocated in the 2.C.1.a.

**Table 4.25: Activity data, emission factors and CO<sub>2</sub> emissions in integrated iron and steel production in 2005 – 2023 that are reported in 2.C.1.a subcategory**

YEAR	Coal Cons.	Coke	NG Cons.	CG Output	BFG Output	Steel Prod.	Limestone Used	CO <sub>2</sub>	IEF (CO <sub>2</sub> )
	kt		mil. M <sup>3</sup>	kt			Gg	t/t	
2005	2 594.52	-20.00	30.67	626.30	3 622.84	4 238.12	829.34	3 893.90	0.919
2006	2 853.64	179.00	37.68	670.28	4 665.12	4 836.49	781.85	4 391.72	0.908
2007	2 960.17	-147.00	26.31	682.77	3 838.94	4 784.81	606.74	4 140.88	0.865
2008	2 867.21	-152.00	22.11	668.56	3 693.60	4 229.40	464.33	3 992.89	0.944
2009	2 455.88	-85.00	20.27	592.13	3 378.26	3 642.28	518.34	3 479.24	0.955
2010	2 516.80	327.63	36.14	657.13	4 227.88	4 401.78	640.47	4 071.97	0.925
2011	2 503.00	-27.00	41.18	645.28	4 025.42	3 961.02	600.73	3 461.85	0.874
2012	2 709.17	-22.00	24.89	618.32	4 135.38	4 236.19	622.03	3 842.85	0.907

YEAR	Coal Cons.	Coke	NG Cons.	CG Output	BFG Output	Steel Prod.	Limestone Used	CO <sub>2</sub>	IEF (CO <sub>2</sub> )
	kt		mil. M <sup>3</sup>	kt			Gg	t/t	
2013	2 482.48	-13.97	22.25	591.42	3 867.60	4 344.25	820.30	3 708.94	0.854
2014	2 606.36	74.98	20.13	604.21	3 958.03	4 439.48	973.80	4 024.91	0.907
2015	2 641.87	-29.98	20.18	657.42	3 586.84	4 310.94	800.39	4 018.99	0.932
2016	2 626.27	99.39	23.31	649.04	3 703.90	4 599.44	942.05	4 326.18	0.941
2017	2 650.44	150.69	19.37	784.45	3 894.35	4 712.96	961.71	4 319.01	0.916
2018	2 637.44	176.76	20.67	792.90	4 097.63	4 641.84	957.39	4 177.19	0.900
2019	2 279.01	28.15	21.04	549.83	3 018.73	3 608.95	749.44	3 543.54	0.982
2020	1 914.80	92.42	17.15	477.18	2 687.21	3 119.01	650.71	3 136.29	1.006
2021	2 677.74	108.19	18.60	578.35	4 002.59	4 560.37	963.27	4 264.48	0.935
2022	2 313.60	-41.41	18.08	538.54	3 197.69	3 558.10	767.63	3 313.75	0.931
2023	2 362.94	186.77	21.47	564.47	3 832.19	4 074.97	805.88	3730.20	0.915

CG = coking gas, BFG = blast furnace gas, con. = consumption, prod. = production

**Table 4.26: Production and CO<sub>2</sub> emissions in steel industry in 1990 – 2004**

YEAR	Steel Production	Limestone Used	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	kt		Gg	t/t
1990	3 561.50	615.78	4 149.82	1.165
1991	3 163.40	540.44	3 015.13	0.953
1992	2 952.40	501.77	2 639.86	0.894
1993	3 205.40	555.13	4 337.65	1.353
1994	3 330.40	581.39	3 815.70	1.146
1995	3 207.40	562.16	4 304.41	1.342
1996	2 920.00	508.61	4 533.89	1.553
1997	3 072.30	542.47	4 547.00	1.480
1998	3 100.00	541.86	4 075.07	1.315
1999	3 420.00	527.61	3 967.28	1.160
2000	3 519.99	713.79	3 326.23	0.945
2001	3 751.85	660.08	3 356.97	0.895
2002	4 103.20	575.05	4 129.07	1.006
2003	4 382.92	608.29	3 956.26	0.903
2004	4 421.14	1 154.75	4 273.53	0.967

**Table 4.27: Activity data, emission factors (below) and CO<sub>2</sub> emissions in individual plants with EAF steel production in particular years**

YEAR	ŽELEZIARNE PODBREZOVÁ			SLOVAKIA STEEL MILLS			METALURG STEEL		
	Steel by EAF	Carbon	CO <sub>2</sub>	Steel by EAF	Carbon	CO <sub>2</sub>	Steel by EAF	Carbon	CO <sub>2</sub>
			Gg			Gg			Gg
1990	C	3.81	13.97	NO	NO	NO	C	1.10	4.02
1995	C	3.88	14.22	NO	NO	NO	C	1.04	3.83
2000	C	3.88	14.22	NO	NO	NO	C	1.12	4.10
2005	C	3.41	12.49	NO	NO	NO	C	0.24	0.89
2010	C	4.47	16.37	NO	NO	NO	C	0.34	1.23
2011	C	7.06	25.88	NO	NO	NO	C	0.30	1.09
2012	C	4.64	17.00	NO	NO	NO	C	0.17	0.62
2013	C	3.97	14.55	C	10.85	39.80	C	0.00	0.01
2014	C	3.00	11.01	C	4.21	15.43	C	0.01	0.05
2015	C	2.49	9.14	NO	NO	NO	NO	NO	NO
2016	C	2.39	8.78	NO	NO	NO	C	0.01	0.04

YEAR	ŽELEZIARNE PODBREZOVÁ			SLOVAKIA STEEL MILLS			METALURG STEEL		
	Steel by EAF	Carbon	CO <sub>2</sub>	Steel by EAF	Carbon	CO <sub>2</sub>	Steel by EAF	Carbon	CO <sub>2</sub>
			Gg			Gg			Gg
2017	C	2.38	8.73	NO	NO	NO	C	0.08	0.28
2018	C	2.83	10.35	NO	NO	NO	C	0.08	0.28
2019	C	2.93	10.74	NO	NO	NO	NO	NO	NO
2020	C	2.60	9.53	NO	NO	NO	NO	NO	NO
2021	C	2.84	10.41	NO	NO	NO	NO	NO	NO
2022	C	2.82	10.35	NO	NO	NO	NO	NO	NO
2023	C	1.02	3.73	NO	NO	NO	NO	NO	NO

YEAR	UNEX, PRAKOVCE		TOTAL		
	Steel by EAF	CO <sub>2</sub>	Steel by EAF	CO <sub>2</sub>	IEF
	kt	Gg	kt	Gg	t/t
1990	C	0.16	310.73	18.15	0.0584
1995	C	0.16	314.64	18.21	0.0579
2000	C	0.17	316.36	18.49	0.0584
2005	C	0.08	356.90	13.46	0.0377
2010	NO	NO	331.25	17.60	0.0531
2011	NO	NO	374.22	26.97	0.0721
2012	NO	NO	372.40	17.62	0.0473
2013	NO	NO	711.34	54.36	0.0764
2014	NO	NO	527.85	26.49	0.0502
2015	NO	NO	315.05	9.14	0.0290
2016	NO	NO	293.80	8.82	0.0300
2017	NO	NO	356.80	9.01	0.0253
2018	NO	NO	380.30	10.63	0.0280
2019	NO	NO	327.78	10.74	0.0328
2020	NO	NO	279.95	9.53	0.0341
2021	NO	NO	370.29	10.41	0.0281
2022	NO	NO	365.53	10.35	0.0283
2023	NO	NO	244.94	3.73	0.0152

Table 4.28: Activity data and CH<sub>4</sub> emissions from integrated iron and steel plant

YEAR	SINTER		COKE		TOTAL
	Production	CH <sub>4</sub> emission	Production	CH <sub>4</sub> emission	CH <sub>4</sub> emission
	kt				
1990	5 532.13	0.3872	1 199.29	0.1067	0.4940
1991	4 913.90	0.3440	1 065.27	0.0948	0.4388
1992	4 586.19	0.3210	994.23	0.0885	0.4095
1993	4 978.99	0.3485	1 079.38	0.0961	0.4446
1994	5 173.30	0.3621	1 121.50	0.0998	0.4619
1995	4 981.95	0.3487	1 080.02	0.0961	0.4449
1996	4 535.41	0.3175	983.22	0.0875	0.4050
1997	4 771.96	0.3340	1 034.50	0.0921	0.4261
1998	4 815.26	0.3371	1 043.89	0.0929	0.4300
1999	5 312.49	0.3719	1 151.68	0.1025	0.4744
2000	5 468.23	0.3828	1 185.44	0.1055	0.4883
2001	5 828.42	0.4080	1 263.52	0.1125	0.5204
2002	6 374.24	0.4462	1 381.85	0.1230	0.5692

YEAR	SINTER		COKE		TOTAL
	Production	CH <sub>4</sub> emission	Production	CH <sub>4</sub> emission	CH <sub>4</sub> emission
	kt				
2003	6 808.94	0.4766	1 476.09	0.1314	0.6080
2004	6 868.35	0.4808	1 488.97	0.1325	0.6133
2005	6 552.13	0.4586	1 420.42	0.1264	0.5851
2006	7 477.21	0.5234	1 620.96	0.1443	0.6677
2007	7 397.32	0.5178	1 603.64	0.1427	0.6605
2008	6 538.65	0.4577	1 417.49	0.1262	0.5839
2009	5 630.97	0.3942	1 220.72	0.1086	0.5028
2010	6 805.16	0.4764	1 475.27	0.1313	0.6077
2011	6 153.96	0.4308	1 334.10	0.1187	0.5495
2012	6 581.34	0.4607	1 426.75	0.1270	0.5877
2013	6 737.70	0.4716	1 460.64	0.1300	0.6016
2014	7 182.50	0.5028	1 446.66	0.1288	0.6315
2015	6 562.73	0.4594	1 504.28	0.1339	0.5933
2016	7 070.81	0.4950	1 512.39	0.1346	0.6296
2017	7 179.50	0.5026	1 472.05	0.1310	0.6336
2018	7 006.23	0.4904	1 483.81	0.1321	0.6225
2019	5 466.96	0.3827	1 306.39	0.1163	0.4990
2020	4 970.97	0.3480	1 110.74	0.0989	0.4468
2021	7 205.65	0.5044	1 524.99	0.1357	0.6401
2022	5 520.99	0.3865	1 385.53	0.1233	0.5098
2023	6 161.68	0.4313	1 409.88	0.1255	0.5568

**Table 4.29: Activity data and N<sub>2</sub>O emissions from integrated iron and steel plant**

YEAR	BLAST FURNACE GAS		CONVERTER GAS		TOTAL
	Quantity flared	N <sub>2</sub> O emission	Quantity flared	N <sub>2</sub> O emission	N <sub>2</sub> O emission
	TJ	t	TJ	t	t
1990	5 385.42	3.0158	1 148.60	0.4594	3.4753
1991	4 783.58	2.6788	1 020.21	0.4081	3.0869
1992	4 464.57	2.5002	952.16	0.3809	2.8810
1993	4 846.94	2.7143	1 033.75	0.4135	3.1278
1994	5 036.11	2.8202	1 074.07	0.4296	3.2498
1995	4 849.83	2.7159	1 034.40	0.4138	3.1297
1996	4 415.13	2.4725	941.71	0.3767	2.8492
1997	4 645.41	2.6014	990.83	0.3963	2.9978
1998	4 687.56	2.6250	999.76	0.3999	3.0249
1999	5 171.60	2.8961	1 102.96	0.4412	3.3373
2000	5 323.22	2.9810	1 135.21	0.4541	3.4351
2001	5 673.85	3.1774	1 209.99	0.4840	3.6614
2002	6 205.19	3.4749	1 323.30	0.5293	4.0042
2003	6 628.37	3.7119	1 413.51	0.5654	4.2773
2004	6 686.20	3.7443	1 425.84	0.5703	4.3146
2005	6 378.37	3.5719	1 366.81	0.5467	4.1186
2006	7 278.92	4.0762	1 559.79	0.6239	4.7001
2007	7 201.14	4.0326	1 543.12	0.6172	4.6499
2008	6 365.24	3.5645	1 364.00	0.5456	4.1101
2009	5 481.64	3.0697	1 174.65	0.4699	3.5396
2010	6 624.68	3.7098	1 419.59	0.5678	4.2777
2011	5 990.75	3.3548	1 277.44	0.5110	3.8658

YEAR	BLAST FURNACE GAS		CONVERTER GAS		TOTAL
	Quantity flared	N <sub>2</sub> O emission	Quantity flared	N <sub>2</sub> O emission	N <sub>2</sub> O emission
	<i>TJ</i>	<i>t</i>	<i>TJ</i>	<i>t</i>	<i>t</i>
2012	6 406.80	3.5878	1 366.19	0.5465	4.1343
2013	6 559.02	3.6730	1 401.04	0.5604	4.2335
2014	7 090.73	3.9708	1 508.14	0.6033	4.5741
2015	5 865.68	3.2848	1 398.03	0.5592	3.8440
2016	6 220.81	3.4837	1 646.25	0.6585	4.1422
2017	7 241.34	4.0552	1 514.47	0.6058	4.6609
2018	8 001.10	4.4806	1 365.17	0.5461	5.0267
2019	4 739.26	2.6540	880.65	0.3523	3.0062
2020	4 433.78	2.4829	979.78	0.3919	2.8748
2021	8 022.07	4.4924	1 753.80	0.7015	5.1939
2022	5 414.26	3.0320	1 288.12	0.5152	3.5472
2023	6 914.44	3.8721	1 481.30	0.5925	4.4646

### Uncertainties and Time-series Consistency

Iron and Steel Production is the significant source of GHG emissions and key category in level and trend assessment, therefore important attention was paid on time series consistency. However, there are several comments to be mentioned:

**Iron and Steel Production in blast furnaces:** Natural gas was also used for heating of blast furnaces since 2000. Therefore, the IEF (CO<sub>2</sub>) decreased from that year. The detailed data for country specific methodology described above are directly available for period 2005 – 2023. The older data (1990 – 2004) has been recalculated in previous submissions by using alternative recalculation techniques (surrogate method). The recalculation was based on the combined driver based on the mass of produced steel and pig iron and the total amount of coking coal used in the plant. Where available, the mass and composition of iron scraps, the mass and composition of iron ore and the mass of pig iron that was not processed to steel were considered to ensure the reliable results. This way of extrapolation provided more consistent data (see comparison of IEF for the boundary years 2003 – 2007). The EU ETS reports are available since 2005, but no disaggregated data on fuel consumption or CO<sub>2</sub> emissions to the very bottom level are presented in these reports. The methodology used by plant operator in the EU ETS report is based on total mass balance and was used for comparison during QA/QC process. The methane and nitrous oxide emissions estimates are reported for the first time in this submission. Detailed activity data are available since 2014. For the period 1990 – 2013, the surrogate method for extrapolation was used. Several drivers were tested. As the best ones, the following drivers were used: (i) driver for the sinter production: the ratio of the sinter produced to the quantity of steel and pig iron produced. The uncertainty of the driver is 2.4%; (ii) driver for the coke production: the ratio of the coke produced to the quantity of steel and pig iron produced. The uncertainty of the driver is 5.0%; (iii) driver for the BFG: the ratio of the BFG flared to the quantity of steel and pig iron produced. The uncertainty of the driver is 1.9%; (iv) driver for the converter gas flaring: the ratio of the converter gas flaring to the quantity of steel produced. The uncertainty of the driver is 6.1%.

**EAF Steel Production:** Emissions estimation is based on the available country specific data and following assumptions

- **Železiarne Podbrezová:** the EU ETS reports are available since 2005. According to the questionnaires sent back by the producer for the period 2000 – 2004, the average value of carbon (in all material inputs) for production is 13.4 kg / 1 t of produced steel.
- **Metalurg Steel:** the EU ETS reports are available since 2005. Until 2006, the CO<sub>2</sub> emission factor was 0.165 t / 1 t of produced steel. This approach was based on the carbon balance made directly by the plant. Since 2007, direct consumption of carbon is available. From the data

---

directly reported in the period 2007 – 2011, carbon consumption was extrapolated using driver method (steel production) back to 1990. The EF (CO<sub>2</sub>) = 0.165 t/t was verified during this exercise. In 2015, the plant did not produce steel. Since 2019, the plant does not produce steel, as well.

- **UNEX Prakovce:** The plant is not included in the EU ETS. The default CO<sub>2</sub> emission factor was used (0.08 t/t) for produced steel. The plant did not produced the steel since 2010.
- **Slovakia Steel Mills:** the EU ETS report with detailed data is available since the start of the production (2013). The production in 2013 was high due to export to the Russian Federation (the RF). In 2014, after economic sanctions put on the RF, the export and subsequently production significantly decreased, too (production from 394 kt in 2013 to 177 kt in 2014 and CO<sub>2</sub> decreased from 40 Gg in 2013 to 15 Gg in 2014). This plant was closed in the end of 2014.

The above-mentioned assumptions were used for the CO<sub>2</sub> emissions estimation in the period 1990 – 1999, as well. Wide range of the EFs for the EAF steel production is based on the content of carbon in the scraps. One of the plants is using low carbon scraps (<0.1% of C). On the other hand, the other plant is using high carbon iron scraps (about 4% of C). Content of carbon in produced steel is approximately 1%. The unequal carbon content results in significantly different EFs.

As described in the **Chapter 4.2**, tier 2 approach to the uncertainty analysis of the subcategories was chosen. Compatible methodology to energy sector was used for uncertainty analyses in this category. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### **4.9.2. Ferrous Alloys Production (CRT 2.C.2)**

Ferrous alloys are produced by the reduction reaction of iron ore and added metal and/or metalloid (Si) oxides in arc furnaces and submerged arc furnaces. Processing CO<sub>2</sub> and CH<sub>4</sub> (only from FeSi alloys) emissions from ferrous alloys production were 7.77 Gg of CO<sub>2</sub> and 4.55 t of CH<sub>4</sub> in 2023. The decrease is by 90% when compared to the year 2022; which is accordance with the decrease of the production (87%). According to the IPCC 2006 GL, also limestone used for the production was included in this estimation.

##### **Methodological Issues**

The CO<sub>2</sub> emissions estimation is based on the carbon material balance (tier 2 approach) described in the IPCC 2006 GL:

$$\text{CO}_2 \text{ emissions} = (\text{C in coal materials} + \text{C in raw materials} + \text{C in carbonates} - \text{C in products}) * 44/12$$

The methane emissions were calculated based on operation specific emission factors (tier 2). The production of FeSi started in 1998. Further information is provided in **Tables 4.30-4.32**.

Plant specific emission factors are estimated annually (based on carbon balance). Methane emission factor is based on the operational specific default value 1.3 kg CH<sub>4</sub>/1 t of FeSi ferrous alloys for whole time series (IPCC 2006 GL). Information on activity data was taken directly from producers of ferrous alloys provided in questionnaires and they are summarized in **Table 4.30**.



**Table 4.30: Activity data used for carbon balance and CO<sub>2</sub> emissions in ferroalloys production in 2023**

Carbon in "Raw Materials"	Carbon in Coals	Limestone Consumed	Carbon in Products	CO <sub>2</sub> Emissions
<i>t</i>				<i>Gg</i>
59.3	2 089.7	NO	29.9	7.770

**Table 4.319: Activity data, CO<sub>2</sub> and CH<sub>4</sub> emissions in ferroalloys production in 1990 – 2001**

YEAR	FERROALLOYS				CaCO <sub>3</sub> Used	Total CO <sub>2</sub>	EF (CO <sub>2</sub> )	Total CH <sub>4</sub>
	Based on Cr	Based on Mn	Based on Si	Total				
	<i>kt</i>							
1990	53.000	116.000	NO	169.000	73.853	296.739	1.756	NO
1991	52.000	113.000	NO	165.000	72.105	289.618	1.755	NO
1992	50.000	110.000	NO	160.000	69.920	281.004	1.756	NO
1993	47.000	103.000	NO	150.000	65.550	263.394	1.756	NO
1994	34.000	111.300	NO	145.300	63.496	259.567	1.786	NO
1995	45.000	89.800	NO	134.800	58.908	235.642	1.748	NO
1996	46.000	84.000	NO	130.000	56.810	226.252	1.740	NO
1997	42.000	78.000	NO	120.000	52.440	209.025	1.742	NO
1998	44.000	81.000	8.666	133.666	58.412	246.984	1.848	11.27
1999	46.700	56.300	13.205	116.205	50.782	220.040	1.894	17.17
2000	17.658	69.458	7.611	94.727	41.396	182.446	1.926	9.89
2001	12.140	69.380	5.200	86.720	37.897	165.901	1.913	6.76

**Table 4.320: Activity data, CO<sub>2</sub> and CH<sub>4</sub> emissions in ferroalloys production in 2002 – 2023**

YEAR	FeSi <sub>75</sub>	FeSi <sub>65</sub>	FeSi <sub>45</sub>	FeSiMn	FeMnC	FeCr	FeSiCa	Total
	<i>kt</i>							
2002	31.208	NO	NO	62.084	56.297	3.521	364	153.474
2003	41.539	NO	NO	52.773	43.434	1.654	1.155	140.555
2004	34.684	NO	NO	64.842	66.959	1.634	1.137	169.256
2005	13.943	1.710	859	47.843	43.458	894	11	108.718
2006	12.319	2.473	1.363	59.128	59.391	NO	NO	134.674
2007	8.417	112	NO	71.587	74.065	NO	NO	154.181
2008	9.510	941	393	59.940	61.194	NO	NO	131.978
2009	4.241	118	278	32.102	20.976	NO	NO	57.715
2010	16.274	9.519	626	34.960	35.449	NO	NO	96.828
2011	22.079	7.174	1.039	25.023	18.180	NO	4.066	77.561
2012	24.658	3.614	201	50.089	12.862	NO	10.168	101.592
2013	30.952	1.761	365	26.794	2.119	NO	3.685	65.676
2014	37.530	1.206	559	29.642	17.554	NO	4.735	91.226
2015	35.761	1.497	929	27.063	25.373	NO	4.898	95.521
2016	27.943	1.799	1.114	35.736	35.589	NO	4.086	106.267
2017	43.117	1.307	210	40.069	42.115	NO	2.661	129.479
2018	39.129	1.543	3.429	37.225	32.364	NO	NO	113.689
2019	27.566	808	1.060	49.897	26.187	NO	NO	105.518
2020	27.679	0.812	1.066	33.812	24.045	NO	1.182	88.596
2021	32.797	0.849	1.145	48.590	30.929	NO	1.478	115.788
2022	10.007	0.250	0.365	15.242	9.106	NO	0.458	35.428
2023	3.504	NO	NO	0.830	0.241	0.079	NO	4.653

YEAR	CaCO <sub>3</sub> Used	Total CO <sub>2</sub>	EF (CO <sub>2</sub> )	Total CH <sub>4</sub>
	<i>kt</i>		<i>t/t</i>	<i>t</i>
2002	67.068	333.657	2.174	40.57
2003	61.423	328.038	2.334	54.00
2004	73.965	371.066	2.192	45.09
2005	47.510	227.646	2.094	20.35
2006	58.853	275.660	2.047	19.23
2007	67.377	301.324	1.954	11.09
2008	57.674	263.043	1.993	13.59
2009	25.221	115.512	2.001	5.67
2010	42.314	219.069	2.262	33.53
2011	33.894	201.979	2.604	38.03
2012	44.396	265.502	2.613	36.75
2013	28.713	165.003	2.512	42.53
2014	41.893	222.894	2.443	50.36
2015	6.428	239.671	2.509	48.43
2016	4.824	237.053	2.482	38.66
2017	4.344	293.887	3.077	57.75
2018	323	281.565	2.948	52.87
2019	NO	239.101	2.503	36.89
2020	NO	214.524	2.421	37.04
2021	NO	251.900	2.843	43.74
2022	NO	75.981	2.145	13.33
2023	NO	7.770	1.670	4.55

### Uncertainties and Time-series Consistency

Carbon balance for CO<sub>2</sub> emissions (and EFs) estimation is used since 2002. Before 2002, a different aggregation of production data is available. EFs in the period 1990 – 2001 are constant and were calculated from available data (1.684 t/t of ferroalloys based on Mn, 1.3 t/t of ferroalloys based on Cr and 3.194 t/t of ferroalloys based on Si). In previous submissions (period 1990 – 2001) verification of emissions calculation was made as follows: (i) the activity data for the period 2002 – 2010 were aggregated in the same way as data available for the years 1900 – 2001; (ii) CO<sub>2</sub> emissions for the period 2002 – 2010 were calculated using the emission factors reported above and compared with the carbon balance method. The difference between these estimations did not exceed 0.6%. Significant increase in emissions since 2002 is caused by the change of the new plant owner’s plans and the new market situation. The using of calcium carbonate in the plant ended during 2018.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.9.3. Aluminium Production (CRT 2.C.3)

Aluminium is produced by the electrolysis of alumina dissolved in cryolite-based melt (t = 950°C). The main additives to cryolite (Na<sub>3</sub>AlF<sub>6</sub>) are aluminium fluoride (AlF<sub>3</sub>) and CaF<sub>2</sub>. In Slovakia, the plants for aluminium production use a modern technology where the majority of HF and other fluorides escaped from the electrolytic cells is absorbed and adsorbed on alumina. Alumina is used subsequently in the

electrolytic process. The anodes are made from graphite. So-called pre-baked anodes for aluminium production are made in separate plants. Due to this technology, emissions are much lower than in the Söderberg process. The release of CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub> emissions can occur at a special technological disturbance (the anode effect). Because of the progress in process control, this irregularity occurs only (1-2) times in a month. Implementation of IPCC 2019 Refinement resulted in the dividing of PFC emission into two sources: (i) high-voltage anode effect (HVAE) that corresponds to the “common” anode effect as described above; (ii) low-voltage anode effect (LVAE) that was not described in IPCC 2006 Guidance, however, it is described in IPCC 2019 Refinement as a new source of PFC emission.

In the middle of 2022, the aluminium plant started to closed due to the high prices of electrical energy. Therefore, the production and emissions decreased very rapidly in this category. At the beginning of January 2023, electrolysis was completely shut down, only the production of anodes remained in operation, which were sold abroad.

### Methodological Issues

Tier 3 in combination with tier 2 approach based on plant specific emission factors and activity data was applied since 2004 in CO<sub>2</sub> and HVAE PFCs emissions estimation. According to the information from producers, 36 t of graphite anodes were used with the sulphur and ash contents 1.59% and 0.20%, respectively, in 2023. The CO<sub>2</sub> emissions from electrolysis were estimated based on the IPCC 2006 GL multiplying the volume of used anodes by carbon content and 44/12 (0.13 Gg CO<sub>2</sub> in 2022). The CO<sub>2</sub> emissions from pitch volatiles combustion and from bake furnace packing material were calculated by the tier 3 method (eq. 4.22 and 4.23 of the IPCC 2006 GL, Volume 3, Chapter 4) and were as follows: 3.13 Gg and 3.62 Gg, respectively. As it was mentioned above, in 2023 electrolysis was shut down, only anode production remained in operation. Before 1996, default EF (CO<sub>2</sub>) = 1.8 t/t for Söderberg process had been used. Since that year, the CO<sub>2</sub> emission factors are evaluated annually in agreement with the tier 3 method described in the IPCC 2006 GL.

The total PFC emissions were 0.002 t (0.013 Gg of CO<sub>2</sub> eq.) in 2023. The HVAE PFC emission was zero. According to the information from the plant operator: *“At the beginning of 2023, only 10 furnaces out of a total of 226 furnaces were in operation in the electrolysis plant, which were shut down after 10 days. PFC emissions from primary aluminum production occur due to the anode effect. The anode effect is an undesirable condition and occurs when an unstable state occurs in an electrolysis furnace, which is caused by various chemical-physical phenomena. During these 10 days, when the instruction to completely shut them down was essentially awaited, these furnaces were in a very stabilized state and no anode effect occurred. This data is also recorded in the Elpos control system at the operator Slovalco”.*

The LVAE PFC emissions was calculated using Tier 1 method (with EF(CF<sub>4</sub>) = 0.009 kg / t of aluminium). There is not methodology for tier 2, and tier 3 requires specific measurements that were not realized in the plant. SF<sub>6</sub> is not used in aluminium castings in the Slovak Republic.

**Table 4.33: CO<sub>2</sub> emissions and EFs in aluminium production in particular years**

YEAR	Aluminium Production	CO <sub>2</sub> (Electrolysis)	CO <sub>2</sub> (Anode Production)	Total CO <sub>2</sub>	EF per Aluminium
	kt			Gg	t/t
1990	67.40	121.32	NE	121.32	1.8000
1995	32.60	58.68	NE	58.68	1.8000
2000	109.81	160.33	16.23	176.56	1.6078
2005	159.20	230.69	23.53	254.22	1.5968
2010	163.00	239.38	24.09	263.47	1.6164
2011	162.84	237.21	24.07	261.28	1.6045
2012	160.66	235.77	23.75	259.52	1.6153
2013	163.30	241.10	24.14	265.24	1.6243

YEAR	Aluminium Production	CO <sub>2</sub> (Electrolysis)	CO <sub>2</sub> (Anode Production)	Total CO <sub>2</sub>	EF per Aluminium
	kt			Gg	t/t
2014	167.67	246.07	19.93	266.00	1.5865
2015	171.33	253.74	22.59	276.33	1.6129
2016	173.64	257.08	14.34	271.41	1.5631
2017	173.49	257.97	16.04	274.01	1.5794
2018	173.72	256.20	19.33	275.53	1.5860
2019	174.79	256.20	18.51	274.71	1.5716
2020	151.87	223.24	15.47	238.71	1.5717
2021	164.00	241.54	17.14	258.68	1.5773
2022	71.93	103.10	8.35	111.45	1.5493
2023	0.21	0.13	6.75	6.88	32.2683

**Table 4.34: PFC emissions and EFs in aluminium production in particular years**

YEAR	HVAE CF <sub>4</sub>	EF per Aluminium	HVAE C <sub>2</sub> F <sub>6</sub>	EF per Aluminium	LVAE CF <sub>4</sub>	Total PFC
	t	kg/t	t	kg/t	t	Gg CO <sub>2</sub> eq.
1990	28.15	0.4176	2.42	0.0359	0.07	28.15
1995	11.86	0.3637	1.02	0.0313	0.03	90.15
2000	1.30	0.0118	0.13	0.0011	0.99	16.56
2005	2.28	0.0143	0.22	0.0014	1.43	27.09
2010	2.41	0.0148	0.23	0.0014	1.47	28.27
2011	1.93	0.0119	0.19	0.0012	1.47	24.63
2012	2.47	0.0154	0.24	0.0015	1.45	28.62
2013	0.94	0.0058	0.09	0.0006	1.47	17.02
2014	1.07	0.0064	0.10	0.0006	1.51	18.27
2015	0.82	0.0048	0.08	0.0005	1.54	16.53
2016	0.62	0.0036	0.06	0.0003	1.56	15.17
2017	0.83	0.0048	0.08	0.0005	1.56	16.75
2018	0.75	0.0043	0.07	0.0004	1.56	16.14
2019	0.50	0.0029	0.05	0.0003	1.57	14.28
2020	0.54	0.0036	0.05	0.0003	1.37	13.22
2021	0.57	0.0035	0.06	0.0003	1.48	14.19
2022	0.21	0.0029	0.02	0.0003	0.65	5.88
2023	NO	NA	NO	NA	0.00	0.01

### Uncertainties and Time-series Consistency

The technology was changed from Søderberg to prebaked technology in 1996. It results in significant decrease of CO<sub>2</sub> and PFC emissions. The CO<sub>2</sub> emissions were calculated by using the tier 1 method in the period 1990 – 1995 due to lack of detailed data. Due to the changes in ownership of the plant (and new producing policy), higher tier method can be implemented since 1996. Input data necessary for tier 3 method are available since 2005. Average CO<sub>2</sub> emission factor calculated from years 2005 and 2010 – 2012 was used also for years 1996 – 2004 (emission factors based on the years 2006 – 2009 could not be used due to technological reasons. Background data about it were made available and accepted by the ERT during in-country review in 2012). According to the questionnaire sent by producer, the significant progress in control of the electrolysis was achieved in 2009 (this information is confidential but was provided together with the reasoning of the IEF (CO<sub>2</sub>) decrease during the in-country review in 2012). The improvements in production resulted also in decrease of PFC emissions after 2009. Further improvement in better performance controlling process of electrolysis cells continues until now. The CO<sub>2</sub> emissions from pitch volatiles combustion and from bake furnace packing material

were calculated in 2013 for the first time (according to the IPCC 2006 GL) and the resulting implied emission factor per produced aluminium was estimated. This IEF was also used for the time series 1996 – 2012. This IEF is almost without change also for next years and recalculation of the time series 1996 – 2012 is not necessary.

In the 2024 submission, the new source of PFC emission was adopted from IPCC 2019 Refinement. The emissions reported so far represented emissions from HVAE as it is now defined in IPCC 2019 Refinement. They were recalculated using new default coefficients of the Slope method. New source of PFC emission from LVAE was calculated using the default emission factors for the corresponding technologies used.

As described in the **Chapter 4.2**, tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.9.4. Magnesium Production (CRT 2.C.4)

This production does not occur in the Slovak Republic, therefore the notation key “NO” for time series was used.

#### 4.9.5. Lead Production (CRT 2.C.5)

Lead is produced only from secondary raw materials in Slovakia. This production started in Imperial Smelt Furnaces in 2011 and is not significant. The CO<sub>2</sub> emission was 84.80 t in 2023.

##### Methodological Issues

This category is not key category and therefore tier 1 method based on the IPCC 2006 GL was used for whole time series. Default EF (0.2 t/t) for CO<sub>2</sub> emissions from treatment of secondary raw materials was used for whole time series. According to the direct information from the plant operator, 424 t of lead was produced from the secondary raw materials in 2023.

**Table 4.35:** The overview of activity data and CO<sub>2</sub> emissions from lead production in 1990 – 2023

YEAR	Lead Production from Secondary Materials	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	t	t	t/t
1990-2010	NO	NO	NA
2011	49.81	9.96	0.2
2012	203.63	40.73	0.2
2013	261.10	52.22	0.2
2014	292.70	58.54	0.2
2015	323.12	64.62	0.2
2016	292.05	58.41	0.2
2017	303.83	60.77	0.2
2018	47.60	9.52	0.2
2019	66.00	13.20	0.2
2020	125.00	25.00	0.2
2021	155.00	31.00	0.2
2022	400.00	80.00	0.2
2023	424.00	84.80	0.2

## Uncertainties and Time-series Consistency

Tier 1 method according to the IPCC 2006 GL is used in whole time series.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.9.6. Zinc Production (CRT 2.C.6)

Zinc is produced by pyrometallurgical process involving the use of an Imperial Smelting Furnace, which allows the simultaneous treatment of lead and zinc concentrates in Slovakia and is not significant. This production started in 2012. Since 2015, the production was not occurring.

#### Methodological Issues

This category is not key category and therefore tier 1 method based on the IPCC 2006 GL was used for whole time series.

Default EF (0.43 t/t) for CO<sub>2</sub> emissions from pyrometallurgical process was used for whole time series. According to the direct information from the plant operator, no zinc was produced in 2023.

**Table 4.36:** The overview of activity data and CO<sub>2</sub> emissions from zinc production in 1990 – 2023

YEAR	Zinc Production (Pyrometallurgical - ISF)	CO <sub>2</sub> Emissions	IEF (CO <sub>2</sub> )
	t		t/t
1990 – 2011	NO	NO	NA
2012	43.90	18.88	0.43
2013	31.45	13.52	0.43
2014	23.94	10.29	0.43
2015 – 2023	NO	NO	NA

## 4.10. Non-energy Products from Fuels and Solvent Use (CRT 2.D)

This category produces emissions of CO<sub>2</sub> and NMVOC. Based on known composition of NMVOC emissions, indirect (potential) CO<sub>2</sub> emissions were calculated in this submission, too. Direct CO<sub>2</sub> emissions were 60.21 Gg in 2023 and increased by approximately 24% compared with the previous year. When comparing with the base year, the increase was 19% caused by the use the SCR in cars.


**Table 4.37:** Emissions in the category 2.D according to subcategories in particular years

YEAR	2.D.1 Lubricant Use	2.D.2 Paraffin Wax Use	2.D.3 Other
	Gg of CO <sub>2</sub> eq.		
1990	48.02	2.46	NO
1995	48.02	2.46	NO
2000	48.02	2.46	NO
2005	28.94	1.23	NO
2010	12.39	2.54	2.01
2011	18.52	1.90	3.48
2012	27.11	2.52	3.93
2013	32.51	2.54	6.09

YEAR	2.D.1 Lubricant Use	2.D.2 Paraffin Wax Use	2.D.3 Other
	Gg of CO <sub>2</sub> eq.		
2014	26.58	3.17	6.50
2015	26.85	2.54	6.32
2016	26.40	2.54	8.77
2017	28.46	2.52	10.67
2018	28.88	1.90	10.57
2019	23.61	2.54	16.70
2020	19.05	2.54	18.95
2021	21.58	2.54	30.01
2022	28.15	2.54	17.98
2023	30.54	2.54	27.13

The major share (50.7%) in emissions belongs to the lubricant use category, 45.1% belongs to the other used (urea use) and 4.2% to the paraffin wax use.

**Figure 4.10:** The share of GHG emissions in individual subcategories of the 2.D in 2023



In 2021 submission, recalculations were focused on the NMVOC emissions from solvent use have been prepared since the base year 1990. Also, harmonization between the GHG a CLRTAP inventories continuing and the completion of the QA/QC process of NMVOC emissions in 2.D.3 categories was finished in 2020 and presented in 2021 submission. The results are summarised in the [Annex 4.4](#) of NID 2022. Moreover, CO<sub>2</sub> emissions resulted from the NMVOC emissions are indirect and are reported according to the document [“Conclusions and recommendations from the 17<sup>th</sup> meeting of greenhouse gas inventory lead reviewers”](#). No recalculation was made in this submission regarding the indirect CO<sub>2</sub> emissions.

#### 4.10.1 Lubricant Use (CRT 2.D.1)

Lubricants are mostly used in industry and transport. The CO<sub>2</sub> emissions estimated in Slovakia from this category were 30.54 Gg in 2023.

##### Methodological Issues

This category is not key category and therefore tier 1 method based on the IPCC 2006 GL was used for whole time series.

Default carbon content (20 t CO<sub>2</sub>/TJ) and ODU (Oxidized During Use) factor (0.2) according to the IPCC 2006 GL was used.

Activity data of non-energy use of lubricants are available from the Statistical Office of the Slovak Republic. Total volume of lubricants for non-energy use in Slovakia was 2 084.1 TJ in 2023. Due to technical reasons, the activity data in this category are presented in CRT Tables in kilotons units. Due to lack of relevant statistics, data for the time series 1990 – 2001 were approximated by the Statistical Office of the Slovak Republic.

**Table 4.38:** *The overview of activity data and CO<sub>2</sub> emissions in lubricant non-energy use in particular years*

YEAR	Lubricant Use	Lubricants Use	CO <sub>2</sub> Emissions
	kt	TJ	Gg
1990	78	3 276.8	48.024
1995	78	3 276.8	48.024
2000	78	3 276.8	48.024
2005	47	1 974.5	28.938
2010	20	845.2	12.388
2011	30	1 263.5	18.517
2012	44	1 849.5	27.106
2013	53	2 218.4	32.513
2014	44	1 813.4	26.577
2015	45	1 831.8	26.847
2016	46	1 801.5	26.402
2017	47	1 941.5	28.455
2018	47	1 970.4	28.878
2019	39	1 611.1	23.612
2020	31	1 300.0	19.053
2021	35	1 472.5	21.581
2022	46	1 920.7	28.150
2023	50	2 084.1	30.544

#### Uncertainties and Time-series Consistency

Tier 1 approach according to the IPCC 2006 GL is used in whole time series. As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.10.2. Paraffin Wax Use (CRT 2.D.2)

Paraffin waxes are mostly derived by combustion of waxes derivate of paraffin (e.g. candles). The CO<sub>2</sub> emissions estimated in Slovakia from this category were 2.54 Gg in 2023.

#### Methodological Issues

This category is not key category and therefore tier 1 method based on the IPCC 2006 GL was used for whole time series. Default carbon content (20 t CO<sub>2</sub>/TJ) and ODU factor (0.2) according to the IPCC 2006 GL was used. Activity data on non-energy use of paraffin wax in Slovakia are available from the Statistical Office of the Slovak Republic. Total volume of paraffin wax for non-energy use in Slovakia was 173.2 TJ (4 kt) in 2023. No paraffin wax was reported in the years 2004 and 2006 (based on the



statistical data). Due to lack of relevant statistics, data for the time series 1990 – 2002 were approximated by the Statistical Office of the Slovak Republic.

**Table 4.39: The overview of activity data and CO<sub>2</sub> emissions in paraffin wax non-energy use in particular years**

YEAR	Paraffin Wax Use	Paraffin Wax Use	CO <sub>2</sub> Emissions
	<i>kt</i>	<i>TJ</i>	<i>Gg</i>
1990	4	168.04	2.46
1995	4	168.04	2.46
2000	4	168.04	2.46
2005	2	84.02	1.23
2010	4	173.20	2.54
2011	3	129.90	1.90
2012	4	172.00	2.52
2013	4	173.20	2.54
2014	5	216.50	3.17
2015	4	173.20	2.54
2016	4	173.20	2.54
2017	4	172.00	2.52
2018	3	129.90	1.90
2019	4	173.20	2.54
2020	4	173.20	2.54
2021	4	173.20	2.54
2022	4	173.20	2.54
2023	4	173.20	2.54

### Uncertainties and time-series consistency

Tier 1 method according to the IPCC 2006 GL is used in whole time series.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.10.3. Other (CRT 2.D.3)

This category includes potential CO<sub>2</sub> and NMVOC emissions from solvent use, road paving with asphalt. CO<sub>2</sub> emissions were calculated from solvent use, where composition of NMVOC emissions is known. It should be noted that CO<sub>2</sub> emissions represent only potential emissions originate from the oxidation of NMVOC emissions. Total NMVOC emissions from solvent use, road paving with asphalt and asphalt roofing were estimated in the frame of the National Program for Emission Reduction of Non-Methane Volatile Organic Compounds in the Slovak Republic.

Collection of available input data of solvents used in industry was the most challenging step in inventory preparation. Official statistical information in this area was insufficient, so it was decided directly requested producers, importers, distributors and users of solvents and other products. This inventory was prepared in the consistency with the CLRTAP inventory. During last several submissions, the CLRTAP inventory was recalculated in several 2.D.3 subcategories. The results of the recalculation were always adopted in GHG inventory, which resulted in the recalculation of NMVOC and CO<sub>2</sub>

emissions in 2.D.3 category since the base year. In 2020, the thorough QA/QC process focused on the harmonisation of the CLRTAP (NECD) and the GHG inventories for the 2.D.3 categories was finished, and the recalculation was necessary in previous submission. More information about the comparison of changes among submissions and detailed activity data can be found in the [Annex 4.4](#) of NID 2022 and detailed information is presented in the CLRTAP submission 2022. The respective indirect CO<sub>2</sub> were calculated on the basis of stoichiometry of NMVOC emissions.

Urea used in catalytic converters is reported in this category. The use of urea in catalytic converters can occur in vehicles and in industrial plants. The CO<sub>2</sub> emissions from urea based catalysts were estimated using COPERT 5 model. The fuel consumption of diesel oil corresponding heavy duty trucks and passenger cars with SCR are included in the category 1.A.3.b.

The use of urea in industrial plants is reported in Slovakia since 2016. This possible use of urea is annually monitored by questionnaires that are sent to the operators. The only NO<sub>x</sub> reduction method used in Slovakia before 2016 occurred in the ammonia plant where ammonia is used for reduction purposes and no CO<sub>2</sub> emissions occur at this method. Since 2016, due the new emission limits for NO<sub>x</sub>, seven plants started using the DeNOx technologies. Three of them are using the ammonia, the rest are using the urea.

Total direct GHG emissions in this category were 27.13 Gg of CO<sub>2</sub> eq. in 2023. Total NMVOC emissions were 18.58 kt. [Table 4.40](#) summarizes CO<sub>2</sub> and NMVOC emissions for particular years of time series.

**Table 4.40: CO<sub>2</sub> and NMVOC emissions (Gg) in 2.D.3 in particular years**

YEAR	NMVOC Emissions	Indirect CO <sub>2</sub> Emissions	Direct CO <sub>2</sub> Emissions
	kt	Gg	
1990	38.495	87.769	NO
1995	35.824	82.085	NO
2000	29.601	65.444	NO
2005	30.732	66.929	NO
2010	22.416	49.201	2.012
2011	26.146	57.615	3.484
2012	21.197	46.484	3.925
2013	21.088	46.413	6.090
2014	22.502	49.541	6.504
2015	25.643	56.344	6.315
2016	23.925	52.517	8.767
2017	21.725	47.481	10.669
2018	24.154	53.114	10.571
2019	20.547	45.301	16.696
2020	20.851	45.875	18.951
2021	19.886	43.666	30.009
2022	17.890	39.505	17.979
2023	18.577	41.079	27.130

### Methodological Issues

In the CLRTAP inventory, 2.D.3 category consists of following subcategories:

- 2.D.3.a Domestic solvent use including fungicides
- 2.D.3.b Road paving with asphalt
- 2.D.3.c Asphalt roofing
- 2.D.3.d Coating application
- 2.D.3.e Degreasing

- 2.D.3.f Dry cleaning
- 2.D.3.g Chemical products
- 2.D.3.h Printing
- 2.D.3.i Other solvent use

In the GHG inventory, all categories except of 2.D.3.b and 2.D.3.c are reported under 2.D.3 Other – Solvent Use. Categories 2.D.3.b and 2.D.3.c are reported separately in 2.D.3 Other – Road paving with asphalt and 2.D.3 Other – Asphalt roofing.

During the QA/QC process performed in last years, a great effort was made to identify the chemical compounds in NMVOC emissions. 97 chemical compounds were identified. Due to this large number, the list of the chemical compounds is not presented in the Document, however, it is available to the ERT. Carbon content in the chemical compounds was calculated based on the stoichiometry of the molecule. For the others NMVOC emissions the carbon content was assumed to be the default value (0.6). The identification of large number of chemical compounds in the NMVOC emissions, made the CO<sub>2</sub> emissions estimate more accurate than in the previous submissions where only several groups of the chemicals were reported. CO<sub>2</sub> emissions were calculated for each subcategory separately (2.D.3.a – 2.D.3.i) since the year 2000. Extrapolation by the trend was used for the years 1990 – 1999 for each category, as well. The results are presented in [Tables 4.41-4.42](#). Detailed data are presented in the [Annex 4.4](#) of NID 2022.

The CO<sub>2</sub> emissions from urea based catalysts from cars were estimated using COPERT 5 model for vehicle category “Heavy duty trucks Euro V 2008 Standards” and “Passenger cars Diesel PC Euro 6 up to 2016” for the years 2010 – 2023. As the number of vehicles with SCR technology is not known, the default value in COPERT model 5 was used. The urea based catalysts were not used before 2010. More information is included in the Chapter 3 of this Document. The CO<sub>2</sub> emissions from urea based catalysts in industry were calculated from the amount of used urea in industrial DeNOx technologies. Activity data on the urea use were reported in the CRT Software as the sum of the urea used for industrial DeNOx technologies and of its use in vehicles. However, the concentration of urea solution in cars is assumed to be 32% in COPERT 5 model, while the concentration of the urea in industrial DeNOx technologies is usually 40%. Therefore, the consumption of urea from use in vehicles was estimated by reverse calculation from the CO<sub>2</sub> emissions in the term of the pure urea and summed with the pure urea (calculated from the 40% solution) from industrial technologies. In the NID, the activity data on the urea use are reported separately according to the ERT recommendation I.5 based on the ARR 2022. The CO<sub>2</sub> emissions from urea based catalysts are presented in [Table 4.43](#).

**Table 4.41: NMVOC and CO<sub>2</sub> emissions in solvent use category in particular years**

YEAR	NMVOC Emissions	Indirect CO <sub>2</sub> Emissions
	kt	Gg
1990	38.386	87.512
1995	35.771	81.961
2000	29.575	65.382
2005	30.708	66.874
2010	22.399	49.164
2011	26.125	57.570
2012	21.179	46.446
2013	21.070	46.373
2014	22.486	49.504
2015	25.622	56.297
2016	23.904	52.471
2017	21.705	47.436
2018	24.132	53.066

YEAR	NMVOE Emissions	Indirect CO <sub>2</sub> Emissions
	kt	Gg
2019	20.529	45.261
2020	20.834	45.837
2021	19.863	43.615
2022	17.867	39.454
2023	18.557	41.034

**Table 4.42: NMVOC and CO<sub>2</sub> emissions from asphalt using in particular years**

YEAR	Road Paving with Asphalt	Asphalt Roofing	Road Paving with Asphalt	Asphalt Roofing	Road Paving with Asphalt	Asphalt Roofing
	Asphalt use in kt		NMVOC in t		Indirect CO <sub>2</sub> in t	
	1990	366.8	130.2	62.355	46.717	154.994
1995	171.0	65.9	29.067	23.659	72.251	52.051
2000	52.5	46.5	10.363	16.323	25.760	35.910
2005	113.0	32.3	19.138	5.773	42.103	12.701
2010	102.4	25.3	14.373	2.402	31.620	5.285
2011	121.0	28.1	18.230	2.411	40.105	5.304
2012	102.3	27.6	14.870	2.340	32.715	5.147
2013	86.0	41.0	15.197	2.907	33.434	6.396
2014	79.2	59.4	13.746	2.635	30.242	5.797
2015	147.3	37.9	20.067	0.973	44.147	2.141
2016	150.8	66.4	18.942	1.959	41.672	4.310
2017	115.0	50.6	18.737	1.290	41.221	2.838
2018	146.0	68.5	19.933	2.096	43.852	4.611
2019	132.5	63.7	16.455	1.900	36.201	4.180
2020	132.9	65.0	15.082	2.186	33.180	4.810
2021	154.8	79.0	20.829	2.229	45.823	4.904
2022	163.1	59.4	20.953	2.114	46.096	4.652
2023	152.5	53.3	18.040	2.398	39.689	5.275

**Table 4.43: CO<sub>2</sub> emissions originating from the use of urea in catalytic converters in 2010 – 2023**

YEAR	Urea Consumption in Industry	CO <sub>2</sub> Emissions in Industry	Urea Consumption in Cars	CO <sub>2</sub> Emissions in Cars	Total CO <sub>2</sub> Emissions
	t				
2010	NO	NO	2 745.8	2 012.2	2 012.2
2011	NO	NO	4 753.6	3 483.5	3 483.5
2012	NO	NO	5 356.5	3 925.3	3 925.3
2013	NO	NO	8 310.2	6 089.8	6 089.8
2014	NO	NO	8 874.8	6 503.6	6 503.6
2015	NO	NO	8 617.6	6 315.1	6 315.1
2016	2 227.8	1 632.6	9 735.0	7 134.0	8 767.0
2017	2 271.0	1 664.2	12 287.8	9 004.7	10 668.9
2018	1 997.8	1 464.0	12 428.0	9 107.4	10 571.4
2019	732.4	536.7	21 916.5	16 060.7	16 696.4
2020	1 568.8	1 149.6	24 291.4	17 801.1	18 950.7
2021	1 661.1	1 217.3	39 289.1	28 791.7	30 009.0
2022	1 623.0	1 189.4	22 911.5	16 789.9	17 979.3
2023	1 364.2	999.7	35 657.0	26 130.0	27 129.7

## Uncertainties and Time-series Consistency

Consistent methodology and tier method was used for the whole time series in all subcategories mentioned in this chapter. The detailed data are available since 2000. The extrapolation was used for the rest of the time series. The extrapolation was based on the average IEF of CO<sub>2</sub> per 1 t of NMVOC from the years 2000 – 2005.

As described in the **Chapter 4.2**, tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

## 4.11. Electronic Industry (CRT 2.E)

No halocarbons, SF<sub>6</sub> or NF<sub>3</sub> were used in the Slovak Republic in 1990 – 2023 in this category, therefore notation key “NO” was used in all 2.E categories.

## 4.12. Product Uses as Substitutes for ODS (CRT 2.F)

### 4.12.1. Source Category Description

F-gases notion means the emissions of substances that, because of their effects, can be added to the greenhouse gases group. However, before COP3 in Kyoto F-gases were not considered in the GHG emissions inventory or GHG emission projections.

At the present, following gases are included into inventory submission of the Slovak Republic:

- HFCs – hydrofluorocarbons (23, 32, 125, 134a, 152a, 143a, 227ea, 236fa, 245fa, 365mfc);
- SF<sub>6</sub> – sulphur hexafluoride;
- PFCs – per fluorocarbons (CF<sub>4</sub> for the period 1997 – 2005; C<sub>2</sub>F<sub>6</sub>).


The PFC emissions (CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub>) from metal production are reported in 2.C.3 – Aluminium Production. The inventory of F-gases is complicated due to a high number of substances. These gases are components of different mixtures and are used in more than 15 different applications. Each application has its own development of consumption and emissions trend. To ensure environmental integrity, the post-2012 agreement includes additional fluorinated gases (NF<sub>3</sub>, hydrofluoroethers and perfluoropolyethers) with lower GWPs. There are two additional HFCs gases already reported in the Slovak inventory under memo items: HFC 245fa and HFC 365mfc. These gases are used in industry as foam agent (polyurethane-foam blowing agent – PU closed cell foam and integral PU-foam) with the highest consumption as PU spray foam for roof insulation.

**Table 4.44:** The overview of actual HFCs and PFCs emissions in particular years

YEAR	2.F.1	2.F.2	2.F.3	2.F.4	2.F.5	2.F.6	Total 2.F
	<i>Gg CO<sub>2</sub> eq.</i>						
1990	NO	NO	NO	NO	NO	NO	NO
1995	10.202	NO	2.179	NO	NO	NO	12.381
2000	82.192	5.597	7.988	2.425	1.274	NO	99.475
2005	252.913	4.803	13.196	6.182	0.398	NO	277.491
2010	543.084	2.114	16.908	7.116	NO	NO	569.222

YEAR	2.F.1	2.F.2	2.F.3	2.F.4	2.F.5	2.F.6	Total 2.F
	Gg CO <sub>2</sub> eq.						
2011	548.226	2.182	18.403	7.624	NO	NO	576.434
2012	574.144	2.542	17.684	7.704	NO	NO	602.074
2013	593.363	2.142	17.391	8.098	NO	NO	620.993
2014	598.021	1.985	17.728	8.406	NO	NO	626.139
2015	674.550	1.800	19.427	9.058	NO	NO	704.835
2016	615.631	1.790	21.279	9.253	NO	NO	647.952
2017	679.098	1.781	20.991	8.325	NO	NO	710.194
2018	645.605	1.772	20.141	8.103	NO	NO	675.621
2019	656.458	1.763	22.155	8.310	NO	NO	688.686
2020	615.045	1.754	21.546	8.303	NO	NO	646.649
2021	642.666	1.745	19.130	8.832	NO	NO	672.373
2022	447.340	1.737	21.931	9.852	NO	NO	480.860
2023	398.920	1.728	27.204	10.038	NO	NO	437.890


Figure 4.11: The share of emissions in the 2.F category according to the subcategories in 2023



Total actual HFCs and PFCs emissions reported in the category 2.F Product uses as substitutes for ODS were 437.89 Gg of CO<sub>2</sub> eq. in 2023 and they decreased by 9% compared to the previous year. The decrease corresponds to the decrease of emissions in 2.F.1 category. The reasons for such high decrease are discussed in the respective categories.

The decrease was expected due the Regulation (EU) No 517/2014 of the European parliament and of the Council on fluorinated greenhouse gases. However, due to the decommissioning of the equipment with the high GWP gases, only small decrease occurred in several last years. Total trend in several last years (since 2017) fluctuated but was slowly decreasing. Since 2022 the implementation of the Regulation (EU) No 517/2014 is fully manifested.

**Figure 4.12: Trend in individual F-gases in 1990 – 2023**



Generally, increasing trend is visible since the base year and is caused by supplying HCFCs gases by the HFCs up to 2010. However, the emissions of F-gases were approximately constant since 2010 because of the almost complete replacement of HCFCs gases. Another reason of the change in trend is the use of HFC-134a in mobile air conditioners (ACs). Coolant R134a showed continuing increasing trend mainly because of rising uses of cars with ACs. This trend stopped in 2010. It was caused by smaller purchases of cars in Slovakia since 2010, which resulted in a smaller bank of HFC-134a in Slovakia. Also HFO-1234yf is used in an increased extent in new cars; while in 2016 the ratio of HFO-1234yf and HFC-134a was ca 1:1, in 2023 the ratio is ca 9:1.

#### 4.12.2. Activity Data

Before year 2009, the activity data had been collected via paper questionnaires addressed to the 250 potential suppliers, users and consumers of the substances based on the description of the substances with GWP (global warming potential). These potential consumers of the substances were requested annually by the letter authorized by the Ministry of Environment. Provided data enabled to determine the rate of emissions and new filling by using the method of approximation. In case of any uncertainty, received data were verified by the provider and they were summarized in tables according to the way of use. Since the year 2009, input data are reported through the new electronic system that includes also a servicing of the installed equipment (more than 1 200 operators report data). Tables used since 1990 were used also in the latest inventory years for data storage and archiving in order to retain the continuity of observing trends.

The implemented electronic system consists of:

- Annual reporting of F-gases (new charges and leakages) by certified companies;
- Annual reporting of F-gases imported in bulks by certified companies;
- Annual reporting of F-gases in products by importers, exporters, producers by companies.

All operators dealing with the F-gases have access to this electronic system based on certification. Advantages of electronic data logging and reporting are in the possibilities of automatic analysis, fault detection and comparison of, fast access to the full history of leak checks and various forms of output. Service engineers get quick survey of the customers, cooling circuits, details of all maintenance work and repairs, refrigerants in store, refrigerants added, recovered, reclaimed, and disposed of it. Value added of electronic logbook is indirect detection of refrigerant leak. The fault detection classifier estimates the probability of refrigerant leak. Electronic way of the data records enables summarizing, reporting and analysing important data in a chosen period.

---

This system is based on the activities of the Slovak Association for Cooling and Air-conditioning Technology (SZ CHKT) and started its operation in the year 2009. The Slovak Association for Cooling and Air-conditioning Technology (the “Notified Body”) is the body officially authorized by the Ministry of Environment to certify companies and organizations for the activities in this area. The electronically led documentation has developed from the previous paper form. Evaluated data were collected from the service organizations. Details on the electronic system and method of data collection are presented in the Annex 4.2 of this Document.

The Slovak Republic reports emissions of HFCs and PFCs gases (use of substances) in the IPPU sector in the following subcategories:

- 2.F.1 Refrigeration and air conditioning
- 2.F.2 Foam blowing agents
- 2.F.3 Fire protection
- 2.F.4 Aerosols

In the following subcategories are emissions not reported in 2019 and the notation key “NO” was used:

- 2.F.5 Solvents – no gases occur in this category since 2006;
- 2.F.6 Other application – no gases occur in this category.

### **4.12.3. Emission Factors**

Emission factors were evaluated in each category for each individual product (or using of gas) to ensure the best available accuracy of inventory. EFs are described in each category.

### **4.12.4. Methods**

The actual emission estimation of time series was performed mainly by tier 2 method that accounts for the time lag between the consumption and the emissions. Detailed description of methodology is provided in the subcategories.

### **4.12.5. Uncertainties and Time-series Consistency**

A consistent time series of the HFCs import-export exists since 1995 and is well documented, based on questionnaires. In previous submissions, the data on banks were not consistent because of use of extrapolation in 2010. The recalculation of banks was made according to the discussion with the ERT during six weeks’ period after in-country review in 2012. In that submission, this extrapolation was useful in order to differentiate the banks among different subcategories. The differentiation was based on the 2010 share of HFCs gases. Data gathered from 2010 onwards on refrigerant use by subcategories support this disaggregation. Therefore, the disaggregation was accepted by the ERT as a final one and further step, removal of inconsistency of bank data, followed. The inconsistency was caused by the two types of formula used for bank calculation. In the 2015 submission, the bank data were recalculated by the same formula (previously used only for data since 2010) for the whole time series. The reported emissions are also influenced by the recalculation of the disposal emissions in last reporting years. A new, consistent method for the estimation of retiring equipment was used in 2015 submission. The main change in 2016 submission was the recalculation of reported recovery (in ETF Software). In previous submissions, the recovery represented the amount of HFCs and PFCs that was recovered and recycled from the disposed systems and could be used again. Since 2016 submission, the recovery represents amount that was recovered, recycled, and destroyed from disposed systems. Emissions were not influenced by this correction. In 2017 submission, the recalculation of operational emissions has been done in 2.F.1 category. This recalculation considered the possibility of no servicing of equipment few years before its decommissioning. Details are presented in 2.F.1 category.

Monte Carlo method for the uncertainty analyses was implemented for F-gases for the first time in 2016 submission. The IPCC default values for uncertainty of activity data and emission factors were used. The results of the simulation are presented in respective subcategories.



---

#### 4.12.6. Source-Specific QA/QC and Verification

Slovakia has a unique reporting system of F-gases in bulks and in products. Due to the reporting system includes all F-gases, the QA/QC of 2.G category is included here, as well. Data processing system and verification is done automatically. The advantages of the system are as follows:

- historical development of reported data in numbers and graphs during the reported years are available,
- reported numbers from importers (wholesalers) of F-gases and reported numbers from service companies are compared,
- Notified Body has access to historical development in all monitored categories and compares it with ex-post and ex-ante projections up to 2030.

This data processing system allows calculating the emissions by top-down approach. However, the differentiation of the reported data into subcategories is rather limited (mainly in 2.F.1 - Refrigeration and air conditioning).

The new internet reporting system Leaklog has been running since 2009 on the legal basis (Act No 286/2009 Coll. and its amendment No 314/2009 Coll.). Increased publicity from the Ministry of Environment and increased number of inspections from the Slovak Inspection of Environment has increased knowledge of the companies to get data that are more accurate. This system allows estimating the emissions by the bottom-up approach with differentiating among subcategories (see the [Annex 4.2](#) for more details). These two sets of data are supplementary to each other and allow comparing the total amounts of F-gases consumed in Slovakia. Data from these two reporting systems are compared annually.

[Refrigeration and Air Conditioning](#) - Verification is a part of electronic database system.

[Fire Extinguishers](#) - The information on fillings and recycling of already used fire extinguishers is realized with the cooperation of the [Association of the Fire Extinguishers Producers](#) in the Slovak Republic based on the Regulation of the Ministry of Environment of the Slovak Republic No 314/2009 Coll. The Association is obliged to provide information from all members. The sector-specific QA/QC activities were performed as described in the [Chapter 4.2](#) of the Document and results are verified by the top-down approach. Verification is a part of electronic database system.

[N<sub>2</sub>O from Product Uses](#) - Due to the lack of appropriate statistical information and methodological advises in this category, inputs were taken directly from the questionnaires sent to distributors of N<sub>2</sub>O liquid gas in the Slovak Republic.

#### 4.12.7. Source-Specific Recalculations

No recalculation was made in this submission.

#### 4.12.8. Source-Specific Planned Improvements

No improvements are planned.

#### 4.12.9. Refrigeration and Air Conditioning Equipment (CRT 2.F.1)

The emissions originating from refrigeration and AC equipment represent more than 90% of emissions from the 2.F category. Therefore, these emissions are significant source. Total actual emissions of HFCs were 398.92 Gg of CO<sub>2</sub> eq. in 2023 and they decreased by 11% in comparison with the previous year. The decrease due the Regulation (EU) No 517/2014 of the European parliament and of the Council on fluorinated greenhouse gases was expected. The expectations are caused by several reasons. One of them is end of using of new R404A gas with GWP 3940 (only recovered gas can be used since now). Another reason is using of new replacements of R404A, R410A, R134a with low GWP blends (in Slovakia new gases R448A, R449A, R454B, R454C, R513A, R514A, R1234yf, R1234ze were


introduced into the market). However, the decommissioning of the old equipment disrupts this expectation in several past years. Since previous submission, expectations about the reduction of emissions were fully met. When compared with the previous year, the reduction in emissions was ca 48 Gg CO<sub>2</sub> eq. At verifying of this decrease, synergistic effect of several facts was determined: (i) significant increase of recovery, mainly R404A and R410A blends since 2022. In comparison with the previous year, the increased recovery has no significant effect on the this year decrease.; (ii) reduction in usage of the blend with high GWP (R404A, R410A, R407C and R134a) and their replacement with the blends with low GWP. This reduction corresponds to the decrease in emissions by ca 11 Gg CO<sub>2</sub> eq.; (iii) decrease in decommissioning of devices in categories 2.F.1.a, 2.F.1.d and 2.F.1.f that corresponds to the decrease in emissions by 40 Gg CO<sub>2</sub> eq. Totally, it corresponds to the decrease in emissions by 51 Gg CO<sub>2</sub> eq.; which is in very solid agreement with the actual decrease of 48 Gg CO<sub>2</sub> eq.

It can be also seen that the small decreasing trend (with fluctuations) occurs since 2017.

The emissions of NF<sub>3</sub> and SF<sub>6</sub> are not occurring in this category. The following gases and subcategories are reported in 2.F.1:

- HFC-23, HFC-32, HFC-125, HFC-134a, HFC-143a, HFC-152a and C<sub>2</sub>F<sub>6</sub> in 2.F.1.a - Commercial refrigeration.
- HFC-134a in 2.F.1.b - Domestic refrigeration.
- HFC-32, HFC-125, HFC-134a, HFC-227ea and HFC-143a in 2.F.1.c - Industrial refrigeration.
- HFC-32, HFC-125, HFC-134a and HFC-143a in 2.F.1.d - Transport refrigeration.
- HFC-134a in 2.F.1.e - Mobile AC.
- HFC-32, HFC-125, HFC-134a, HFC-143a and C<sub>4</sub>F<sub>8</sub> in 2.F.1.f - Stationary AC.

**Figure 4.13:** The share of individual subcategories within the category 2.F.1 in 2023




The products designed for coolants R22, R134a and R404A were usually imported up to the year 1998. Only in 1999, the indications of import of products containing coolants R407C and R410A are emerging. When the Act No. 76/1998 on the Protection of the Ozone Layer of the Earth entered into force on April 1, 1998, the use of the alternative coolants started. Consumption of alternative coolants R401A and R409A in order to supply R22 started to decrease in the year 2002. Coolants R407C and R410A show the increase since 1999. Coolant R134a shows continuing increasing trend mainly because of rising use of cars with AC; the increasing trend stopped in 2010. It is caused by smaller purchases of cars in Slovakia and lower amount of gas in AC since then, which results in smaller bank of HFC-134a in Slovakia. General increasing trend of HFCs emissions visible since the base year has been caused by

supplying HCFCs gases by the HFCs. However, because of the almost complete replacement of HCFCs gases, the F-gases emissions were approximately constant since 2010. Rising trend since 2014 is caused by increased decommissioning of refrigerant units, while the decreasing trend since 2016 is caused by increased using of HFCs with lower GWP. Servicing of the MACs with HFC-134a is lower than in previous years, therefore the operational emissions decreased. On the other hand, the servicing with HFO-1234YF increased.

The decreased in 2018 was followed after a peak in 2017. This can be explained by the decreasing of share of mixtures containing major share of HFC-134a and HFC-125 increasing of the share of mixtures with a higher HFC-32 content. This is mostly visible in subcategory 2.F.1.f. In 2023, the replacement of HFC-404A with the blends HFC-448A and HFC-452A occurred in an increased extent. Also the replacement of HFC-410A with HFC-452B was more significant than in the previous year. The replacement of HFC-134a with the HFC-513A blend is negligible, but using of R1234yf, R1234ze is of increased importance. The use of natural refrigerants is of increased importance in this submission, as well.

Approximately 36% of total F-gases emissions (in CO<sub>2</sub> eq.) are allocated in 2.F.1.a – Commercial Refrigeration followed by 2.F.1.e – Mobile AC (32%) in 2023 (*Figure 4.13*). This relates to the high share of automotive industry in last years in Slovakia. About 20% emissions are allocated in 2.F.1.f – Stationary AC, 9% in 2.F.1c, 3% in 2.F.1.d and below 1% in 2.F.1.b – Domestic Refrigeration. Time series of F-gases consumption in the category 2.F.1 is summarized in the following *Tables 4.45-4.52*.

*Figure 4.14: The share of individual F-gases in the category 2.F.1 in 2023*



*Table 4.45: Aggregated data on HFCs use in the subcategory 2.F.1.a in particular years*

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
<i>Gg CO<sub>2</sub> eq.</i>									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	2.304	4.056	4.056	NO	0.023	0.767	NO	NO	0.790
2000	34.303	59.940	121.623	NO	0.343	17.941	NO	NO	18.284
2005	84.295	110.225	625.353	3.130	0.843	89.274	2.381	0.749	92.497
2010	117.202	136.799	1338.124	35.500	1.172	200.687	29.086	6.415	230.945
2011	84.876	135.435	1401.600	53.453	0.849	185.001	43.892	9.562	229.741
2012	72.657	141.886	1454.765	67.925	0.727	204.650	55.846	12.079	261.222
2013	78.970	153.151	1504.833	80.128	0.790	214.583	65.974	14.154	281.347

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
<i>Gg CO<sub>2</sub> eq.</i>									
2014	100.696	91.000	1484.649	86.239	1.007	212.735	68.497	17.742	282.242
2015	111.966	104.003	1463.439	97.401	1.120	236.280	82.012	15.389	319.419
2016	101.551	120.448	1449.254	102.160	1.016	227.965	73.842	28.318	302.829
2017	70.463	78.876	1373.565	120.336	0.705	216.399	111.025	9.311	328.139
2018	50.753	47.398	1247.467	139.448	0.508	194.369	125.379	14.069	320.266
2019	40.706	39.999	1120.365	134.361	0.407	179.086	125.335	9.026	304.864
2020	12.383	32.743	985.480	137.959	0.124	139.440	119.734	18.225	259.327
2021	4.900	20.181	854.497	123.962	0.049	153.227	107.159	16.803	260.478
2022	14.468	16.530	732.367	113.386	0.145	117.009	44.635	68.751	161.815
2023	19.747	21.425	625.600	104.236	0.197	110.295	33.108	71.128	143.627

**Table 4.46: Aggregated data on PFCs use in the subcategory 2.F.1.a in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
<i>Gg CO<sub>2</sub> eq.</i>									
1990-2018	NO	NO	NO	NO	NO	NO	NO	NO	NO
2019	NO	0.707	0.707	NO	NO	NO	NO	NO	NO
2020	NO	NO	0.707	NO	NO	NO	NO	NO	NO
2021	NO	NO	0.707	NO	NO	0.036	NO	NO	0.036
2022	NO	NO	0.707	NO	NO	0.029	NO	NO	0.029
2023	NO	NO	0.707	NO	NO	NO	NO	NO	NO

**Table 4.47: Aggregated data on HFCs use in the sub-category 2.F.1.b in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
<i>Gg CO<sub>2</sub> eq.</i>									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	NO	NO	NO	NO	NO	NO	NO	NO	NO
2000	13.130	12.998	56.741	NO	0.131	0.284	NO	NO	0.415
2005	1.586	1.359	74.942	NO	0.016	0.375	NO	NO	0.391
2010	NO	0.189	69.975	4.290	NO	0.350	2.574	1.716	2.924
2011	NO	5.779	64.386	8.814	NO	0.322	5.288	3.526	5.610
2012	NO	9.566	59.364	11.414	NO	0.297	6.848	4.565	7.145
2013	NO	12.486	54.848	13.363	NO	0.274	8.018	5.345	8.292
2014	NO	1.661	43.933	9.842	NO	0.220	4.468	5.374	4.688
2015	NO	0.018	36.211	6.016	NO	0.181	4.314	1.703	4.495
2016	NO	NO	31.576	3.708	NO	0.158	2.610	1.097	2.768
2017	NO	NO	29.038	2.030	NO	0.145	1.441	0.589	1.587
2018	NO	NO	27.106	1.546	NO	0.136	1.283	0.263	1.418
2019	NO	NO	25.990	0.893	NO	0.130	0.850	0.043	0.980
2020	NO	NO	25.179	0.649	NO	0.126	0.598	0.051	0.724
2021	NO	NO	24.655	0.419	NO	0.123	0.383	0.037	0.506
2022	NO	NO	24.438	0.173	NO	0.122	0.140	0.033	0.262
2023	NO	NO	22.835	1.283	NO	0.114	0.532	0.751	0.646

**Table 4.48: Aggregated data on HFCs using in the sub-category 2.F.1.c in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	3.423	0.655	0.655	NO	0.034	0.097	NO	NO	0.131
2000	39.834	8.680	17.306	NO	0.398	2.278	NO	NO	2.676
2005	96.958	16.088	91.300	NO	0.970	11.776	NO	NO	12.746
2010	134.340	13.730	203.361	0.105	1.343	25.980	0.064	0.041	27.387
2011	145.366	142.015	344.924	0.323	1.454	32.488	0.218	0.105	34.160
2012	91.984	73.893	418.077	0.517	0.920	53.625	0.366	0.152	54.910
2013	83.479	54.283	471.121	0.894	0.835	56.148	0.665	0.229	57.647
2014	51.382	53.585	522.659	1.380	0.514	46.648	0.965	0.415	48.126
2015	48.774	51.768	570.689	2.664	0.488	53.420	2.203	0.461	56.111
2016	47.303	51.108	615.858	4.295	0.473	54.831	3.105	1.190	58.408
2017	47.643	49.815	657.014	6.575	0.476	59.532	6.040	0.535	66.049
2018	54.514	91.715	737.732	8.334	0.545	49.106	7.500	0.834	57.151
2019	20.907	60.144	784.263	10.654	0.209	44.038	9.978	0.676	54.225
2020	6.014	14.431	783.419	11.839	0.060	43.186	10.305	1.534	53.552
2021	4.294	18.233	784.433	13.744	0.043	46.431	11.905	1.839	58.380
2022	42.621	87.606	853.999	13.898	0.426	27.610	5.310	8.588	33.346
2023	61.484	80.407	913.248	16.573	0.615	28.263	5.100	11.473	33.978

**Table 4.49: Aggregated data on HFCs using in the sub-category 2.F.1.d in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	NO	1.156	1.156	NO	NO	0.243	NO	NO	0.243
2000	2.843	4.299	12.348	NO	0.028	2.180	NO	NO	2.208
2005	6.676	9.527	47.177	0.703	0.067	7.039	0.447	0.256	7.552
2010	10.582	12.772	93.762	3.282	0.106	14.595	2.504	0.778	17.205
2011	13.192	18.099	103.691	3.714	0.132	14.301	2.830	0.884	17.263
2012	11.283	15.449	109.836	4.194	0.113	15.786	3.207	0.987	19.106
2013	5.267	6.388	105.809	4.718	0.053	15.363	3.627	1.091	19.043
2014	1.759	1.752	95.709	5.503	0.018	12.500	3.967	1.535	16.485
2015	4.888	4.888	87.149	5.889	0.049	22.433	4.770	1.119	27.252
2016	3.423	9.424	81.880	6.811	0.034	23.584	4.823	1.988	28.442
2017	2.857	6.204	72.189	8.308	0.029	20.874	7.368	0.939	28.271
2018	3.628	4.491	61.170	7.458	0.036	20.187	6.575	0.883	26.798
2019	1.191	2.074	48.603	7.718	0.012	20.197	7.209	0.509	27.418
2020	0.611	1.285	37.754	8.387	0.006	14.870	7.355	1.032	22.231
2021	NO	0.352	30.799	5.459	NO	17.994	4.748	0.711	22.742
2022	3.249	3.249	29.395	2.035	0.032	14.818	0.821	1.214	15.672
2023	4.660	4.819	28.811	1.660	0.047	13.253	0.533	1.127	13.833

**Table 4.50: Aggregated data on HFCs using in the sub-category 2.F.1.e in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	21.479	59.207	59.207	NO	0.215	6.477	NO	NO	6.692
2000	64.437	76.822	386.615	NO	0.644	42.296	NO	NO	42.940
2005	107.395	116.232	896.882	NO	1.074	98.119	NO	NO	99.193
2010	240.630	116.719	1314.360	30.053	2.406	165.772	18.032	12.021	186.211
2011	275.132	130.330	1376.452	31.323	2.751	149.845	18.794	12.529	171.390
2012	398.192	69.680	1372.319	32.362	3.982	136.871	19.417	12.945	160.271
2013	412.867	56.810	1346.229	36.903	4.129	131.095	22.142	14.761	157.365
2014	324.723	56.633	1311.294	41.455	3.247	150.464	18.820	22.634	172.532
2015	484.940	67.508	1279.104	45.971	4.849	154.210	32.961	13.010	192.021
2016	404.641	31.178	1203.225	50.536	4.046	105.939	35.578	14.959	145.563
2017	196.198	24.065	1114.561	53.803	1.962	124.123	38.200	15.603	164.285
2018	260.464	12.165	1008.764	56.810	2.605	106.876	47.152	9.658	156.633
2019	225.507	22.569	909.850	58.914	2.255	107.765	56.057	2.857	166.076
2020	159.243	16.684	801.903	61.017	1.592	105.410	56.202	4.814	163.205
2021	93.142	9.800	686.895	63.492	0.931	106.809	57.965	5.528	165.706
2022	79.969	8.516	576.428	63.157	0.800	105.530	50.984	12.173	157.313
2023	89.676	9.377	474.149	64.156	0.897	100.676	26.586	37.570	128.159

**Table 4.51: Aggregated data on HFCs using in the sub-category 2.F.1.f in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	12.711	12.711	12.711	NO	0.127	2.219	NO	NO	2.346
2000	32.330	21.297	94.095	NO	0.323	15.345	NO	NO	15.669
2005	71.963	37.700	268.953	NO	0.720	39.815	NO	NO	40.535
2010	107.306	37.149	515.882	11.465	1.073	70.306	7.034	4.431	78.413
2011	106.682	101.112	601.346	12.017	1.067	81.501	7.493	4.524	90.061
2012	121.818	114.980	698.620	13.234	1.218	61.773	8.498	4.736	71.489
2013	90.259	81.340	758.645	15.809	0.903	58.391	10.374	5.435	69.668
2014	46.751	86.929	819.165	19.962	0.468	61.813	11.667	8.295	73.948
2015	98.800	131.979	919.897	24.080	0.988	56.564	17.700	6.380	75.252
2016	54.706	95.748	980.302	27.495	0.548	59.031	18.041	9.454	77.620
2017	251.464	177.959	1 119.831	29.843	2.515	64.252	24.001	5.842	90.768
2018	95.527	142.743	1 219.699	32.944	0.955	55.477	26.907	6.037	83.339
2019	65.819	78.825	1 251.088	35.751	0.658	70.266	31.971	3.780	102.896
2020	53.862	50.168	1 245.502	43.697	0.539	77.947	37.520	6.177	116.006
2021	703.689	124.957	1 307.739	49.773	7.037	86.083	41.734	8.039	134.854
2022	107.113	81.197	1 327.107	46.686	1.071	59.878	17.983	28.703	78.932
2023	68.479	70.143	1 323.964	56.888	0.685	64.053	13.940	42.948	78.678

**Table 4.52: Aggregated data on PFCs use in the subcategory 2.F.1.f in particular years**

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions From:			Recovery	Total
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990-2022	NO	NO	NO	NO	NO	NO	NO	NO	NO
2023	NO	3.535	3.535	NO	NO	NO	NO	NO	NO

### Methodological Issues

The IPCC 2006 GL describe two tiers for estimating emissions. The bottom-up approach takes into account the time lag between consumption and emissions explicitly through emission factors. The top-down approach, the time lag is taking into account implicitly, by tracking the amount of virgin chemicals consumed in a year that replaces emissions from the previous year.

The web reporting system used in Slovakia allows calculating emissions in both approaches. The bottom-up approach is combined with the top-down approach. The procedure is as follows:

1. Using the bottom-up approach (tier 2a) from the Logbook Leaklog;
2. Calculation of the total consumptions of individual gases in Slovakia based on the Leaklog (tier 2a);
3. Calculation of the total consumption of individual gases in Slovakia according to the top-down approach (tier 2b);
4. Comparison of the total consumptions calculated by these two approaches;
5. If differences above 2% occur, the data for bottom-up approach are corrected as follows (expert judgement based on the QA activity introduced in 2011):
  - R134a: Difference is added to leakage from mobile AC,
  - R404A: Difference is added between new charge/recharge 0.2/0.8,
  - R407C: Difference is added to new charge of stationary AC,
  - R410A: Difference is added to leakage from industrial refrigeration and stationary AC 0.1/0.9.
6. If differences below 2% occur, the data for bottom-up approach are corrected proportionally according to the operational emissions.
7. Calculation of emission estimates by the bottom-up approach using the corrected data.

In 2023, no significant corrections were necessary, the differences between top-down and bottom-up approaches were up to 2%. Following formulas (tier 2b, top-down approach) based on the structure of the reporting systems were used:

Emissions = Annual Sales of New Refrigerant – Total Charge of New Equipment + Disposal Emissions  
 where *Annual Sales* and *Total Charge of New Equipment* are calculated by formulas presented in the IPCC 2006 Guidelines, Chapter 3, p. 7.54 (not simplified formulas).

Following formulas (tier 2a, bottom-up approach) based on the structure of the reporting systems were used: Emissions = Emissions from new fillings + Operational emissions + Disposal emissions

where: *Emissions from new fillings* represent 1% (EF) of the new charges filled in Slovakia (*Chemicals to Charge Domestically Manufactured and Assembled Equipment + Chemical to Charge Equipment that is not Factory-Charged*).

*Operational emissions*: The approach described in IPCC 2006 GL assumes that servicing of equipment restocks the bank of single chemical and thus the amount of gas used for servicing represents the

operational emissions. Slovakia adopted this assumption with a modification. The servicing of equipment restocks the bank of chemical and its amount used at servicing equals to the emissions. However, equipment that is few years before decommissioning is not serviced and bank is not restock at this equipment. Therefore, the operational emissions are composed from two terms in this submission: (i) data from servicing of equipment; (ii) emissions from non-serviced equipment few years before its decommissioning. The first term in the operational emissions represents the consumption of gases for servicing and container management (these data are reported in Leaklog). It is assumed that the chemical used for servicing restocks the emissions from the bank and thus the bank of the chemical remains constant. The second term in operational emissions represents emissions from non-serviced equipment few years before its decommissioning. These emissions decrease the amount of chemical in equipment and the equipment contains only a part of the chemical at its decommissioning. The emissions are calculated by using product life factor that are presented in [Table 4.3](#) The product life factors, number of years when the equipment is not serviced and fraction of gas remaining at the decommissioning of equipment presented in [Table 4.53](#) are consistent and they are based on the default factors presented in IPCC 2006 GL. These emissions do not restore the bank of the chemical and are subtracted from the bank.

**Table 4.53: Product life factor of not serviced equipment; number of years, when the equipment is not serviced and ratio of initial charge that is remaining at decommissioning of equipment**

Category	Product Life Factor	Years Before Retirement	Initial Charge Remaining at Retirement
2.F.1.a	10%	2	80%
2.F.1.b	0.5%*	12-15*	80%*
2.F.1.c	20%	1	80%
2.F.1.d	25%	2	50%
2.F.1.e	16.67%	3	50%
2.F.1.f	10%	2	80%

\* Default IPCC 2006 GL values

*Disposal emissions* represent the emissions from the retired equipment. Since 2014, the recycling companies report the data about recovery of gases in database Leaklog. There is available amount of gas that is recovered, reused and destroyed in recycling factories. All these terms are covered in CRT term "recovery". The amount of recovered gas is known and comparison with the amount of gas in decommissioned equipment can be made. The fractions of gases that are recovered from disposed equipment in 2023 are presented in [Table 4.54](#). Differentiating of the gases among the subcategories is not possible, only total data for each gas is available. Therefore, the same fraction of recovered gas is assumed in all categories. The annual data of the recovery ratio of the individual gases for whole time series is presented in [Table 4.55](#). For years before 2013, the average value of the years 2014 and 2015 is assumed.

**Table 4.54: Comparison of amount of gases in retired units and amount of recovered gases in 2023**

F-GAS	Amount in Retired Equipment	Recovery Amount	Ratio
	<i>t</i>		
HFC-23	0.017	NO	0.0%
HFC-32	14.872	12.7018	85.4%
HFC-125	25.214	19.238	76.3%
HFC-134a	63.604	37.247	58.6%
HFC-143a	14.979	9.7904	65.4%
HFC-152a	NO	NO	-



**Table 4.55: Aggregated data on HFCs recovery ratio (%) in the category 2.F.1. in particular years**

YEAR	HFC-23	HFC-32	HFC-125	HFC-134a	HFC-143a	HFC-152a
	%					
2013 and before	-	55.0	25.0	40.0	13.0	-
2014	-	68.2	27.3	54.6	15.6	-
2015	-	43.7	22.7	28.3	11.9	-
2016	-	49.5	34.6	29.6	24.4	-
2017	-	29.8	12.6	29.0	3.9	-
2018	-	30.5	16.0	17.0	6.3	-
2019	-	17.5	11.1	4.9	4.0	-
2020	-	15.9	15.9	7.9	11.7	-
2021	-	17.0	18.1	8.7	11.1	-
2022	-	72.0	68.9	19.3	60.0	-
2023	-	85.4	76.3	58.6	65.4	-

For the consistency of operational emissions, it is necessary to follow the bank of chemical. The bank is calculated as follows:

$$\text{Bank}_{\text{in year } t} = \text{Bank}_{\text{in year } t-1} + \text{New additions to bank} - \text{Chemical in retired equipment} - \text{Operational emissions from non-serviced equipment}$$

Where: New additions to bank = Chemicals to Charge Domestically Manufactured and Assembled Equipment + Chemicals to Charge Equipment that is not Factory-Charged + Chemicals Contained in Imported Equipment Already Charged – Chemicals Contained in Exported Equipment Already Charged.

It should be mentioned that due to the last two terms in the above relationship the using of the data about new fillings from CRT software is not possible for the calculation of the bank (stock). Calculation of the bank has to contain data that includes import and export of already filled equipment.

Emissions factor from the filling chemicals into new equipment (product manufacturing factor) is assumed 1% (based on the producers' data) for all categories and gases in the 2.F.1.

Operational emission factor (product life factor) is calculated annually based on the reported operational emissions and the respective bank.

Disposal emission factor (disposal loss factor) is based on the data of recycling companies. The fractions of gases that are recovered from disposed equipment are presented in [Table 4.55](#). The average value of the years 2014 and 2015 is assumed since 2013 and back to base year 2013.

Activity data were collected via web reporting system and treated as described above and in the [Annex 4.2](#) of this Document.

**2.F.1.a – Commercial Refrigeration:** This category includes emissions from manufacturing, assembly, installation of small refrigeration equipment mostly for export (“stand-alone” commercial application including also some equipment for domestic refrigeration) and emissions from refrigeration in supermarkets and other commercial refrigeration. Only one company manufactures smaller “stand-alone” equipment for commercial and refrigeration (fridges, freezers) with the HFC R-134a and R-404A as cooling agents. This equipment is mostly exported. Data on F-gas consumption are reported through web reporter. Emissions from commercial refrigeration manufacturing, assembly, installation are estimated to equal 1%. No detailed figure on the installed equipment is available. Data on consumption for new systems and refilling were provided by the main service companies through web reporting; the stocks were calculated accordingly. Used refrigerants R-32, R-404A, R-410A, R-134a, R-600A, R-407C, R-717, R-723, R-449A, R-290, R-513A, R-452A, R-507, R-508B, R-407F, R-407H, R-417A, R-448A, R-407A, C<sub>5</sub>H<sub>12</sub>, R-1234yf, R-23, CO<sub>2</sub>, R-1234ze, R-422D, R-437A, R-22, R-143a, R-

---

454B. Lifetime of equipment was assumed 9-12 years. Nameplate capacity of retired equipment was calculated as follows:

Retired equipment<sub>in year t</sub> = New addition to stock<sub>in year t-12</sub> / 4 + New addition to stock<sub>in year t-11</sub> / 4 + New addition to stock<sub>in year t-10</sub> / 4 + New addition to stock<sub>in year t-9</sub> / 4

The fraction of the gas that remained in the retired equipment is presented in [Table 4.53](#) and the recovered fraction is presented in [Table 4.55](#).

**2.F.1.b – Domestic Refrigeration:** Partially also the HFC-134a is used for domestic use as refrigerant in refrigerators (fridges and freezers). HFC-134a as refrigerant was introduced by industry at the end of 1995 as replacement of CFC-12. In the following years (starting in 1999) it was gradually replaced by R600a (isobutane). Charging of refrigerators with R134a was stopped by the end of 2006. The calculation of operational emissions is different in this category. The domestic refrigeration units are not serviced usually. Therefore, we used the default product life factor (0.5%) and it is assumed that the emissions decrease the bank of the chemical. Lifetime of domestic refrigeration equipment was assumed 12-15 years.

Nameplate capacity of retired equipment was calculated as follows:

Retired equipment<sub>in year t</sub> = New addition to stock<sub>in year t-15</sub> / 4 + New addition to stock<sub>in year t-14</sub> / 4 + New addition to stock<sub>in year t-13</sub> / 4 + New addition to stock<sub>in year t-12</sub> / 4

The fraction of the gas that remained in the retired equipment is presented in [Table 4.53](#) and the recovered fraction is presented in [Table 4.55](#).

**2.F.1.c – Industrial Refrigeration:** In industrial refrigeration, refrigerants are used for production process, e.g. in chemical industry to keep definite process temperatures or in food industry for cooling/freezing partly inoculating with commercial refrigeration. In contrast to commercial refrigeration, in the **IPPU sector** not only HFC/HCFC refrigerants play the significant role, but also NH<sub>3</sub> and CO<sub>2</sub>, as well. The refrigeration systems are normally maintained by service companies. Refrigerants are R-134a, R-513A, R-404A, R-407C, R-452A, R-32, R-410A, R-449A, R-1234yf, R-22, R-407A, R-417A, R-422A, R-448A, R-143A, R-407H, R-507, R-152a, R-407F, R-600A, CO<sub>2</sub>, R-290, R-425A, R-452B, R-401A, R-170, R-50, R-23, R-1234ze, R-514A, R-515B, R-422D, R-227ea, R-454B;R-450A. Lifetime of equipment was assumed 15-19 years. Nameplate capacity of retired equipment was calculated as follows:

Retired equipment<sub>in year t</sub> = New addition to stock<sub>in year t-19</sub> / 5 + New addition to stock<sub>in year t-18</sub> / 5 + New addition to stock<sub>in year t-17</sub> / 5 + New addition to stock<sub>in year t-16</sub> / 5 + New addition to stock<sub>in year t-15</sub> / 5

The fraction of the gas that remained in the retired equipment is presented in [Table 4.53](#) and the recovered fraction is presented in [Table 4.55](#).

**2.F.1.d – Transport Refrigeration:** This group includes refrigerated road vehicles. Recently used refrigerants are: R-134a, R-600A, R-404A, R-143a, R-123, CO<sub>2</sub>, R-452A, R-32, R-410A, R-449A, R-1234yf, R-507, R-407C, R-448A, R-290, R-124, R-417A, R-422D, R-23, R-407F. Manufacturing of refrigeration units takes place in Slovakia only in very small scale. Emissions occur mainly from stock and from disposal. Lifetime of equipment was assumed 8-9 years. Nameplate capacity of retired equipment was calculated as follows:

Retired equipment<sub>in year t</sub> = New addition to stock<sub>in year t-9</sub> / 2 + New addition to stock<sub>in year t-8</sub> / 2

The fraction of the gas that remained in the retired equipment is presented in [Table 4.53](#) and the recovered fraction is presented in [Table 4.55](#).

**2.F.1.e – Mobile AC:** Mobile air conditioning includes passenger cars, trucks, busses, agricultural machines, rail and manufacturing of vehicles for construction sites. The use of R-134a for mobile air conditioning started in 1995. New charges filled into vehicles are taken from the automobile producers in Slovakia. New additions to stock are calculated from the registrations of vehicles in Slovakia and

compared with the records of official manufacturers and importers of cars. The following time series of the share of cars with MAC is assumed: (i) in years 1995 – 1999, 70% of registered cars contained MAC; (ii) in 2000 – 2003, 80% of registered cars contained MAC; (iii) in 2004 – 2011, 90% of registered cars contained MAC; (iv) 100% of registered cars contained MAC since 2012. The presented shares are based on the data of car manufacturers in Slovakia. We assume that the share is a typical one and it is applied to the rest of cars.

In 2023, 97 110 new vehicles were registered in Slovakia. In these vehicles, the average charge was assumed 0.062 kg of HFC-134a per one new car in 2023. The average charge is based on the data from car manufacturers in Slovakia (number of produced cars; consumption of HFC-134a and HFO-1234yf necessary to fill them. In 2023, the average charge was 0.624 kg while the share of HFC-134a was 10.0%). We assume that a similar average charge can be used for cars that are not produced in Slovakia. The number of imported and registered second-hand vehicles was 18 497 pcs. HFC-134a charge in these vehicles was assumed to be as in new registered vehicles. The time series of the HFC load into new vehicles is presented in [Table 4.56](#).

**Table 4.56: Loads of HFCs into new vehicles**

YEAR	Number of Produced Vehicles	Amount of HFC-134a Used In New Vehicles	Amount of HFC-1234yf Used in New Vehicles	Fraction of HFC-134a From Total HFC Use	Average HFC Load per One Vehicle	Average HFC-134a Load per One Vehicle
	No	t	t		kg	kg
2016	1 095 191	310.517	354.577	0.4669	0.607	0.284
2017	1 266 289	150.15	386.606	0.2797	0.424	0.119
2018	1 093 215	199.95	532	0.2732	0.67	0.183
2019	1 122 067	173.113	461.324	0.2729	0.565	0.154
2020	990 598	122.211	444.738	0.2156	0.572	0.123
2021	1 000 030	71.648	508.351	0.1235	0.580	0.072
2022	970 275	61.128	541.71	0.1014	0.621	0.063
2023	1 080 000	67.384	606.12	0.1001	0.624	0.062

Lifetime of equipment was assumed 12-15 years. Nameplate capacity of retired equipment was calculated as follows:

Retired equipment<sub>in year t</sub> = New addition to stock<sub>in year t-15</sub> / 4 + New addition to stock<sub>in year t-14</sub> / 4 + New addition to stock<sub>in year t-13</sub> / 4 + New addition to stock<sub>in year t-12</sub> / 4

The fraction of the gas that remained in the retired equipment is presented in [Table 4.53](#) and the recovered fraction is presented in [Table 4.55](#).

**2.F.1.f – Stationary AC:** This category includes stationary air conditions, room air conditions and heat pumps. Plants for waste heat recovery are included in this category, as well (we are not able to distinguish between them and heat pumps). Stationary air conditions include large equipment >20 kW. Data on consumption for new systems and refilling are provided by service organizations since 2009 via web reporting, the stocks are calculated accordingly. Room air conditions are in the contrast with the stationary AC (a comparable sector in terms of HFCs consumption for new and refilling). Room AC systems include small mobile and compact equipment to be installed at windows or walls, fixed split- and multisplit systems up to 20 kW and larger Variable Refrigerant Flow (VRF) or Multi Air Conditioning systems. Small equipment, split- and multisplit systems and VRF systems are imported already charged with refrigerant. The installation of heat pumps with the HFCs started in Slovakia mainly in the 2004. Heat pumps are manufactured in Slovakia and imported, as well. Used F-gases in this subcategory are: R-407C, R-32, R-134a, R-410A, R-1234yf, R-449A, R-404A, R-407H, R-507, CO<sub>2</sub>, R-417A, R-290, R-600A, R-407A, R-437A, R-448A, R-452A, R-401A, R-513A, R-23, R-22, R-143a, R-422D, R-422A, R-407F, R-405A, R-454B, R-452B, R-424A, R-1234ze. Lifetime of air conditioning equipment and heat pumps was assumed 12-15 years. Nameplate capacity of retired equipment was calculated as follows:

---

Retired equipment<sub>in year t</sub> = New addition to stock<sub>in year t-15</sub> / 4 + New addition to stock<sub>in year t-14</sub> / 4 + New addition to stock<sub>in year t-13</sub> / 4 + New addition to stock<sub>in year t-12</sub> / 4

The fraction of the gas that remained in the retired equipment is presented in [Table 4.53](#) and the recovered fraction is presented in [Table 4.55](#).

### Uncertainties and Time-series Consistency

A consistent time series of HFCs import-export exists since 1995 and is well documented. In previous submissions, the data on banks were not consistent because of use of extrapolation in 2010. The recalculation of banks was made according to the discussion with the ERT during six weeks' period after in-country review in 2012. In that submission, this extrapolation was useful in order to differentiate the banks among different subcategories. The inconsistency was caused by two types of formula used for bank calculation. In the 2015 submission, the inconsistent bank data were corrected, the bank data were recalculated by the same formula (as presented above) in the whole time series.

In 2017 submission, the bank data were recalculated again. The reason of recalculation was new way of operational emission estimation. It was assumed that equipment few years before its decommissioning is not serviced and the operational emission from this equipment has to be subtract from the bank. New product life factors were estimated based on this assumption. Product life factors for the time series 1990 – 2009 were assumed average of product life factors in the period 2010 – 2013 (outliers were excluded from the average). The used product life factors are presented in [Table 4.57](#) and they are within the range presented in the IPCC 2006 GL. The reported emissions are also influenced by the recalculation of the disposal emissions in last reporting years. Amounts of HFCs in retiring equipment were calculated consistently according to the previous presented formulas for all subcategories.

The changes in trend in new fillings in 2.F.1.e are caused by manufacturers of cars. Three factories exist in Slovakia. One of them has been producing cars since 1995, the others since 2006. Since 2015 submission the new fillings emissions are calculated from the data provided by the car producers (HFCs used for new fillings) for the years since 2009. For the rest of the time series the new fillings were estimated based on car production. The following time series of the share of cars with MAC is assumed: (i) in years 1995 – 1999, 70% of registered cars contained MAC; (ii) in 2000 – 2003, 80% of registered cars contained MAC; (iii) in 2004 – 2011, 90% of registered cars contained MAC; (iv) 100% of registered cars contained MAC since 2012.

The emissions in the category 2.F.1.f have stable trend since 2012 (inter-annual changes are up to 5%). The decrease in 2.F.1.a in 2011 was caused by the economic situation on the market and decrease in use of R404A in that year from service. The decrease in 2.F.1.c in 2014 is caused by decrease in service of units containing R404A and by starting to use units with the mixtures containing R152a with lower GWP.

Generally, the decrease due the Regulation (EU) No 517/2014 of the European parliament and of the Council on fluorinated greenhouse gases is expected. The expectations are caused by several reasons. One of them is end of using of new R404A gas with GWP 3940 (only recovered gas can be used since now). Another reason is using of new replacements of R404A, R410A, R134a with low GWP blends (in Slovakia new gases R448A, R449A, R454B, R454C, R513A, R514A, R1234yf, R1234ze were introduced into the market). However, the decommissioning of the old equipment disrupts this expectation in several past years. In this submission, expectations about the reduction of emissions were fully met. In 2022, when compared with the previous year, the reduction in emissions was ca 195 Gg CO<sub>2</sub> eq. At verifying of this decrease, synergistic effect of several facts was determined: (i) significant increase of recovery, mainly R404A and R410A blends. The increased recovery corresponds to the decrease in emissions by ca 80 Gg CO<sub>2</sub> eq.; (ii) reduction in usage of the blend with high GWP (R404A, R410A, R407C and R134a) and their replacement with the blends with low GWP. This reduction corresponds to the decrease in emissions by ca 90 Gg CO<sub>2</sub> eq.; (iii) decrease in decommissioning of

devices in categories 2.F.1.a, 2.F.1.d and 2.F.1.f that corresponds to the decrease in emissions by 13 Gg CO<sub>2</sub> eq. Totally, it corresponds to the decrease in emissions by 183 Gg CO<sub>2</sub> eq.; which is in very solid agreement with the actual decrease of 195 Gg CO<sub>2</sub> eq.

**Table 4.57: Product life factors of individual gases in the category 2.F.1 in 1990 – 2009**

CATEGORY	HFC-125	HFC-134a	HFC-143a	HFC-152a	HFC-23	HFC-32
	%					
2.F.1.a	14.20	19.20	13.93	22.30	10	NO
2.F.1.b	NO	0.50	NO	NO	NO	NO
2.F.1.c	12.46	15.00	12.92	NO	NO	9.72
2.F.1.d	12.59	21.04	12.28	NO	NO	12.95
2.F.1.e	NO	10.94	NO	NO	NO	NO
2.F.1.f	12.97	17.48	8.61	NO	NO	9.62

A detailed look at the product life factors in the subcategories 2.F.1.a, 2.F.1.c, 2.F.1.d and 2.F.1.f shows large variations as well as inconsistent development of reported product life factors. It is due to several factors:

- (i) The methodology used (hybrid tier 2a/tier 2b). It is assumed that all leaks in a given year will be replenished in a given year during equipment servicing. The practice of having all equipment serviced every year is not common in Slovakia. A typical example is MAC in cars, where, according to the automotive association, MAC service is performed on average at 2-4 annual intervals. This results in year-on-year jumps in servicing emissions, i.e. PLF. This is also very noticeable with gasses that have a small bank, each small increase in servicing will be reflected in a jump in PLF.
- (ii) By reporting organizations in the Leak log database. Organizations do not report each individual activity to the database but they report it at certain time intervals (e.g. monthly). Then they report the type of gas and all the operations they performed with it in summary and assign it to the category in which they performed the most activities. For example, the organization can carry out multiple replacements for leaks with R410A gas, with 60% of these replacements falling into subcategory 2.F.1.f, but the remaining 40% into other categories. However, the organization reports all additions to leaks in one record and assigns it to category 2.F.1.f, because that's where the gas was used the most. We're working with the assumption (since there are several hundreds of organizations involved in this activity) that when processing the database for the whole year, these deviations compensate each other. However, they can only partially compensate. Thus, the division of F-gases into individual subcategories may not be completely correct, and this may cause year-to-year changes in individual subcategories. On the other hand, the emissions calculated by our procedure are certainly more accurate than the calculation using some more general emission factor. It should also be noted that although the division into categories 2.F.1.a, 2.F.1.c, 2.F.1.d and 2.F.1.f may not be completely correctly allocated for every year, the output from the overall category 2.F.1 is correct, because all movements of F-gases are already summarized there.

Due to this large variations and inconsistencies, the QA/QC procedure has been performed in this submission. The details about the procedure and results can be found in [Annex 4.4](#).

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission.

Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.12.10. Foam Blowing (CRT 2.F.2)

This category is not significant and includes F-gases used in industry as follow:

- PU foam appliances (transferred from blowing agent R141b directly to cyclopentane in 1998).
- Injected PU foams in commercial cooling (started in 1999 and transferred from blowing agent R134a to water in 2007).
- Sprayed PU foams for roofs (transferred directly from ODS to HFC245fa and 365mfc in 2002).
- PU panels for containers, store rooms, etc. Big importers imported only panels with hydrocarbons, water blowing agents; smaller importers (in opened market) imported panels with R134a from 1999 up to 2007. In the main application areas of PU hard foam (rigid foam insulating panels, flexibly coated; rigidly faced sandwich panels) hydrocarbons and CO<sub>2</sub> are usually used as blowing agent. In the area of PU insulating foam for pipes HFC-245fa and HFC-365mfc cover a small share of the market whilst CO<sub>2</sub> and pentane are dominating.

Total HFCs emissions in this category were 1.73 Gg CO<sub>2</sub> eq. in 2023 ([Table 4.58](#)).

#### Methodological Issues

HFCs emissions from open cells are not occurring in Slovakia. For closed cells, the blowing agents remain longer in foam; the half life time is calculated with >20 years and depends on the panel's thickness. According to the IPCC 2006 GL, product life of the used cells should be 50 years except of injected foams where product life is 15 years. These values are used in the calculation of emissions estimates. Emissions estimates are calculated based on first-year emissions and annual losses as described in the IPCC 2006 GL (emissions from decommissioning do not occur in Slovakia, yet).

Bank of used HFCs is monitored since the first year of their use as follows: Bank in year t = Bank in year t-1 + New fillings in year t-1 – Emissions from new fillings in year t-1 – Emissions from bank in year t-1 – Decommissioned equipment in year t

The relationship for bank is different from that used for refrigeration category because no servicing occurs for foams.

Emission factors are based on the data provided by producers. First-year losses are assumed to be 10%, annual losses 0.5%. These values are the same or close to the default values according to the IPCC 2006 GL. Activity data were collected via the web reporting system as described in the [Annex 4.2](#) of this Document. Import-export of bulk chemicals and products were collected in order to obtain annual sales data, which were then assumed be equal to new fillings. It was decided to use conservative approach for the first-year emissions from all new fillings occurred in Slovakia. In 2014, the using of HFC-227ea was reported for the first time in Slovakia.

**Table 4.58: Aggregated data on HFCs using in the category 2.F.2 in particular years**

YEAR	New Fillings	Bank	Disposal	New Fillings			Bank	Disposal
				New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.								
1990	NO	NO	NO	NO	NO	NO	NO	NO
1995	NO	NO	NO	NO	NO	NO	NO	NO
2000	53.560	48.204	NO	5.356	0.241	NO	NO	5.597
2005	34.282	274.942	NO	3.428	1.375	NO	NO	4.803
2010	3.656	349.693	NO	0.366	1.748	NO	NO	2.114

YEAR	New Fillings	Bank	Disposal	New Fillings			Bank	Disposal
				New Fillings	Bank	Disposal		
	Gg CO <sub>2</sub> eq.							
2011	4.257	351.236	NO	0.426	1.756	NO	NO	2.182
2012	7.752	353.311	NO	0.775	1.767	NO	NO	2.542
2013	3.489	358.521	NO	0.349	1.793	NO	NO	2.142
2014	1.853	359.869	NO	0.185	1.799	NO	NO	1.985
2015	0.012	359.737	NO	0.001	1.799	NO	NO	1.800
2016	NO	357.949	NO	NO	1.790	NO	NO	1.790
2017	NO	356.159	NO	NO	1.781	NO	NO	1.781
2018	NO	354.378	NO	NO	1.772	NO	NO	1.772
2019	NO	352.606	NO	NO	1.763	NO	NO	1.763
2020	NO	350.843	NO	NO	1.754	NO	NO	1.754
2021	NO	349.089	NO	NO	1.745	NO	NO	1.745
2022	NO	347.344	NO	NO	1.737	NO	NO	1.737
2023	NO	345.607	NO	NO	1.728	NO	NO	1.728

### Uncertainties and Time-series Consistency

A consistent time series of HFCs import-export exists since the first years of HFCs using in foams and is well documented (the collection of data started in 1995 by using questionnaires – before the start of using of HFCs in foams). The same method was used for the whole time series. The decrease in emissions (and new fillings) in 2008 was caused by replacing of blowing agent R134a with water in new injected PU foams in 2007.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.12.11. Fire Protection (CRT 2.F.3)

This category is not significant and includes F-gases used in the following industry:

- HFC134a used as fluid in operating systems since 1994 in very little amount;
- HFC227ea (*FM 200*) is used as extinguishing media and suitable alternative for halon H1301 in fixed extinguishing systems since 2004. After 1993, halons are not imported into Slovakia;
- HFC 236fa (*FE36*) started to be used for portable extinguishing systems since the year 2000;
- PFCs extinguishing media are not imported into Slovakia. PFC 410 and PFC 614 have been never used in stable extinguishing equipment.

Prices of new extinguishing medias are quite high (approx. 40 Euro/kg), so the consumption and emissions are minimal. Stationary fire protection systems for flooding indoor spaces mainly use inert gases at present. Formerly used ozone layer depleting halons have been replaced in some cases by HFCs. HFC-227ea in the fire extinguishers was firstly introduced on the Slovak market in 1994. F-gases for firefighting are imported in cylinders and filled in fixed installed systems.

Total HFCs emissions in this category were 27.20 Gg CO<sub>2</sub> eq. in 2023.

## Methodological Issues

Annual sales of single HFC gases are calculated based on import – export of bulk chemicals and products. Detailed data on consumption for new equipment, the stock in existing fixed flooding systems, annual losses (refilling) and recovered F-gases from disposal were obtained directly from the fire protection companies. Stable extinguishing systems (flooding a streaming systems) used to protect electronic equipment have pressure vessels with lifetime from 10-12 years (given by the producer). After this time, extinguishing media are recovered, recycled and used again. In systems with working pressure 25 or 40 bar, the lifetime of pressure vessels is supposed to be at least up to 25 years.

HFC emissions occur from filling in fixed systems, from the bank (*in case of false alarm, fire, leakage, accidents etc.*) and from disposal. Test flooding, in former times an important source of emissions, have not taken place since 2000. The product manufacturing emission factor for filling of fixed systems is 1%. The emissions from bank are equalized with the company reports for refilling of losses. The product life factor from bank is 5% based on this assumption. Both factors were consulted with the fire protection companies and are in agreement with references. Used product life factor was used as a country specific one and it is slightly higher than the default value provided in the IPCC 2006 GL for installed flooding systems (1-3% per year). Emissions from disposal are reported since 2016.

Activity data were collected via web reporting system as described the [Annex 4.2](#) of this Document. Import-export of bulk chemicals and products data were collected in order to obtain annual sales data (which are equal to new fillings + service). Detailed data on consumption for the new equipment, the stock in existing fixed flooding systems, annual losses (refilling) and recovered F-gases from disposal were obtained directly from the fire protection companies and the Association of the Fire Extinguishers Producers in the Slovak Republic. These data served as tools for differentiating of annual sales data into new fillings and operational emissions (from bank).

**Table 4.59: Aggregated data on HFCs used in the category 2.F.3 in particular years**

YEAR	New Fillings	Bank	Disposal	New Fillings	Bank			Disposal	New Fillings
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	33.812	33.668	36.823	NO	0.338	1.841	NO	NO	2.179
2000	31.270	31.152	153.507	NO	0.313	7.675	NO	NO	7.988
2005	19.098	19.034	260.099	NO	0.191	13.005	NO	NO	13.196
2010	15.527	15.152	335.054	NO	0.155	16.753	NO	NO	16.908
2011	41.793	41.548	359.694	NO	0.418	17.985	NO	NO	18.403
2012	10.737	10.241	351.533	NO	0.107	17.577	NO	NO	17.684
2013	12.047	11.565	345.414	NO	0.120	17.271	NO	NO	17.391
2014	21.318	22.274	350.296	NO	0.213	17.515	NO	NO	17.728
2015	39.530	48.071	380.639	NO	0.395	19.032	NO	NO	19.427
2016	27.211	27.084	379.918	6.702	0.272	18.996	2.011	4.692	21.279
2017	31.536	16.428	367.488	7.672	0.315	18.374	2.302	5.370	20.991
2018	20.062	12.044	351.243	7.680	0.201	17.562	2.378	5.302	20.141
2019	34.131	28.065	340.927	16.495	0.341	17.046	4.767	11.728	22.155
2020	24.564	8.086	324.752	5.499	0.246	16.238	5.063	0.436	21.546
2021	27.038	5.425	309.165	3.623	0.270	15.458	3.402	0.221	19.130
2022	22.708	7.610	290.536	8.408	0.227	14.527	7.177	1.231	21.931
2023	22.323	6.880	259.945	18.174	0.223	12.997	13.984	4.190	27.204



---

## Uncertainties and Time-series Consistency

A consistent time series of HFCs import-export data exists since 1995 and is well documented by using questionnaires. The same method was used for the whole time series. The increasing trend (since 1994) in actual emissions from HFC-227ea and HFC-236fa was stabilized and emissions are approximately at the same level. The purchase of new fire extinguishers depends mostly on the building of new server rooms.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

### 4.12.12. Aerosols (CRT 2.F.4)

The producers of aerosols in Slovakia changed directly from ODS to mechanical principles and use of hydrocarbons and dimethyl ether in 1990. The group of aerosols gases includes medical aerosols, i.e. Metered Dose Inhalers (MDIs), only. The HFC-134a and HFC227ea are used as propellant for such aerosols in Slovakia. However, since 2015, HFC-134a occurs only in Slovakia. Total HFCs emissions in this category are not significant and were 10.04 Gg of CO<sub>2</sub> eq. in 2023. The production of MDI does not occur in Slovakia.

#### Methodological Issues

Aerosol emissions are considered prompt because all the initial charge escapes within the first two years after manufacture, typically six months after sale for most sub-applications (due to the short expiration time). It is assumed that the initial charge escapes approximately during two years. Therefore, the total amount of aerosol initially charged in product containers prior to sale in actual year and year before were taken into consideration in emissions estimation.

The production of MDI does not occur in Slovakia. The initial charge (new fillings, import to Slovakia) escapes during two years. This calculation of emission estimates corresponds to the equation:

$$\text{Emissions}_{\text{in year } t} = \text{Initial charge}_{\text{in year } t-1} * (1-EF) + \text{Initial charge}_{\text{in year } t} * EF$$

In a similar way a bank of chemicals is calculated:

$$\text{Bank}_{\text{in year } t} = \text{Initial charge}_{\text{in year } t-1} * (1-EF) + \text{Initial charge}_{\text{in year } t} * (1-EF)$$

EF is the same in both equation and equals to 0.5.

The basic philosophy of the calculation of bank is that the bank refers to the amount of gas that is not released as an emission in the previous and current year. In order to increase transparency, the numerical exercise is provided.

The content of HFC-134a in sold MDI in 2017 and 2018 were 6.175 t and 6.292 t, respectively. For emission calculation in 2018 the following way is used:

1. Due the fact that EF=0.5, the half of the amount sold in 2017 was used in 2017 and this amount is not of interest for 2018 calculation.
2. The rest of the gas sold and not used in 2017 was moved to bank of chemicals (3.087 t).
3. Emission calculation for 2018: the term  $\text{Initial charge}_{\text{in year } t-1} * (1-EF)$  in the equation represents the gas that was moved to the bank in 2017 (3.087 t). The term  $\text{Initial charge}_{\text{in year } t} * EF$  in the equation represents the gas that was used in 2018 (the half of the gas sold in 2018: 3.146 t).

4. Bank calculation for 2018: the term  $Initial\ charge_{in\ year\ t-1} * (1-EF)$  in the equation represents the gas that was moved to the bank in 2017 (3.087 t). The rest of the gas that was sold in 2018 (and not used this year) is also added to the bank (3.146 t) and will be used for emission calculation in 2019.
5. It should be noted that the same numbers for emissions and bank are due the fact that  $EF=0.5$ . E.g. if we assume that  $EF=0.6$  the values for emissions and bank will not be the same.

The State Institute for Drug Control of Slovakia is in the position to provide activity relevant data for emissions estimation. The activity data represents the number of containers with aerosols imported to Slovakia. The State Institute for Drug Control (ŠÚKL) provided this data on behalf of the Act No 286/2010 Coll. Data are available since 2000. Based on the statement of the ŠÚKL experts, no MDIs had been imported to Slovakia before the year 2000.

**Table 4.60: Aggregated data on HFCs using in the category 2.F.4 in particular years**

YEAR	Filled Into New Products	Bank	Emissions From:		Total Emissions
			New Fillings	Bank	
<i>Gg CO<sub>2</sub> eq.</i>					
1990	NO	NO	NO	NO	NO
1995	NO	NO	NO	NO	NO
2000	NO	2.425	NO	2.425	2.425
2005	NO	6.182	NO	6.182	6.182
2010	NO	7.116	NO	7.116	7.116
2011	NO	7.624	NO	7.624	7.624
2012	NO	7.704	NO	7.704	7.704
2013	NO	8.098	NO	8.098	8.098
2014	NO	8.406	NO	8.406	8.406
2015	NO	9.058	NO	9.058	9.058
2016	NO	9.253	NO	9.253	9.253
2017	NO	8.325	NO	8.325	8.325
2018	NO	8.103	NO	8.103	8.103
2019	NO	8.310	NO	8.310	8.310
2020	NO	8.303	NO	8.303	8.303
2021	NO	8.832	NO	8.832	8.832
2022	NO	9.852	NO	9.852	9.852
2023	NO	10.038	NO	10.038	10.038

### Uncertainties and Time-series Consistency

A consistent time series of HFCs import-export data exists since the first years of MDIs use (2000) and is well documented. The same method for emissions estimation is used for the whole time series. HFC-134a is used since 2000. MDIs containing HFC-227ea are imported into Slovakia since 2008 and ended in 2015.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### 4.12.13. Solvents (CRT 2.F.5)

The HFCs emissions are not occurring in this category, recently. There is no import of F-solvents to Slovakia because they are rather expensive. SP-255, which contains distilled oil and methyl acetate, is used as a flushing material. The solvents L113 and S316 used in Slovakia are not obliged to include in the emissions inventory. The solvents with HFCs are not used in cleaning machines for flushing refrigeration circuits. The emissions of PFC14 (CF<sub>4</sub>) in solvents were estimated for the years 1997 – 2006 and then PFC14 was replaced with the SF<sub>6</sub> gas. No PFCs or SF<sub>6</sub> emissions were occurring in this category in 2022. Used amount of SF<sub>6</sub> is negligible (below 0.2 t/year). In the production process, the SF<sub>6</sub> is used for Si wafers etching after previous operation (Si wafers cutting on chips). Technological process can be described as follows:

- Si wafers are put into chamber of plasma equipment and after that air is exhausted from chamber for required vacuum,
- etching process starts with high-frequency burning SF<sub>6</sub> what cause etching of Si wafers surface,
- SF<sub>6</sub> and remains after etching process are exhausted from plasma equipment,
- these by-products go into special washing tank with NaOH where HF is neutralized.

According to the measuring of the semiconductor producer Semicron Vrbové, SF<sub>6</sub> emissions during etching are not emitted into atmosphere. Therefore, notation key “NO” is used for time series. PFC14 emissions from the solvents use are reported for the period 1997 – 2006.

**Table 4.61: PFC14 emissions in the category 2.F.5 in 1997 – 2006**

YEAR	Filled Into New Products	Bank	Emissions From:		Total Emissions
			New Fillings	Bank	
<i>Gg CO<sub>2</sub> eq.</i>					
1997	NO	0.610	NO	0.610	0.610
1998	NO	2.021	NO	2.021	2.021
1999	NO	2.563	NO	2.563	2.563
2000	NO	1.274	NO	1.274	1.274
2001	NO	2.244	NO	2.244	2.244
2002	NO	3.315	NO	3.315	3.315
2003	NO	1.591	NO	1.591	1.591
2004	NO	0.696	NO	0.696	0.696
2005	NO	0.398	NO	0.398	0.398
2006	NO	0.099	NO	0.099	0.099

Emissions are considered prompt. It was considered that the new fillings escape during two years. Therefore, the total amount of PFC114 used in actual year and year before was used for emissions estimation. Due to this fact (the rest of the previous year’s new fillings has to escape in the next year), the emission factor from bank is 100% (the bank is calculated in the same way as described in the Chapter 4.12.12). The emission calculation corresponds to the equation:

$$\text{Emissions}_{\text{in year } t} = \text{New fillings}_{\text{in year } t-1} * (1-\text{EF}) + \text{New fillings}_{\text{in year } t} * \text{EF}, \text{ where EF}=0.5.$$

#### 4.12.14. Other Applications (CRT 2.F.6)

Emissions in this category are not occurring for the time series 1990 – 2023.

## 4.13. Other Product Manufacture (CRT 2.G)


### 4.13.1. Source Category Description

Emissions of SF<sub>6</sub> from the high voltage switchgears and emissions of N<sub>2</sub>O from use for anaesthesia and in food industry (aerosol cans) are reported in this category. Total emissions in CO<sub>2</sub> eq. were 71.41 Gg in 2023, decreased by 1% in comparison with the previous year. The decrease is caused by decreased service emissions of electrical equipment. Comparing with the base year, the increase is nearly 500%. This increase is mostly caused by the increase in N<sub>2</sub>O emissions from aerosol cans. Emissions from SF<sub>6</sub> from other product use (2.G.2) are included in 2.G.1 electrical equipment.

**Table 4.62:** Emissions in the category 2.G according to the subcategories in particular years


YEAR	2.G.1 Electrical Equipment	2.G.2 SF <sub>6</sub> and PFCs from Other Product Use	2.G.3 N <sub>2</sub> O from Product Use
	Gg of CO <sub>2</sub> eq.		
1990	0.06	IE	14.58
1995	10.47	IE	26.50
2000	13.44	IE	17.21
2005	16.89	IE	73.82
2010	20.23	IE	69.06
2011	21.44	IE	64.84
2012	21.90	IE	78.04
2013	22.99	IE	108.69
2014	14.60	IE	71.17
2015	14.75	IE	60.57
2016	6.00	IE	61.75
2017	7.30	IE	61.96
2018	9.68	IE	61.80
2019	9.14	IE	58.54
2020	17.73	IE	58.01
2021	17.44	IE	57.48
2022	15.38	IE	57.00
2023	14.70	IE	56.71

**Figure 4.15:** The trend of individual subcategories in the category 2.G in 1990 – 2023



The major share (79.4%) in emissions belongs to the N<sub>2</sub>O emissions from the product use, 20.6% belongs to SF<sub>6</sub> emissions from electrical equipment.

**Figure 4.16:** The share in GHG emissions on individual categories of the 2.G in 2023



#### 4.13.2. Electrical Equipment (CRT 2.G.1)

Emissions of SF<sub>6</sub> from the thermal insulation of windows and from the high voltage switchgears are reported in this category. The Nitrasklo Ltd. company for windows used SF<sub>6</sub> since 1994 for anti-noise and thermal isolation. It was mixed with argon in the rate 30:70. Due the more effective production, consumption decreased. It was filled in close cycles without emissions from production. Consumption of SF<sub>6</sub> in Nitrasklo Ltd. continually decreased and it was phased out in the year 2002. Amount of stored gas annually in windows in the Slovak Republic was 10 kg from 80 kg filled into windows annually (70 kg were exported in windows). For the stock of gas remaining inside, an annual leakage rate is 1%. SF<sub>6</sub> emissions from window insulation are very negligible when compared to the emissions from electrical equipment (approx. 0.09% of total SF<sub>6</sub> emissions. Since the production of windows stopped in 2002, we considered it unfeasible to report disaggregated emissions. Data on windows are reported together with the emissions from isolating gas in high voltage switchers.

Most of the SF<sub>6</sub> is used as insulation media in high and low voltage electric equipment because of higher safety level and enable to reduce dimension of equipment. SF<sub>6</sub> is used as an arc quenching and insulating gas in high-voltage (>36 kV [110–380 kV]) and medium-voltage (1–36 kV) switchgear and control gear. The equipment – mainly Gas-Insulates Systems, GIS – has not been manufactured during the report period in Slovakia, but has been completely imported. High-voltage GIS (HV GIS) operate with a high operating pressure (up to seven bar) and large gas quantities. They are imported with a transport filling and are filled up on site. The systems are “closed for life” and have to be replenished in their lifetime. Emissions from operating HV systems are higher than emissions from the medium-voltage GIS (MV GIS). These operate with lower overpressure and small gas quantities of only some kg per system. They are already charged with SF<sub>6</sub> when imported and are hermetically closed (“sealed for life”).

Total actual emissions of SF<sub>6</sub> were 14.70 Gg CO<sub>2</sub> eq. (0.625 t SF<sub>6</sub>) in 2023 ([Table 4.63](#)). In 2013, old equipment started to be disposed. Servicing of the electrical equipment was lower than in previous years, therefore the operational emissions decreased. It was verified by top-down approach (balance of annual sales etc. of SF<sub>6</sub>).

**Table 4.63:** SF<sub>6</sub> emissions in the category 2.G.1 in particular years

YEAR	New Fillings	New Addition to Bank	Bank	Retired Equip.	Emissions from:			Recovery	Total
					New Fillings	Bank	Disposal		
Gg CO <sub>2</sub> eq.									
1990	3.008	3.008	3.008	NO	0.030	0.030	NO	NO	0.060
1995	72.122	63.008	974.526	NO	0.721	9.745	NO	NO	10.466
2000	54.779	42.307	1 289.668	NO	0.548	12.897	NO	NO	13.444
2005	88.595	73.329	1 600.124	NO	0.886	16.001	NO	NO	16.887
2010	69.584	50.520	1 957.101	NO	0.696	19.531	NO	NO	20.227
2011	102.319	82.751	2 039.852	NO	1.023	20.417	NO	NO	21.440
2012	85.164	64.768	2 104.620	NO	0.852	21.044	NO	NO	21.896
2013	64.390	49.350	2 143.160	10.810	0.644	22.161	0.184	10.626	22.988
2014	47.215	62.731	2 068.376	137.515	0.472	11.793	2.338	135.177	14.603
2015	121.035	152.786	2 210.024	11.138	1.210	13.354	0.189	10.949	14.753
2016	6.705	165.196	2 357.065	18.154	0.067	5.621	0.309	17.846	5.997
2017	17.631	86.571	2 433.818	9.818	0.176	6.957	0.167	9.651	7.300
2018	40.361	74.491	2 485.782	22.527	0.404	8.883	0.394	22.132	9.681
2019	9.338	46.841	2 511.102	21.521	0.093	8.680	0.364	21.157	9.137
2020	4.196	38.688	2 536.423	13.367	0.042	17.455	0.233	13.134	17.729
2021	NO	47.888	2 575.092	9.219	NO	17.283	0.155	9.064	17.438
2022	14.558	22.674	2 568.187	29.579	0.146	14.707	0.524	29.055	15.377
2023	16.450	297.552	2 823.573	42.166	0.165	13.802	0.728	41.438	14.695

### Methodological Issues

The IPCC 2006 GL describe two general approaches for estimating emissions, which occur during the year mass-balance (top-down) and emission-factor (bottom-up) approach, respectively. The bottom-up approach takes into account the time lag between consumption and emissions explicitly by the emission factors. The top-down approach takes the time lag into account implicitly, by tracking the amount of virgin chemical consumed in a year that replaces emissions from the previous year.

The web reporting system allows calculating emissions in both approaches. The bottom-up approach in a combination with the top-down approach was used. The procedure is as follows:

1. Using the bottom-up approach from the Logbook Leaklog (described in the [Annex 4.2](#));
2. Calculation of the total consumption of SF<sub>6</sub> in Slovakia based on the Leaklog;
3. Calculation of the total consumption of SF<sub>6</sub> in Slovakia according to the top-down approach;
4. Comparison of calculated results by different approaches;
5. If differences occur, the data in bottom-up approach are corrected by correction of operational emissions (no correction was necessary in 2022);
6. Calculation of emission estimates by the bottom-up approach using the corrected data.

For the top-down approach, the following formula based on the structure of the reporting systems was used:

$$\text{Emissions} = \text{Annual sales of SF}_6 - \text{Total charge of new equipment} + \text{Disposal emissions}$$

where: *Annual sales* and *Total charge of new equipment* are calculated by formulas presented in the IPCC 2006 Guidelines, Chapter 3, p. 7.54 (not simplified formulas). This formula corresponds to the formula described in Chapter 3, p. 8.14.

For bottom-up approach, the following formulas are used:

$$\text{Emissions} = \text{Emissions from new fillings} + \text{Operational emissions} + \text{Disposal emissions}$$

---

where: *Emissions from new fillings* represent 1% (EF) of the new charges filled in Slovakia (*SF<sub>6</sub> to Charge domestically manufactured and Assembled equipment + SF<sub>6</sub> to Charge equipment that is not Factory-Charged*).

*Operational emissions* represent the consumption of gases for servicing (these data are reported in Leaklog). It is assumed that the SF<sub>6</sub> used for servicing restocks the emissions from the bank and thus the bank of the chemical remains constant.

*Disposal emissions* represent the emissions from the retired equipment.

For the consistency of operational emissions, the bank of SF<sub>6</sub> is necessary to follow. The bank is calculated as follows:

$\text{Bank}_{\text{in year } t} = \text{Bank}_{\text{in year } t-1} + \text{New additions to bank} - \text{SF}_6 \text{ in retired equipment}$

where: *New additions to bank = SF<sub>6</sub> to Charge Domestically Manufactured and Assembled Equipment + SF<sub>6</sub> to Charge Equipment that is not Factory-Charged + SF<sub>6</sub> Contained in Imported Equipment Already Charged – SF<sub>6</sub> Contained in Exported Equipment Already Charged.*

Emission factors from the filling SF<sub>6</sub> into new equipment (product manufacturing factor) is assumed 1% (based on the producers' data). Operational emission factor (product life factor) is calculated yearly based on the reported operational emissions and the respective bank.

Disposal emission factor (disposal loss factor) is based on the survey of recycling factories. It follows that 98.3% of SF<sub>6</sub> is recovered for repeated used or destroyed (in 2023, 0.421 t was destroyed). Thus, the disposal loss factor is 1.7%. The activity data are collected together with the other F-gases data as described in the category 2.F and in the [Annex 4.2](#) of this Document. Amount of SF<sub>6</sub> in disposed systems was taken directly from recycling factories.

#### **Uncertainties and Time-series Consistency**

A consistent time series of SF<sub>6</sub> import-export data exists since 1993 and is well documented. Data were collected in questionnaires. In previous submissions, the data on banks were not consistent because of use of extrapolation in 2010. The inconsistency was caused by two types of formula used for bank calculation. In the 2015 submission, the inconsistent bank data were corrected, the bank data were recalculated by the same formula (as presented above) in the whole time series. Product life factor for the time series 1990 – 2009 was assumed average of product life factors of 2010 – 2014 (1%). Product life factor is higher than the default value (0.2%) provided in the IPCC 2006 GL.

In 1994, the owner of the Slovak electrical power system began with the modernization of grids and transformer stations. Therefore, the sharp increase in SF<sub>6</sub> emissions is visible in 1994.

As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.

#### **4.13.3. Use of SF<sub>6</sub> and PFCs in Other Products (CRT 2.G.2)**

SF<sub>6</sub> can be used as an extinguishing medium in electronics, protection against explosion, isolation, sterilization, detection gas, alloying of Al and Mg and in tobacco production. SF<sub>6</sub> gas is rather expensive and therefore it was never used as an extinguishing medium in industry in Slovakia. Shoes and tires with F-gas cushions are not manufactured or imported to Slovakia for the time series 1990 – 2023. Emissions from in windows insulation are reported in 2.G.1.

#### 4.13.4. N<sub>2</sub>O from Product Uses (CRT 2.G.3)

Medicine (anaesthesia) and food industry (aerosol cans) N<sub>2</sub>O emissions are reported in this category in 2020. There is also the consumption of N<sub>2</sub>O for analytical purposes, but the gas is burned after use, so this source is not included into inventory. Total N<sub>2</sub>O emissions from aerosol cans were 194.6 t and total N<sub>2</sub>O emissions from anaesthesia were 19.4 t in 2023.

##### Methodological issues

The methodology is based on the default tier 1 due to less significant of this category (it is not a key category). The final N<sub>2</sub>O emissions from these sources are equal to the consumed gas in medicine and food industry in the reporting year. The time series was reconstructed based on the statistical data on production. The N<sub>2</sub>O emissions according to the categories are summarized in [Table 4.64](#).

**Table 4.64:** N<sub>2</sub>O emissions from product use in particular years

YEAR	Total N <sub>2</sub> O		
	2.G	2.G.3.a Medical Application (Anaesthesia)	2.G.3.b Other (Aerosol Cans)
	Gg		
1990	0.0550	0.0550	NO
1995	0.1000	0.1000	NO
2000	0.0650	0.0650	NO
2005	0.2785	0.0656	0.2129
2010	0.2606	0.0528	0.2078
2011	0.2447	0.0490	0.1957
2012	0.2945	0.0445	0.2500
2013	0.4102	0.0190	0.3912
2014	0.2686	0.0176	0.2510
2015	0.2285	0.0275	0.2010
2016	0.2330	0.0190	0.2140
2017	0.2338	0.0178	0.2160
2018	0.2332	0.0182	0.2150
2019	0.2209	0.0186	0.2023
2020	0.2189	0.0198	0.1991
2021	0.2169	0.0196	0.1973
2022	0.2151	0.0195	0.1956
2023	0.2140	0.0194	0.1946

Used N<sub>2</sub>O EFs in medicine and food industry are based on approximation, that emissions are equal to consumed gas (EF = 1 t/t). It is assumed that all gas is evaporated into the atmosphere in the reporting year. This assumption is in line with the IPCC 2006 GL for medical applications and aerosol cans in food industry. The activity data in the category 2.G.3 come from the distributors of N<sub>2</sub>O liquid gas – Messer-Tatragas, Linde, Air Products and SIAD companies. The disaggregation of gas utilization is based on direct information from the gas distributors.

##### Uncertainties and Time-series Consistency

Consistent methodology and tier method were used for the whole time series. As described in the [Chapter 4.2](#), tier 2 approach to the uncertainty analysis of the subcategories was chosen. The uncertainty analysis is presented in the SVK NID 2024. This approach has been chosen because no changes of uncertainties in IPPU subcategories and no changes in methodology of any subcategory have been made. The results of the uncertainty analysis of the subcategories (probability density function, percentiles, overall uncertainty) are nearly the same as in the previous submission. Therefore, the calculation of the statistical characteristics in the subcategories will be made every fifth year since this submission or it has been made in the case when the methodology or uncertainties of any used data is changed.



---

## 4.14. Other Production (CRT 2.H)


The NMVOC emissions mainly from food industry were reported in this category in 2023. Total emissions of NMVOC were 3 336 t and are consistent with the CLRTAP inventory. No GHG emissions occurred in the time series 1990 – 2023.

# Annex 4.1. CO<sub>2</sub> Reference Approach and Comparison with the Sectoral Approach, and Relevant Information on the National Energy Balance

## A4.1.1 Methodology for Carbon Balance of Iron and Steel Production

The country specific methodology is implemented in the inventory (see Chapter 4.9.1 of this Document). Pig iron and steel are produced in iron and steel integrated plant and by the EAF method. Iron and steel integrated production is a complex with many energy-related installations (coke ovens, heating plant, etc.). Several available data for integrated iron and steel can be found in: (i) questionnaires provided by the producers (data on raw materials, pig iron and steel produced and limestone used); (ii) the NEIS database (detailed data on fuels used and their flows); (iii) the EU ETS reports (data on total carbon balance of all inputs and outputs). The EU ETS reports were used during QA/QC process to verify estimates. The allocation of sources into IPCC subcategories cannot be provided based on data available in the EU ETS reports. In order to prepare carbon balance, the simplified scheme of the plant was proposed (*Figure A4.1.1*). Occasional sale of produced pig iron was considered, too. In some cases, parts of coking gas and blast furnace gas were sold to the nearby brickyard plant, which was also considered during estimation. Total carbon balance was calculated according to the proposals depicted in the Scheme. All the streams were estimated using plant specific conversion units and carbon EFs taken from the category 1.A.2.a of the Energy sector or based on carbon content in materials.

*Figure A4.1.1: The simplified distribution scheme of the complex plant for pig iron and steel production*



Carbon balance consists of four steps: (1) balance of 2.C.1, (2) balance of 1.A.1.c, (3) balance of 1.A.2.a and (4) balance of 1.A.2.g.viii - Other.

**Table A4.1.1: Balance of the category 2.C.1 in 2023**

STREAM	AD	NCV	EF (C)	CARBON
	kt; mil. m <sup>3</sup>	TJ /m.u.	t/TJ; mass fraction	kt
Coking coal	2362.94	29.126	25.703	1768.96
Anthracite	0.00	0.000	0.000	0.00
Coke surplus	186.77	27.062	29.871	150.98
Natural gas	21.47	35.477	15.331	11.68
Tar and wastes	-1913.87	-	0.037	-70.43
Coking gas	-564.47	16.501	11.435	-106.51
Blast furnace gas	-3832.19	2.911	76.618	-854.71
Iron ore	6847.65	-	3.552E-03	24.32
Steel	-4074.96	-	8.900E-04	-3.63
Pig iron sold	-17.56	-	1.991E-03	-0.03
Limestone used	805.88	-	1.201E-01	96.78
TOTAL			1 017.40	

CO<sub>2</sub> emissions estimation in the 2.C.1 is based on the carbon balance (from that plant) and represents the value 3 730.20 Gg (total carbon × 44/12).

**Table A4.1.2: Balance of the category 1.A.1.c in 2023**

STREAM	AD	NCV	EF (C)	CARBON
	kt; mil. m <sup>3</sup>	TJ /m.u.	t/TJ; mass fraction	kt
Natural gas	2.049	35.477	15.33	1.11
Coking gas	122.30	16.50	11.44	23.08
Blast furnace gas	1402.08	2.91	76.62	312.71
TOTAL			336.90	

CO<sub>2</sub> emissions estimation in 1.A.1.c is based on the carbon balance (from that plant, not total 1.A.1.c) and represents the value 1 235.31 Gg (total carbon × 44/12).

**Table A4.1.3: Balance of the category 1.A.2.a in 2023**

STREAM	AD	NCV	EF (C)	CARBON
	kt; mil. m <sup>3</sup>	TJ /m.u.	t/TJ; mass fraction	kt
Other bituminous coal	194.16	28.206	25.385	139.02
Natural gas	11.75	35.477	15.331	6.39
Coking gas	236.37	16.501	11.435	44.60
Blast furnace gas	2201.35	2.911	76.618	490.98
TOTAL			680.99	

CO<sub>2</sub> emissions estimation in 1.A.2.a is based on the carbon balance (from that plant, not total 1.A.2.a) and represents the value 2.496.97 Gg (total carbon × 44/12).

**Table A4.1.4: Balance of 1.A.2.g.viii – Other in 2023**

STREAM	AD	NCV	EF (C)	CARBON
	kt; mil. m <sup>3</sup>	TJ /m.u.	t/TJ; mass fraction	kt
Natural gas	88.00	35.477	15.331	47.86
Coking gas	205.56	16.501	11.435	38.79
Blast furnace gas	228.76	2.911	76.618	51.02
TOTAL			137.67	

---

CO<sub>2</sub> emissions estimation in 1.A.2.g.viii - Other is based on the carbon balance (from that plant, not total 1.A.2.g.viii - Other) and represents the value 504.79 Gg (total carbon × 44/12).

The output from the plant was 0.242 mil. m<sup>3</sup> of coking gas and 0 mil. m<sup>3</sup> of blast furnace gas in 2023. In the years, when output is reported from the iron and steel plant, it means, that gases are sold to nearby brickyard and they are balanced in the category 1.A.2.g.viii - Other.

Carbon balance presented in this Annex is only from the integrated iron and steel plant. The CO<sub>2</sub> emissions estimation presented here is allocated in the categories 2.C.1, 1.A.1.c, 1.A.2.a and 1.A.2.g.viii - Other. The presented Energy sector includes also other productions or technologies in Slovakia. Therefore, total CO<sub>2</sub> emissions calculated via this approach will be lower than those presented in each individual CRT table. In comparison with the verified CO<sub>2</sub> emissions under the EU ETS, the emissions estimated for the integrated iron and steel plant by using this country specific input-output approach differ by 0.10%: (i) NID: 7 972.83 Gg CO<sub>2</sub>; (ii) EU ETS: 7 964.71 Gg CO<sub>2</sub>. It should be noted that in both values compared the CO<sub>2</sub> from desulphurization and DENOX applications are included (5.07 and 0.48 Gg CO<sub>2</sub>, respectively).

---

## Annex 4.2. Methodology of Acquisition and Data Processing on F-gases Consumption in the Categories 2.F, 2.G.1 and 2.G.2









Fluorinated greenhouse gases (F-gases) are used in numerous applications and include three types of gases: HFCs, PFCs and SF<sub>6</sub>. F-gases emissions are mainly released from refrigeration and air conditioning equipment, foams, aerosols, solvents, fire protection equipment, from halocarbon production, from certain industrial processes in semiconductor and non-ferrous metal industry and from equipment for transmission of electricity during manufacture, use and at disposal.

Due to their relatively high global warming potentials, F-gases are addressed by international conventions such as the United Nations Framework Convention on Climate Change (UNFCCC) and its Kyoto Protocol (with post-2012 amendment), as well as by policies at the European and at national levels. The EU committed to reduce overall greenhouse gas emissions by 20% compared to the base year 1990 during the second commitment period 2013 – 2020.

The EU policy targets are based on further reduction of halocarbon refrigerant usage, on the substantially decreased leakage percentage and energetically efficient operation of air conditioning systems, heat pumps and refrigeration installations. Success of the EU Regulation No. 517/2014 depends on effective measures. This new regulation, which replaces the first EU Regulation No. 842/2006 was applied from 1<sup>st</sup> January 2015, strengthens the existing measures and introduces a number of far-reaching changes. By 2030, it will cut the EU's F-gas emissions by two-thirds compared with 2014 levels. Described solutions are based on data recorded in the log-book according to EN 378 Regulation (EC) No 1516/2007. Advantages of electronic data logging and reporting are shown on the possibilities of automatic analysis, fault detection and comparison of, fast access to the full history of leak checks and various forms of output. Value added of electronic logbook is indirect detection of refrigerant leak. The fault detection classifier estimates the probability of refrigerant leak.

In the year 2003, Slovakia started software with access to the processing and data assessment. This software system is based on the activities of the Slovak Association for Cooling and Air-conditioning Technology (SZCHKT). The electronically led documentations have been developed from the previous paper questionnaires. Evaluated data were collected from the service organizations and customers. The backward running contact with inventoried companies enabled cooperation that is more effective. The companies can find their data reported in the previous years in the questionnaires. It enables the mutual control of the used data. Next step was data processing in Access database.


*Database of original data was processed in following tables:*

 01 Adresy organizacii s pohybom latok	01 Addresses of companies with move of substances
 02 Kody druhu importu a exportu latok	02 Code of the type of import and export
 03 Latky HFC SF6 PFC	03 Substances
 04 Zlozky zmesi latok	04 Components of the substances (mixtures)
 05 Druh latky	05 Type of substance
 06 Emisne koeficienty podla pouzitia latky	06 Emission factors
 07 Roky	07 Inventory years
 08 Pohyby latok za rok	

Database was prepared for processing according to the suggested algorithm. This way of data reporting was the only one used up to the end of the year 2009. In 2009, a new internet reporting system Leaklog started. This system is based on the activities of the Slovak Association for Cooling and Air-conditioning Technology (SZCHKT) and is available. The SZCHKT is the “Notified Body”, the body officially

authorized by the Ministry of Environment to certified companies and organizations for the activities in this area. Evaluated data are collected from the service organizations.

**Figure A4.2.1: System of data transfer from customer to Notified Body**



The reporting and data processing system consists of:

- Annual reporting of F-gases (new charges and leakages) by certified companies;
- Annual reporting of F-gases imported in bulks by certified companies;
- Annual reporting of F-gases in products by importers, exporters, producers by companies.


All companies dealing with the F-gases have access to the electronic system based on certificate provided by the Notified Body. Advantages of electronic data logging and reporting are in the possibilities of automatic analysis, fault detection and comparison of, fast access to the full history of leak checks and various forms of output. Service engineers get quick survey of the customers, cooling circuits, details of all maintenance work and repairs, refrigerants in store, refrigerants added, recovered, reclaimed, and disposed of. Value added of electronic logbook is indirect detection of refrigerant leak. The fault detection classifier estimates the probability of refrigerant leak. Electronic way of the data records enables summarizing, reporting and analysing important data in a chosen period in connection with the internet (**Figure A4.2.2**). Documented, consistent time series of HFCs import-export data exists since 1995. They were collected using the questionnaires (more than 250 companies). The institutions included in data collection are:

- Refrigerants, air-conditioning, heat pumps: SZCHKT. This institution is appointed for personnel and company certification required by 842/2006/EC. This certification activity was started by the Slovak association for cooling and AC Technology (SZCHKT) in the year 2009;
- Firefighting: Association of extinguishing appliances producers (ZVHP);
- MDI: State Institute for Drug Control (SUKL);
- Mobile AC: Automotive Industry Association (ZAP);
- Solvents: (SZCHKT);
- SF<sub>6</sub> use: (SZCHKT).

Reports to the database in subcategories refrigeration, air-conditioning and heat pumps, solvents and SF<sub>6</sub> include two web systems:

- import, export, sales data of bulk chemicals and products (database used since 2003),
- data on type of use (for new equipment or for recharge/service, recovery, reclaimed, disposal)

**Figure A4.2.2: Diagram of data flow in reporting (data flow direction: top to down)**



### A4.2.1 Reporting of F-gases Imported in Bulks

Refrigerant movements reporting is required according to EU legislation. Every certified company shall restore its certificate annually. Company has to enter website of the Notified Body with its name and password. Table on [Figure A4.2.3](#) is showing front-pages which appear after the signing up of company to the system. In this table, the certified company has to declare the competencies of the employees, possession of technical equipment, regular checking of electronic detectors, and movement of

refrigerants from the previous year. The confirmed data are saved and sent to the Notified Body until the end of January annually. After receiving the report, the Notified Body will restore the certificate. Certified companies and competent persons are listed on the website of the Notified Body. This is a part of the web system used since 2003.

**Figure A4.2.3:** Declaration of certified company with the legal status in the EU and in Slovakia about competencies of the employees, technical equipment, regular checking of electronic detectors, and refrigerant management categorized by field of application on the website of notified body

Kategória certifikátu: **I, MobKlim**

**Technické prostriedky a vybavenie**

Druh	Počet
Odberové zariadenie:	3
Zberné nádoby na zhodnotenie chladiva:	2
Elektronický detektor úniku chladiva s citlivosťou do 5g/rok:	2
Dvojstupňové vákuové čerpadlo:	2
Manometrický mostík:	2
Digitálna váha:	2
Nástroje bežne potrebné na odborný výkon servisnej činnosti:	2

Bola vykonaná kontrola funkcie elektronického detektora(ov) v predchádzajúcom roku podľa Nariadenia (ES) 1516/2007 § 6(2)

**Zamestnanci**

Číslo osvedčenia o odbornej spôsobilosti	Kvalifikácia	Meno	Priezvisko
3308	MXXXX	Ondrej	Fegyveres
2776	AAXXX	Michal	Feketevízi

+ Pridať zamestnanca

Hľadať číslo osvedčenia podľa priezviska

**Nakladanie s fluórovanými skleníkovými plynmi v roku 2012**

Servisná organizácia  
  Dovožca/vývozca  
  Servisná organizácia a zároveň dovozca/vývozca

**Hodnoty uvádzajte v kilogramoch.**

**Predané nové/zhodnotené: len inej certifikovanej organizácii!**

F plyn	Dovoz nové	Dovoz zhodnotené	Vývoz nové	Vývoz zhodnotené	Kúpené v SR nové	Kúpené v SR zhodnotené	Predané v SR nové	Predané v SR zhodnotené	Regenerované	Zničené	Únik nové	Únik zhodnotené
R404A	0.00	0.00	0.00	0.00	0.00	0.00	10.00	0.00	0.00	0.00	0.00	0.00
R134a												

+ Pridať riadok

**Použitie fluórovaných skleníkových plynov v roku 2012**

**Hodnoty uvádzajte v kilogramoch.**

F plyn	Doplnené nová náplň	Doplnené únik	Zhodnotené
R404A – Komerčné chladenie	0.00	0.00	0.00
R134a – Priemyselné chladenie			

+ Pridať riadok

Uložiť

Reporting of refrigerant movements in products is required according legislation. Every importer, producer or exporter shall report annually. Company has to enter the website of notified body with its name and password.

Figure A4.2.4 presents table of data reporting for products, which will be shown to the company after entering its account and this shall be filled in. In this table, the company has to report movements of



refrigerant in products from the previous year. The confirmed data are saved and shall be sent to the Notified Body until the end of January. After receiving the report, data are automatically processed. Reporting companies are listed on the website. All reported data are available for the reporting organizations. Historical development in all monitored refrigerants with emission projections up to 2025 are part of the web system since 2003.

Figure A4.2.4: Data reporting of importers, producers and exporters on products used

**Data reporting for 2012**  
Production, import and export of products

[Return without saving](#)

Product	Refrigerant / extinguishing medium	Charge (kg/pc)	Imported (pcs)	Imported from	Exported (pcs)	Exported to	Produced (pcs)
Aerosols	R227ea						
Air conditioi	R404A						
PUR insulati	R134a						
MobKlim	R134a						
Commercial	R407C						
Transport re	R404A						
Heat pumps	R407C						
SF6	SF6						
Other	L113						

[Add product](#)

Date filled in  
Day: 24 Month: 03 Year: 2013

Place filled in

[Save](#)

Click Save to save your changes  
You will still be able to modify the report afterwards

*Important notice: Producers have to confirm, that they filed into products only refrigerants from certified companies (bought in Slovakia or by own import). In this way doubled counting of refrigerants and reported amounts from products and containers is avoiding.*

### A4.2.3 Reporting of Type of Use (for New Equipment or for Recharge/Service, Recovery, Reclaimed, Disposal) – Logbook Leaklog

Almost complete activity data used for inventory preparation in the category 2.F is covered by the web reporting system Leaklog. Especially the refrigeration is very complex, there are numerous of small enterprises. This web reporting system receives data from more than 1 200 companies. This system was introduced in 2009 and is still in operation. Therefore, also trends are consistent.

Reporting is made by the Logbook software Leaklog. It includes:

- Quick overview, survey;
- List of customers;
- Cooling circuits;
- Details of all maintenance work and repairs;
- Leakage ratio;
- Refrigerants in store;
- Refrigerants added, recovered, reclaimed and disposed.

Each contractor has to enter the website of notified body with its name and password. Which data are filled in and all details are listed above (*Figures A4.2.5 and A4.2.6*).

**Figure A4.2.5: Main outputs of logbook**




Figure A4.2.6: Procedure of data reporting of F-gases

**Table of Inspections: CHLJ-1**

ID	Company	Address	E-mail	Phone
00000001	OBAL a.s. Nové Mesto nad Váhom			

ID	Name	Device	Manufacturer	Type	Year of purchase	Commissioned on	Refrigerant	Oil
00001	CHLJ-1	Proxy-zvárača linka	Proxy		1996	07/05/2009	8 kg R22	1 kg AB (Alky/benzene oil)

Date	Visual and aural check	Direct leak check (location)	Refrigerant addition	Annual leakage	Refrigerant recovery	Oil addition	Inspector	Operator	Remedies	Assembly record No.
15/12/1999 09:52	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci	Vyčistenie výparníka	
13/12/2000 10:25	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci		
30/03/2001 12:04	No	Yes	0.0	0	0.0	0.0	Peter	Karol		
18/02/2002 12:12	No	Yes	0.5	6.25	0.0	0.0	Matuš	vedúci	8 vedný schreder...	2012-1-8-oprava
13/08/2002 12:14	No	Yes	0.0	0	0.0	0.0	Peter	Karol	Vyčistenie ventil...	
02/03/2003 12:30	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci		
04/11/2003 12:31	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci		
03/02/2004 12:39	No	Yes	0.5	18.75	0.0	0.0	Matuš	vedúci	čistenie kondenz...	
08/09/2004 12:40	No	Yes	1.0	0	0.0	0.0	Matuš	vedúci	výmena matice a t...	
03/03/2005 12:42	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci		
03/03/2005 12:58	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci		
03/03/2006 13:00	No	Yes	1.5	43.75	0.0	0.0	Matuš	vedúci	vadný pertel 80...	
24/11/2006 13:01	No	Yes	2.0	0	0.0	0.0	Matuš	vedúci	praskla trubka	
04/04/2007 13:13	No	Yes	0.0	0	0.0	0.0	Matuš	vedúci		
13/05/2007 13:09	No	Yes	0.0	18.75	0.0	0.0	Matuš	vedúci		
15/05/2007 13:06	No	Yes	1.5	0	0.0	0.0	Jozef Mrkvica	vedúci	výmena tesnenia	
<b>Sum</b>			<b>7</b>	<b>9.72222</b>	<b>0</b>	<b>0</b>				


  

**Warnings**

Date	Warning
18/02/2002 12:12	Refrigerant leakage above limit, *Únik chladiva
03/02/2004 12:39	Refrigerant leakage above limit
08/09/2004 12:40	Refrigerant leakage above limit
03/03/2006 13:00	Refrigerant leakage above limit
24/11/2006 13:01	Refrigerant leakage above limit
04/04/2007 13:13	*Netesnosť ventilov kompresora, *Zanesenie kondenzátora
15/05/2007 13:06	Refrigerant leakage above limit, *Únik chladiva

The inserted data can be presented in table with differentiation by category (Figure A4.2.7).

Figure A4.2.7: Table of leakages by application



**Figure A4.2.8: Conversion of mixtures to single substances according to the new charges (NN), leaks (Ú) and calculated leaks (ÚV) for different subcategories – Leaklog**

Chladivá / Sklad / Organizácie / Certifikáty / Nové náplne a úniky podľa druhu s výrobkami / Spolu s výrobkami / Trend vývoja

Chladivá / Kilogramy / Tony / Koefficienty: Zohľadniť / Nezohľadniť / Kategórie oznámené firmami: Zohľadniť / Nezohľadniť

Upraviť údaje v tabuľkách pre nádoby pre rok 2013: Štandardné

Upraviť údaje v tabuľkách pre nádoby pre rok 2013: Rozdiel vypočítaného úniku

Oznámené nové náplne a úniky podľa druhu zariadení za rok 2013

Skratky: NN - Nová náplň, Ú - Únik, ÚV - Únik vypočítaný

Chlad.	Hobklim			Komerčné chladenie			Priemyselné chladenie			Prepravné chladenie			Klimatizácia a TČ			Domáce chladenie			Hasenie			PUR izolácie			Aerosoly			SF6			Iné			Σ					
	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV	NN	Ú	ÚV		Σ				
CSH12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	392.64	0	0	392.64				
CF4	0	0	0	0	0	0	0	0	0	0	0	18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	23	0	0	18					
CO2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1200	0	0	1200						
Ethen	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3.7	0	0	3.7						
L113	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0						
R11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0						
R115	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0						
R116	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1.4	4.9	0	1.41	4.86	4.86				
R12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
R123	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
R1234yf 2287.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7.3	35.6	2287.47	7.34	35.6	2323.07				
R124	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	546	3.7	-546	728.81	3.73	-546	182.81			
R125	0.2	20.3	20.3	7804.8	13429	13429	7564.3	12257.7	12257.7	751	1626.3	1626.3	21410.1	4170.9	4170.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	816.1	2206.4	8837.7	38346.59	33710.62	40341.89	78688.49			
R13	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
R134a	44447.1	25878.9	25878.9	12361.6	9184	9184	18415	10827.4	10827.4	221.6	854.3	854.3	6209.7	7394.8	7394.8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6548.5	0	0	824.8	1417.9	34568.8	89028.38	55557.23	88708.18	177736.56
R141b	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
R142b	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2.1	0	0	2.15	0	0			
R143a	0	8.3	8.3	7781.4	14351.4	14351.4	5256.4	12460.8	12460.8	760.5	1907.9	1907.9	169	73.2	73.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	68.4	588.9	2139.6	14035.69	29390.55	30941.21	44976.9			
R152a	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1.2	0	69.9	1.24	0	69.9				
R218	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1.2	5.4	9.1	1.17	5.45	9.09	10.26			
R22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	370.2	260.5	-273.2	655.17	260.5	-273.2	381.97			
R227ea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	312	94.2	94	4506.16	94.2	94	4600.16			
R23	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2.5	19.7	43.2	2.5	19.65	43.24	45.74			
R236fa	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	61	0	9	81	0	9	90			
R245fa	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1520.2	0	0	1520.2	0	0	1520.2			
R290	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6.7	269	232.19	6.65	269	501.19				
R318	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
R32	0.2	13	13	1191.1	1222.7	1222.7	3039.1	1603	1603	106.9	11.3	11.3	21149.7	3984.6	3984.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	447.2	36.1	4676.7	25934.18	6870.62	11511.24	37445.42			
R365mfc	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2718	0	0	2718	0	0	2718		
R423A	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
R425A	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
R428A	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
R600	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	9.5	61.6	80.8	9.5	61.55	80.75	90.25			
R600a	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	32.5	54.8	166.7	32.5	54.81	166.7	199.21			
R601	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.3	0.3	0.03	0.34	0.33	0.36				
R601a	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.7	0	0	0.66	0.66	0.66				
S316	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
SF6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1635.3	0	1099.8	439.5	280	2735.05	439.49	280	3015.05	
Viermel	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

39 chl. 46735 25920.5 25920.5 29369 38187.1 38187.1 34274.8 37148.9 37148.9 1840 4399.8 4399.8 49500.3 15623.5 0 0 0 3991.7 0 0 4832.8 0 0 6865 0 0 1635.3 0 0 4591.2 5230.7 51600.6 183335.13 126510.53 172880.25 356215.4

#### A4.2.4 Data Processing – Inventory Preparation

The 2006 IPCC GL describe two tiers for estimating emissions. The bottom-up approach takes into account the time lag between consumption and emissions explicitly through emission factors. The top-down approach (mass balance) takes the time lag into account implicitly, by tracking the amount of virgin chemical consumed in a year that replaces emissions from the previous year. The using of two web-reporting systems allows estimation emissions in both approaches. The bottom-up approach combined with the top-down approach was used during emissions estimation in Slovakia. The process was based on the following steps:

1. Using the bottom-up approach based on the Logbook Leaklog;
2. Calculation of the total consumptions of individual gases in Slovakia based on the Leaklog;
3. Calculation of the total consumption of individual gases in Slovakia according to the top-down approach based on the older web reporting system available since 2003 (import, export, sales data of bulk chemicals and products);

4. Comparing of the total consumptions calculated by these two approaches;
5. If differences occur, the data for bottom-up approach will be corrected as follows (expert judgement based on the QA process in 2011):
  - R134a: Difference is added to leakage from mobile AC;
  - R404A: Difference is added between new charge/recharge 0.2/0.8;
  - R407C: Difference is added to new charge of stationary AC;
  - R410A: Difference is added to leakage from industrial refrigeration and stationary AC 0.1/0.9;
6. If differences below 2% occur, the data for bottom-up approach are corrected proportionally according to the operational emissions.
7. Calculation of emissions inventory by the bottom-up approach using the corrected data.

For the top-down approach the following formulas based on the structure of the reporting systems are:

*Emissions = Annual Sales of New Refrigerant – Total Charge of New Equipment + Disposal Emissions*  
 where *Annual Sales* and *Total Charge of New Equipment* are calculated by formulas presented in the 2006 IPCC Guidelines, Chapter 3, p. 7.54 (not simplified formulas).

For bottom-up approach the following formulas are used:

*Emissions = Emissions from new fillings + Operational emissions + Disposal emissions*

where: *Emissions from new fillings* represent 1% (EF) of the new charges filled in Slovakia (*Chemicals to Charge Domestically Manufactured and Assembled Equipment + Chemical to Charge Equipment that is not Factory-Charged*).

*Operational emissions*: The approach described in IPCC 2006 GL assumes that servicing of equipment restocks the bank of single chemical and thus the amount of gas used for servicing represents the operational emissions. Slovakia adopted this assumption with a modification in 2017 submission. The servicing of equipment restocks the bank of chemical and its amount used at servicing equals to the emissions. However, equipment that is few years before decommissioning is not serviced and bank is not restock at this equipment. Therefore, the operational emissions are composed from two terms in this submission: (i) data from servicing of equipment; (ii) emissions from non-serviced equipment few years before its decommissioning. The first term in the operational emissions represents the consumption of gases for servicing and container management (these data are reported in Leaklog). It is assumed that the chemical used for servicing restocks the emissions from the bank and thus the bank of the chemical remains constant. The second term in operational emissions represents emissions from non-serviced equipment few years before its decommissioning. These emissions decrease the amount of chemical in equipment and the equipment contains only a part of the chemical at its decommissioning. The product life factors, number of years when the equipment is not serviced and fraction of gas remaining at its decommissioning is consistent. These emissions do not restore the bank of the chemical and are subtracted from the bank.

*Disposal emissions* represent the emissions from the retired equipment.

For the consistency of operational emissions, the bank of chemical is necessary to follow. The bank is calculated as follows:

$Bank_{in\ year\ t} = Bank_{in\ year\ t-1} + New\ additions\ to\ bank - Chemical\ in\ retired\ equipment - Operational\ emissions\ from\ non-serviced\ equipment$

where: *New additions to bank = Chemicals to Charge Domestically Manufactured and Assembled Equipment + Chemicals to Charge Equipment that is not Factory-Charged + Chemicals Contained in Imported Equipment Already Charged – Chemicals Contained in Exported Equipment Already Charged.*

## Annex 4.3. Balance of Urea: Import-Export-Production-Use Balance

In the GHG inventory, the downstream of CO<sub>2</sub> emission from ammonia production to urea production is reported. The comparison of CO<sub>2</sub> emissions from the ammonia production and net CO<sub>2</sub> emissions reported is shown in [Table A4.3.1](#). The difference is caused by using of the part of “produced” CO<sub>2</sub> to urea production. In Slovakia, the urea is used in the agriculture as fertilizer (reported under 3.H) and DeNO<sub>x</sub> application (in cars and in plants, reported in 2.D.3). The difference among the CO<sub>2</sub> used for urea production and CO<sub>2</sub> reported is shown in [Table A4.3.2](#). This difference is attributed to the export of urea in Slovakia. This Annex deals with the comparison of “CO<sub>2</sub> exported in urea” from Slovakia and the above-mentioned difference. The comparison was made since 2010 because no older data were obtained from the Statistical Office of the Slovak Republic due to the change in statistical methodology of import-export data.

**Table A4.3.1: Comparison of technological and net CO<sub>2</sub> emissions since 2010**

YEAR	Ammonia Production	CO <sub>2</sub> Emissions from Ammonia Production	Net CO <sub>2</sub> Emissions
	kt		
2010	233.56	484.65	387.58
2011	455.48	779.42	577.96
2012	377.30	717.42	545.98
2013	474.91	888.08	674.48
2014	346.27	660.68	529.65
2015	476.94	884.82	638.58
2016	403.96	787.01	563.81
2017	458.88	873.80	632.94
2018	516.74	1 028.79	790.46
2019	491.95	822.68	688.35
2020	545.23	883.52	703.09
2021	580.51	930.46	768.62
2022	462.12	750.41	637.81
2023	413.54	693.10	509.92

**Table A4.3.2: Comparison of CO<sub>2</sub> used for urea production and CO<sub>2</sub> reported from the use of urea**

YEAR	CO <sub>2</sub> Used for Urea Production	CO <sub>2</sub> Emissions Reported in 3.H Category	CO <sub>2</sub> Emissions Reported in 2.D.3 Category	CO <sub>2</sub> Emissions Reported in Slovakia From Use of Urea	Difference (“Missing CO <sub>2</sub> ”)
	kt				
2010	97.074	30.939	2.012	32.951	64.123
2011	201.465	39.708	3.484	43.191	158.274
2012	171.446	45.418	3.925	49.344	122.102
2013	213.603	51.993	6.090	58.083	155.520
2014	131.033	57.941	6.504	64.444	66.589
2015	246.239	60.920	6.315	67.235	179.004
2016	223.200	63.071	8.767	71.838	151.362
2017	240.860	63.534	10.669	74.202	166.658
2018	238.324	65.966	10.571	76.538	161.786
2019	134.339	63.539	16.696	80.235	54.104
2020	180.420	63.666	18.951	82.617	97.806
2021	161.834	63.633	30.009	93.642	68.191

YEAR	CO <sub>2</sub> Used for Urea Production	CO <sub>2</sub> Emissions Reported in 3.H Category	CO <sub>2</sub> Emissions Reported in 2.D.3 Category	CO <sub>2</sub> Emissions Reported in Slovakia From Use of Urea	Difference ("Missing CO <sub>2</sub> ")
	kt				
2022	112.594	56.619	17.979	74.598	37.996
2023	183.185	56.070	27.130	83.200	99.985

Data for the comparison were obtained from the urea producer and from the Statistical Office of the Slovak Republic. Data provided by the producer deals with the use of urea for DeNOx application and the composition of urea containing fertilizers. Urea is used for DeNOx application as the product AdBlue (solution containing approx. 30% of urea) and as the so-called technical urea (solution containing 40% of urea). Data were provided as pure urea ([Table A4.3.3](#)). According to the producer it can be assumed that all urea for DeNOx application was exported (except of data that are reported in the NID in 2.D.3 category). Import and export data about fertilizers were obtained from the Statistical Office of the Slovak Republic under the commodity codes: (i) 31021010, "Urea containing more than 45% by weight of nitrogen on the dry anhydrous product"; (ii) 31028000, "Mixtures of urea and ammonium nitrate in aqueous or ammoniacal solution". Because urea contains 46% of nitrogen, it can be assumed that the commodity code 31021010 represents the pure urea and export-import difference can be easily calculated from the export and import data ([Table A4.3.3](#)). On the other hand, the content of urea in products reported under commodity code 3102800 can vary. According to the Slovak law 555/2002 Z. z. the fertilizers with the different content of urea can be used. The nitrogen originating from the urea can be in the range 11-51%. Because import data are reported as kilograms of nitrogen, the amount of urea imported to Slovakia was calculated using this range. According to the data provided by the Slovak fertilizer producer, the fertilizer DAM-390 represents more than 98% of the export of this commodity. It is the mixture of ammonium nitrate and urea containing 29-30% of N, from which 15.5% N originates from urea, the rest is from AN. To ensure conservatism we assumed that 50% of nitrogen originates from urea. Data about import and export of the commodity 31028000 are provided in [Table A4.3.4](#).

**Table A4.3.3:** Amounts of exported urea for DeNOx application and import-export data for the commodity code 31021010 since 2010

YEAR	Urea Exported for DENOX Application	Import of the Commodity Code 31021010	Export of the Commodity Code 31021010	Export-Import
	kt		kt N	
2010	24.781	63.758	87.885	24.127
2011	51.43	51.999	110.524	58.525
2012	42.538	61.218	95.638	34.419
2013	52.945	42.736	127.442	84.706
2014	32.195	75.848	77.108	1.259
2015	56.651	67.233	159.628	92.395
2016	47.307	88.352	139.278	50.926
2017	62.67	88.158	144.782	56.623
2018	67.838	63.520	107.337	43.817
2019	45.982	85.887	78.164	-7.723
2020	21.028	61.421	91.333	29.912
2021	26.994	106.510	112.078	5.568
2022	31.283	224.272	162.133	-62.139
2023	1.777	146.940	152.321	5.381

**Table A4.3.4: Import-export data for the commodity code 31021010 since 2010**

YEAR	Import of the Commodity Code 31028000	Export of the Commodity Code 31028000	Imported Urea (Range Based on the Possible Urea Content)	Exported Urea	Export-Import (Range Based on the Possible Urea Content)
	kt N		kt		
2010	8.622	25.367	2.062-9.559	27.573	18.014-25.512
2011	8.145	46.889	1.948-9.031	50.966	41.935-49.018
2012	7.970	37.384	1.906-8.837	40.635	31.799-38.729
2013	3.929	51.481	0.939-4.356	55.957	51.602-55.018
2014	4.519	36.075	1.081-5.01	39.212	34.202-38.131
2015	5.540	63.135	1.325-6.142	68.625	62.483-67.300
2016	6.242	54.192	1.493-6.92	58.904	51.983-57.411
2017	6.242	54.110	1.493-6.92	58.816	51.895-57.323
2018	5.243	64.114	1.254-5.813	69.689	63.876-68.436
2019	4.306	50.128	1.030-4.774	54.487	49.713-53.458
2020	1.741	50.121	0.416-1.930	54.479	52.549-54.063
2021	4.076	54.087	0.975-4.519	58.790	54.271-57.815
2022	4.366	145.670	1.044-4.841	158.337	153.497-157.293
2023	4.390	121.158	1.050-4.868	131.69	126.825-130.643

Emission factor of CO<sub>2</sub> from urea is based on the stoichiometry and it is 0.73 t CO<sub>2</sub> / t of urea. Calculated data on the “CO<sub>2</sub> exported” based on the data presented in [Table A4.3.4](#) and their comparison with the difference in the reporting data (so called “missing CO<sub>2</sub>” in [Table A4.3.2](#)) are listed in [Table A4.3.5](#). The negative values in the last column represent the “good” result, it means that there is not missing CO<sub>2</sub> in this balance. In an ideal balance the difference should be zero, however, there were made several assumptions in this balance and change in stocks were also not considered. The red values (for years 2012 and 2014) mean that there is missing CO<sub>2</sub> in this import-export balance. However, when looking to the difference in years 2013 and 2015, the difference is much higher than usual. It can be assumed that the positive value of missing CO<sub>2</sub> is caused by the time lag between the production and export of the urea products.

**Table A4.3.5: Balance of the “export/import CO<sub>2</sub>” from the use of urea**

YEAR	CO <sub>2</sub> from the Exported DENOX Applications	CO <sub>2</sub> from the Commodity Code 31021010	CO <sub>2</sub> from the Commodity Code 31028000	“CO <sub>2</sub> Exported”	“Missing CO <sub>2</sub> ”	Difference
	Gg					
2010	18.09	38.289	13.150-18.623	69.529-75.002	64.123	(-5.406)-(-10.879)
2011	37.544	92.877	30.613-35.783	161.033-166.204	158.274	(-2.760)-(-7.930)
2012	31.053	54.621	23.213-28.272	108.887-113.946	122.102	13.215-8.156
2013	38.650	134.425	37.669-40.163	210.743-213.237	155.520	(-57.717)-(-55.223)
2014	23.502	1.998	24.968-27.836	50.468-53.337	66.589	13.252-16.121
2015	41.355	146.627	45.612-49.129	233.594-237.111	179.004	(-58.107)-(-54.59)
2016	34.534	80.817	37.948-41.910	153.299-157.261	151.362	(-5.899)-(-1.937)
2017	45.749	89.858	37.884-41.846	173.491-177.454	166.658	(-10.796)-(-6.834)
2018	49.522	69.536	46.630-49.958	165.687-169.015	161.786	(-7.229)-(-3.901)
2019	33.567	-12.256	36.291-39.024	57.601-60.335	54.104	(-6.231)-(-3.497)
2020	15.350	47.4659	38.361-39.466	101.18-102.285	97.806	(-4.479)-(-3.374)
2021	19.706	8.835	39.618-42.205	68.159-70.746	68.191	(-2.555)-0.032
2022	22.837	-98.612	112.052-114.824	36.277-39.049	37.996	(-1.053)-1.718
2023	1.297	8.539	92.582-95.369	102.419-105.206	99.985	(-5.220)-(-2.433)



---


## Annex 4.4. QA/QC of Product Life Factors in 2.F.1 category

A detailed look at the product life factors in the subcategories 2.F.1.a, 2.F.1.c, 2.F.1.d and 2.F.1.f shows large variations as well as inconsistent development of reported product life factors. It is due to several factors:


- (i) The methodology used (hybrid Tier 2a/Tier 2b). It is assumed that all leaks in a given year will be replenished in a given year during equipment servicing. The practice of having all equipment serviced every year is not common in Slovakia. A typical example is MAC in cars, where, according to the automotive association, MAC service is performed on average at 2-4 annual intervals. This results in year-on-year jumps in servicing emissions, i.e. PLF. This is also very noticeable with gasses that have a small bank, each small increase in servicing will be reflected in a jump in PLF.
- (ii) It was tried to partially compensate the above mentioned factor using assumption that the equipment is not serviced a few years before the planned end of operations. These years and the PLFs used are listed in [Table 4.51](#).
- (iii) By reporting organizations in the Leak log database. Organizations do not report each individual activity to the database but they report it at certain time intervals (e.g. monthly). Then they report the type of gas and all the operations they performed with it in summary and assign it to the category in which they performed the most activities. For example, the organization can carry out multiple replacements for leaks with R410A gas, with 60% of these replacements falling into subcategory 2.F.1.f, but the remaining 40% into other categories. However, the organization reports all additions to leaks in one record and assigns it to category 2.F.1.f, because that's where the gas was used the most. We're working with the assumption (since there are several hundreds of organizations involved in this activity) that when processing the database for the whole year, these deviations compensate each other. However, they can only partially compensate. Thus, the division of F-gases into individual subcategories may not be completely correct, and this may cause year-to-year changes in individual subcategories. On the other hand, the emissions calculated by our procedure are certainly more accurate than the calculation using some more general emission factor. It should also be noted that although the division into categories 2.F.1.a, 2.F.1.c, 2.F.1.d and 2.F.1.f may not be completely correctly allocated for every year, the output from the overall category 2.F.1 is correct, because all movements of F-gases are already summarized there.
- (iv) PLFs prior 2011 do not show outliers or inconsistencies. It is due to the recalculation of those emissions to ensure the consistency of older emissions data as outlined in the Chapter 4.12.5 and 4.12.9. Since 2012, emissions have been estimated using the Tier 2a/Tier 2b hybrid methodology.

To transparently describe the reasons for the large variations and inconsistent development of reported product life factors of 2.F.1 subcategories, the aggregated product life factors of F-gases were calculated ([Figures A4.4.1 – A4.4.4](#)). The aggregation was chosen on the level of 2.F.1 category due to the factor described in the bullet (iii).


**Figure A4.4.1:** Comparison of time series product life factor of R134a in the subcategories with the product life factor on the aggregated level of 2.F.1 category




**Figure A4.4.2:** Comparison of time series product life factor of R125 in the subcategories with the product life factor on the aggregated level of 2.F.1 category



**Figure A4.4.3:** Comparison of time series product life factor of R143a in the subcategories with the product life factor on the aggregated level of 2.F.1 category



**Figure A4.4.4:** Comparison of time series product life factor of R32 in the subcategories with the product life factor on the aggregated level of 2.F.1 category



---

It can be seen from *Figures A4.4.1 – A4.4.4* that on the aggregation level of 2.F.1 category, the product life factors of all gases do not show large variations and inconsistent development. Product life factors of R134a and R143a are approximately the same for the whole time series while product life factors of R125 and R32 slowly decreases. However, this decrease is without significant outliers.

It can be concluded that behind the major changes in product life factors is the method of reporting data to the Leaklog database in aggregated form (bullet (iii)). Small changes in aggregated product life factors are caused by the practice of no servicing the equipment every year.

In conclusion, it can be stated that the product life factors are consistent at the 2.F.1 category aggregation level and thus the emission data fulfil TACCC.

**Table A5: Explanation of empty cell (in red) – due to special condition of the ETF Software, NK were not included in CRT.**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	Unspecified mix of HFCs and PFCs	SF <sub>6</sub>	NF <sub>3</sub>	NO <sub>x</sub>	CO	NMVOC	SO <sub>x</sub>	Total GHG emissions
	(kt)												CO <sub>2</sub> equivalent (kt)
<b>2. Total industrial processes</b>	6 718.99	0.57	0.40	437.89	0.01	NO	0.00	NO	5.27	70.11	26.29	4.71	7 292.59
<b>2.A. Mineral industry</b>	1 991.90	NO	NO						0.31	0.82	0.15	0.35	1 991.90
2.A.1. Cement production	1 294.23											IE	1 294.23
2.A.2. Lime production	475.06												475.06
2.A.3. Glass production	12.52												12.52
2.A.4. Other process uses of carbonates	210.09	NO	NO						0.31	0.82	0.15	0.35	210.09
<b>2.B. Chemical industry</b>	918.20	0.01	0.18	NO	NO	NO	NO	NO	0.83	1.02	2.03	1.51	965.60
2.B.1. Ammonia production	509.92	0.01	0.00						0.08	0.00	0.00	0.01	510.59
2.B.2. Nitric acid production			0.18						0.24				46.73
2.B.3. Adipic acid production	NO		NO						NO	NO	NO		NO
2.B.4. Caprolactam, glyoxal and glyoxylic acid production	NO		NO								NO	NO	NO
2.B.5. Carbide production	31.82	NO							0.05	0.13	0.00	0.01	31.82
2.B.6. Titanium dioxide production	NO												NO
2.B.7. Soda ash production	NO												NO
2.B.8. Petrochemical and carbon black production	376.47	NA,NO							NO	NO	NO	NO	376.47
2.B.9. Fluorochemical production				NO	NO	NO	NO	NO					NO
2.B.10. Other	NO	NO	NO	NO	NO	NO	NO	NO	0.46	0.89	2.03	1.49	NO
<b>2.C. Metal industry</b>	3 748.67	0.56	0.00	NO	0.01	NO	NO	NO	3.90	67.43	0.67	2.82	3 765.58
2.C.1. Iron and steel production	3 733.94	0.56							2.93	58.66	0.36	1.65	3 749.53
2.C.2. Ferroalloys production	7.77	0.00							0.00	0.11	0.01	0.00	7.90
2.C.3. Aluminium production	6.88				0.01		NO		0.40	6.95	0.02	0.91	6.89
2.C.4. Magnesium production	NO			NO	NO	NO	NO		NO	NO	NO	NO	NO
2.C.5. Lead production	0.08								0.00	0.00	0.00	0.00	0.08
2.C.6. Zinc production	NO								NO	NO	NO	NO	NO
2.C.7. Other	NO	NO	0.00	NO	NO	NO	NO	NO	0.57	1.70	0.28	0.25	1.18
<b>2.D. Non-energy products from fuels and solvent use</b>	60.21	NA,NE,NO	NA,NE,NO						NA,NE	NA,NE	18.58	0.03	60.21
2.D.1. Lubricant use	30.54	NE	NE						NA	NA	NA	NA	30.54
2.D.2. Paraffin wax use	2.54	NE	NE						NA	NA	NA	NA	2.54
2.D.3. Other	27.13	NA,NO	NA,NO						NE	NE	18.58	0.03	27.13
<b>2.E. Electronics industry</b>			NO	NO	NO	NO	NO	NO					NO
2.E.1. Integrated circuit or semiconductor			NO	NO	NO	NO	NO	NO					NO
2.E.2. TFT flat panel display			NO	NO	NO	NO	NO	NO					NO
2.E.3. Photovoltaics				NO	NO	NO	NO	NO					NO
2.E.4. Heat transfer fluid				NO	NO	NO	NO	NO					NO
2.E.5. Other			NO	NO	NO	NO	NO	NO					NO
<b>2.F. Product uses as substitutes for ODS</b>				437.89	NO	NO	NO	NO					437.89
2.F.1. Refrigeration and air conditioning				398.92	NO	NO	NO	NO					398.92
2.F.2. Foam blowing agents				1.73	NO	NO	NO	NO					1.73
2.F.3. Fire protection				27.20	NO	NO	NO	NO					27.20
2.F.4. Aerosols				10.04	NO	NO	NO	NO					10.04
2.F.5. Solvents				NO	NO	NO	NO	NO					NO
2.F.6. Other applications				NO	NO	NO	NO	NO					NO
<b>2.G. Other product manufacture and use</b>	NO	NO	0.21	NO	NO	NO	0.00	NO	0.02	0.50	0.04	0.00	71.41
2.G.1. Electrical equipment				NO	NO	NO	0.00	NO					14.70

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	Unspecified mix of HFCs and PFCs	SF <sub>6</sub>	NF <sub>3</sub>	NO <sub>x</sub>	CO	NMVOC	SO <sub>x</sub>	Total GHG emissions
	(kt)			CO <sub>2</sub> equivalent (kt)			(kt)						CO <sub>2</sub> equivalents (kt)
2.G.2. SF <sub>6</sub> and PFCs from other product use					NO		NO						NO
2.G.3. N <sub>2</sub> O from product uses			0.21										56.71
2.G.4. Other	NO	NO	NO	NO	NO	NO	NO	NO	0.02	0.50	0.04	0.00	NO
2.H. Other <sup>(5)</sup>	NO	NA,NO	NA,NO	NO	NO	NO	NO	NO	0.22	0.34	4.82	0.00	NA,NO
2.H.1. Pulp and paper	NO	NO	NO						NO	NO	NO	NO	NO
2.H.2. Food and beverages industry	NO	NA	NA						NA	NA	4.16	NA	NA,NO
2.H.3. Other (please specify)	NO	NO	NO	NO	NO	NO	NO	NO	0.22	0.34	0.66	0.00	NO

<b>CHAPTER 5. AGRICULTURE (CRF 3)</b> .....	<b>232</b>
5.1 Overview of the Agriculture Sector.....	232
5.2 Category-specific Improvements and Implementation of Recommendations .....	237
5.3 Category-Specific QA/QC and Verification.....	238
5.4 Category-specific Recalculations.....	242
5.5 National Circumstances and Time-series Consistency.....	246
5.6 Uncertainties .....	249
5.7 Enteric Fermentation (CRF 3.A).....	256
5.8 Manure Management (CRF 3.B.1) – CH <sub>4</sub> Emissions.....	266
5.9 Manure Management (CRF 3.B.2) – N <sub>2</sub> O Emissions .....	275
5.10 Indirect N <sub>2</sub> O Emissions from Manure Management (CRF 3.B.2.5).....	282
5.11 Rice Cultivation (CRF 3.C) .....	284
5.12 Agricultural Soils (CRF 3.D).....	285
5.13 Prescribed Burning of Savannas (CRF 3.E) .....	304
5.14 Field Burning of Agricultural Residues (CRF 3.F) .....	304
5.15 Liminig (CRF 3.G) .....	304
5.16 Urea Application (CRF 3.H).....	306
5.17 Other Carbon-containing Fertilizers (CRF 3.I) .....	307

## CHAPTER 5. Agriculture (CRF 3)

This Chapter was prepared using GWP<sub>100</sub> taken from the [5<sup>th</sup> Assessment Report of the IPCC](#) by the sectoral experts and institutions involved in the National Inventory System of the Slovak Republic:

INSTITUTE	CHAPTER	SECTORAL EXPERT
Slovak Hydrometeorological Institute	all	Kristína Tonhauzer
Research Institute for Animal Production	3.A and 3.B supported calculation and background data	Zuzana Palkovičová Ondrej Pastierik
Slovak Hydrometeorological Institute	3.D Estimation of humid area of the Slovak Republic	Peter Tonhauzer

The Agriculture sector is the fourth largest sector of the GHG emissions inventory of the Slovak Republic with the contribution equal to 6.5% on the total GHG emissions.

The emissions of greenhouse gases from agricultural activities include:

CH<sub>4</sub> emissions from the Enteric Fermentation (3.A) and the Manure Management (3.B);

N<sub>2</sub>O emissions from the Manure Management (3.B) and the Agricultural Soil (3.D);

CO<sub>2</sub> emissions from the Liming (3.H) and the Urea Application (3.G);

Emissions inventory of NVMOC and NO<sub>x</sub> were estimated and information is provided in the [Informative Inventory Report](#) of the Slovak Republic.

Categories 3.C and 3.E are not reported due to the weather conditions and climatic zone of Slovakia. Category 3.F is reported as not occurring, burning of fields is prohibited by the law.

### 5.1. Overview of the Agriculture Sector

The share of the agriculture and food industry in the national economy has decreased in the macro-economic indicators (intermediate consumption, sectoral employment, gross production) except gross value added, and sectoral employment in 2022 compared to 2021 and increased in parameter employee's average wage by 0.08%. The share of foreign agri-food trade in exports (0.05%) and imports decreased by 0.37%. The increase in gross agricultural production was caused by a higher value of plant production by 12.2% with a simultaneous increase in animal production by 3.8%, while the overall increase in gross agricultural production was mainly contributed by an increase in prices by 8.9% with a simultaneous increase in the number of subsidies for products. Agriculture, according to data, achieved a negative moderate interannual economic result in 2023. The subsidies from the Common Agricultural Policies (CAP) played the stabilized role of financial support for Slovak agriculture, which help the majority of the farmers avoid the negative economic situation. The subsidies from the CAP decreased by 12.3% due to a decrease of the EU resources by 19%.


Crop production had the continuing dominant share in the economy compared to animal production (60% to 40%). Decrease in the production of most commodities of crop production mainly due to the decrease in harvesting areas and the decrease or stagnation of harvests per hectare, except for sunflower, soybean, sugar beet, fruit and vegetables. Number of livestock decreased in all species with impacts decrease of animal products except slaughter poultry (1.8%), cattle (9.8%) and pigs (3.9%) ([Green Report 2023](#)).



The emissions balance is compiled annually based on the sectoral statistics on animal livestock, animal performance and consumption of organic and inorganic fertilizers, in recent years on the regional level. The Ministry of Agriculture and Rural Development of the Slovak Republic (MPRV SR) issues annual agricultural statistics in the Green Report, part of which is dedicated to agriculture and food. Activity data are also available in the Statistical Yearbooks published by the Statistical Office of the Slovak Republic (ŠÚ SR).

The emissions inventory in agriculture is prepared in the cooperation with the National Agricultural and Food Centre - the Research Institute for Animal Production in Nitra (NPPC - VÚŽV). The NPPC - VÚŽV provided activity data and parameters, improved the methodology and ensured QA/QC activities in animal inventory in the CRF categories 3.A and 3.B. Activity data on number of the livestock and animal productions are provided annually by the ŠÚ SR. The Central Control and Testing Institute in Agriculture (UKSÚP) provides the soil data to the SHMÚ annually, based on cooperation agreement between the both institutions. Emission Inventory System in the Agriculture sector is described on [Figure 5.1](#).

**Figure 5.1: Emission Inventory System in the Agriculture sector**




The largest share of methane emissions was generated by enteric fermentation of cattle, which produced 32.52 Gg (91%) of methane within the sector in 2023. The major source of N<sub>2</sub>O emissions is agricultural soils with a share of 78%, followed by the category 3.B representing 22% on the total N<sub>2</sub>O emissions. Regarding N<sub>2</sub>O, direct emissions from synthetic fertilization are the most significant emissions source and it produced 0.845 Gg of N<sub>2</sub>O (22%) within the sector in 2023.

CH<sub>4</sub> emissions are calculated separately for each animal sub-category in methane model. For categories 3.B and 3.D, N<sub>2</sub>O emissions are calculated based on an N-flow concept, more information is in the Chapter 5.9. In categories 3.G and 3.H, CO<sub>2</sub> emissions are estimated for liming and urea application in line with the IPCC 2006 GL.

[Figures 5.2](#) and [5.3](#) and [Tables 5.1](#) and [5.2](#) show overall emission trends since the base year 1990 according to gases and major categories. [Table 5.3](#) shows an overview of the GHG gases and tiers. In the Slovak Republic, agricultural production stopped increasing in the late '90s. The decrease was followed by a drop during the years 1990 – 2002, because of the economic and political transition of the country. After entering the EU, agriculture was stabilized. Improving conditions in the Agriculture sector,

regeneration of crop production and mineral fertilizers use caused that emissions have increased in the last six years. The inter-annual growth of emissions was caused due to increase of organic nitrogen fertilizers mainly in categories 3.D.1.4 Crop residues and 3.D.1.2C Other nitrogen organic fertilizers into soils. Increase of nitrogen application into soils had positive effect on increase of yield of selected crops (cereals, legumes and oil plants).

**Figure 5.2:** Trend in aggregated emissions (in Gg of CO<sub>2</sub> eq.) by categories within the Agriculture sector in 1990 – 2023




**Table 5.1:** Trend of GHG emissions by gases in the Agriculture sector in particular years

YEAR	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	NM VOC	NO <sub>x</sub>
	Gg				
1990	61.017	137.008	7.448	21.189	13.619
1995	53.291	86.038	4.122	13.440	6.149
2000	46.442	63.886	3.593	11.268	6.203
2005	29.590	54.739	3.927	10.112	6.523
2010	39.170	48.294	4.086	8.358	6.512
2011	54.850	47.256	3.239	8.281	6.986
2012	56.721	48.142	3.023	8.145	6.287
2013	63.944	47.361	3.517	8.132	6.794
2014	69.903	46.426	4.001	8.108	7.144
2015	73.333	46.319	3.098	8.189	6.857
2016	69.845	44.746	3.905	7.987	7.322
2017	66.152	44.934	3.403	8.046	7.170
2018	70.177	44.895	3.273	7.660	7.537
2019	68.250	43.621	3.538	7.412	7.443
2020	72.117	42.749	3.656	6.984	7.144
2021	69.570	40.927	3.282	6.95	7.104
2022	60.841	40.709	2.888	6.657	6.531
2023	72.468	40.663	3.663	7.300	5.970

**Table 5.2:** Trend of GHG emissions by categories in the Agriculture sector in particular years

YEAR	3.A ENTERIC FERMENTATION	3.B MANURE MANAGEMENT	3.D AGRICUL. SOILS	3.G LIMING	3.H UREA APPLICATION
	Gg of CO <sub>2</sub> eq. (AR 5)				
1990	3 120.016	1 276.877	1 413.122	43.786	15.288
1995	1 992.56	801.508	707.342	36.255	15.288
2000	1 481.335	633.697	625.921	32.035	12.101
2005	1 290.374	529.717	753.369	8.838	20.313
2010	1 169.152	428.057	837.823	7.715	30.939
2011	1 158.863	401.912	620.829	14.137	39.708
2012	1 177.210	413.260	558.500	10.550	45.418
2013	1 166.653	395.030	696.454	11.159	51.993
2014	1 139.133	399.572	821.365	11.187	57.941
2015	1 142.595	388.728	586.500	11.581	60.920
2016	1 111.450	368.310	808.017	4.967	63.071
2017	1 109.946	373.422	676.569	1.967	63.534
2018	1 102.821	384.862	636.724	3.332	65.966
2019	1 078.228	371.094	709.723	3.629	63.539
2020	1 072.755	344.402	748.348	6.251	63.666
2021	1 037.591	318.488	659.556	3.935	63.633
2022	1 038.751	309.493	556.885	0.888	56.619
2023	1 037.254	308.857	763.266	9.250	56.068

**Figure 5.3:** The share of aggregated emissions by main categories within the Agriculture sector



**Table 5.3:** Overview of the gases, methodology and tiers reported in the Agriculture sector according to the CRF categories in 2023

CATEGORY (CODE AND NAME)	METHODOLOGY/TIER	GHG GASES REPORTED
3.A.1 DAIRY CATTLE	T2/CS	CH <sub>4</sub>
3.A.1 NON-DAIRY CATTLE	T2/CS	CH <sub>4</sub>
3.A.2 MATURE EWES	T2/CS	CH <sub>4</sub>
3.A.2 GROWING LAMBS	T2/CS	CH <sub>4</sub>
3.A.2 OTHER MATURE SHEEP	T2/CS	CH <sub>4</sub>
3.A.3 SWINE	T1/D	CH <sub>4</sub>
3.A.4 GOATS	T1/D	CH <sub>4</sub>
3.A.4 HORSES	T1/D	CH <sub>4</sub>
3.A.4 RABBITS	T1/CS	CH <sub>4</sub>
3.B.1.1 DAIRY CATTLE	T2/CS	CH <sub>4</sub>

CATEGORY (CODE AND NAME)	METHODOLOGY/TIER	GHG GASES REPORTED
3.B.1.1 NON-DAIRY CATTLE	T2/CS	CH <sub>4</sub>
3.B.1.2 MATURE EWES	T2/CS	CH <sub>4</sub>
3.B.1.2 GROWING LAMBS	T2/CS	CH <sub>4</sub>
3.B.1.2 OTHER MATURE SHEEP	T2/CS	CH <sub>4</sub>
3.B.1.3 SWINE	T2/CS	CH <sub>4</sub>
3.B.1.4 GOATS	T1/CS	CH <sub>4</sub>
3.B.1.4 HORSES	T1/CS	CH <sub>4</sub>
3.B.1.4 POULTRY	T2/CS	CH <sub>4</sub>
3.B.1.4 RABBITS	T2/CS	CH <sub>4</sub>
3.B.2.1 DAIRY CATTLE	T2/CS	N <sub>2</sub> O
3.B.2.1 NON-DAIRY CATTLE	T2/CS	N <sub>2</sub> O
3.B.2.2 MATURE EWES	T1/CS	N <sub>2</sub> O
3.B.2.2 GROWING LAMBS	T1/CS	N <sub>2</sub> O
3.B.2.2 OTHER MATURE SHEEP	T1/CS	N <sub>2</sub> O
3.B.2.3 SWINE	T2/CS	N <sub>2</sub> O
3.B.2.4 GOATS	T1/CS	N <sub>2</sub> O
3.B.2.4 HORSES	T1/CS	N <sub>2</sub> O
3.B.2.4 POULTRY	T2/CS	N <sub>2</sub> O
3.B.2.4 RABBITS	T2/CS	N <sub>2</sub> O
3.B.2.5 INDIRECT N <sub>2</sub> O EMISSIONS	T1/D	N <sub>2</sub> O
3.C RICE CULTIVATION	NO	N <sub>2</sub> O
3.D.1.1 INORGANIC N FERTILIZERS	T1/D	N <sub>2</sub> O
3.D.1.2.A ANIMAL MANURE APPLIED TO SOILS	T1/CS	N <sub>2</sub> O
3.D.1.2.B SEWAGE SLUDGE APPLIED TO SOILS	T1/D (NO 2023)	N <sub>2</sub> O
3.D.1.2.C OTHER ORGANIC FERTILIZERS APPLIED TO SOILS	T1/D	N <sub>2</sub> O
3.D.1.3 URINE AND DUNG DEPOSITED BY GRAZING ANIMALS	T1/CS	N <sub>2</sub> O
3.D.1.4 CROP RESIDUES	T2/CS	N <sub>2</sub> O
3.D.1.5 MINERALIZATION/IMMOBILIZATION ASSOCIATED WITH LOSS/GAIN OF SOIL ORGANIC MATTER	T1/D	N <sub>2</sub> O
3.D.1.6 CULTIVATION OF ORGANIC SOILS	NA	NE
3.D.2.1 ATMOSPHERIC DEPOSITION	T1/D	N <sub>2</sub> O
3.D.2.2 NITROGEN LEACHING AND RUN-OFF	T2/CS	N <sub>2</sub> O
3.E PRESCRIBED BURNING OF SAVANNAHS	NA	NO
3.F FIELD BURNING OF AGRICULTURAL RESIDUES	NA	NO
3.G LIMING	T1/D	CO <sub>2</sub>
3.H UREA APPLICATION	T1/D	CO <sub>2</sub>
3.I OTHER CARBON-CONTAINING FERTILIZERS	NA	NO

## 5.2. Category-specific Improvements and Implementation of Recommendations

Improvement and implemented recommendations are described in this chapter.

### Ad 1. Revision of N losses before application of animal manure to agricultural soils


The method to derive two EF for the estimation of  $FRAC_{LOSS}$  has been changed. Specifically, EF3 is implied EF (IEF) for direct  $N_2O$  emissions from all MMS instead of using default value (Table 10.21, IPCC 2019 RF). As the consequence, loss of N through  $N_2$  ( $FRAC_{N_2}$ ) is calculated as 3 times EF3 (IEF). Remainder of EF included in the estimate  $FRAC_{LOSS}$  have not undergone any changes. The amount of N applied to soil in 3.D.1.b Organic N fertilizers and 3.D.2 Indirect  $N_2O$  emissions from managed soils categories were slightly impacted.

### Ad 2. Addition of new emission source from rabbits

Recommendation to develop new category of emissions from has been given during the 2023 review of the inventories submitted under CLRTAP. Successful implementation enabled to estimate GHG emissions from rabbits as well. New emission sources include  $CH_4$  from enteric fermentation, and both  $CH_4$  and  $N_2O$  from manure management.

As there is no official statistics (ŠÚ SR) on rabbits population representative survey has been carried out in 2023 to identify number of animals per type of breeding purpose among the adult inhabitants of Slovakia and animal waste management systems. This source of data was complemented by the rabbit farms reported themselves in the NEIS (National Emission Inventory System) as mid-, large sources of pollution (since 2006). Data from the survey has been applied consistently across the entire timeseries. It has been found that proportion of pasture on total manure excreted represent 32 % remaining part (68%) is excreted in housing.

**Figure 5.4:** Presents structure and development of rabbit population in the time series



**Table 5.4** shows shares of manure management system types on total animal waste of rabbit population. MCFs and  $N_2O$  EF were taken from Table 10.17 and Table 10.21 of IPCC 2019 RF, respectively. Method used for estimation  $CH_4$  from enteric fermentation was Tier 1 in combination with EF of 0.08 kg  $CH_4$  per head  $^{-1}$  year  $^{-1}$  taken from submission of Italian National Report in 2023 assuming that feed intake and animal characteristics are comparable with the one's occurring in Slovakia.

**Table 5.4:** Structure of MMS and parameters used for estimation of  $CH_4$  and  $N_2O$  emissions from MM

Type of MM system	Share	MCF	EF $N_2O$
-------------------	-------	-----	-----------

	% of total	% of	kg N <sub>2</sub> O-N/kg
Solid storage - unconfined piles or stacks	8.06	2.0	0.010
Solid storage - covered/compacted	53.54	2.0	0.010
Composting - Passive windrow	24.80	1.0	0.005
Municipal waste	12.80	0.0	0.000
Solid storage – Bulking agent addition	0.80	0.5	0.005

Resulting share of AWMS is 61.6% solid, composting 24.8% and 13.6% was assigned to other. Last share of AWMS aggregates Municipal waste (12.8%) and Solid storage - Bulking agent addition (0.8%). NEX used for calculation was 4.6 kg N animal<sup>-1</sup> yr<sup>-1</sup>.

Tier 2 methods of IPCC 2019 RF and country-specific EF were used for estimation N<sub>2</sub>O (Eq. 10.23) and CH<sub>4</sub> (Eq. 10.23) from manure management. Parameters VS (0.1 kg VS day<sup>-1</sup>) and B<sub>0</sub> (0.32 m<sup>3</sup> kg VS) were taken from IPCC 2019 RF, Table 10.15.

Addition of new emission sources from rabbits contributed 54.4 Kt in 2023 which represents 2.51% of total CO<sub>2</sub> eq. from Agriculture sector emissions. The highest share emissions from rabbits in CO<sub>2</sub> eq. is estimated to come from N<sub>2</sub>O from MM as shown in the [Table 5.5](#).

**Table 5.5: Contribution of rabbit categories to subsector emissions in CO<sub>2</sub>eq**

Subsector	Categories	Emissions in 2025 submission	Rabbits
		Gg CO <sub>2</sub> eq.	% of subsector total CO <sub>2</sub> eq.
3.A Enteric fermentation		1037	
	3.A.4.h.i. Rabbit (CH <sub>4</sub> )	10.2	0.98
3.B Manure management		453	
	3.A.4.h.i. Rabbit (CH <sub>4</sub> )	0.2	0.03
	3.A.4.h.i. Rabbit (N <sub>2</sub> O)	44.0	9.7

## 5.3. Category-specific QA/QC and Verification

### 5.3.1 Comparison of the National Activity Data with the FAOSTAT

According to the QA/QC Long-term Plan for agriculture in the area of consistency with the international bodies and statistics, several presentations were made on international and national conferences, publications and references were published in the [Meteorological Journal 2017](#). Results of this article were presented at the international conference [Air Protection 2017](#). Detailed information was presented in the SVK NIR 2018 ([Chapter 5.3.1](#)). The data comparison is provided annually until full consistency will be achieved. In the 2019 submission, new corrected national data on livestock, harvest and fertilisers were sent to the FAO by the national body (ŠÚ SR).

**Inorganic N-fertilizers:** The Slovak Republic has had a long-term issue in inorganic nitrogen fertilizers reporting to the world and European institutions. Data inconsistencies cause problems during inventory preparation of greenhouse gases and pollutants. The first expert panel for data providers for agricultural data took place in 2022.

The Central Agricultural Testing and Controlling Institute (ÚKSÚP) reported inconsistencies in their data of utilisation of nitrogen fertilizers. Fertilization activity is detected on 90% of the agricultural land. Calculations are provided by the ÚKSÚP each year. 90% of data are collected electronically at the farm level and subsequently reported to the ŠÚ SR which reports data to FAOSTAT and EUROSTAT. Revision of data was done in 2022 submission, the data was harmonized with [EUROSTAT](#) database in partial years 2000 and 2010, where the inconsistencies were identified ([Table 5.6](#)).

The quality control comparison of nitrogen was done. Main inconsistencies between the FAOSTAT 2023 database and national inventory (**Table 5.6**) were identified huge inconsistency from 2000 to 2012 (*cursive*). Databases after 2013 are harmonised except for data from IFASTAT. IFASTAT data are different throughout the time-series (***cursive bold***). Different rounding is a common problem in all datasets (IFASTAT, FAOSTAT, and EUROSTAT). Consumption for the year 2021 was not available in the FAOSTAT and IFASTAT at the time of this exercise.

The number of livestock: The number of animals is the most important input parameter into the emissions inventory. The differences can be recognized in the methodological approach of data collection used by the FAOSTAT and by the ŠÚ SR. FAOSTAT grouped livestock in 12-months periods ending on 30<sup>th</sup> September each year. On the other hand, the ŠÚ SR provides annual national data on livestock by 31<sup>st</sup> December of a given year. The statistical survey is based on data collected from selected farms, animal census, by selected animals' categories, up to the regional level and finally up to national level. Therefore, the animal population 2021 in the FAOSTAT is different. In addition, detailed analysis of the data provides **Table 5.7**. It shows a shift in the timeline of goats (since 1994), sheep (since 1994), horses (since 1994) and swine (since 1994) (*cursive*). In 2019, FAOSTAT revised number of cattle (dairy and non-dairy cattle). The timeline is shifted since 2000 (*cursive*). Different allocation of cattle population (***cursive bold***) is visible in the years 1994 – 1997 (***cursive bold***). This inconsistency is caused by the different rules for distribution between dairy and non-dairy cattle. Revision of livestock mentioned above led to unification of cattle data between two databases in 2019, but different allocation of dairy and non-dairy and shift in the timeline were corrected partially. In addition, the FAO prepares its own estimates of broilers and layers number, annually. Therefore, the inconsistencies are visible in bold values. The revision of poultry population provided by the ŠÚ SR was not taken into consideration within the FAOSTAT. The ŠÚ SR as a partner of the EUROSTAT collects, processes and disseminates statistical data in line with the current national and EU legislation. Therefore, use of statistical data is considered as the most appropriate and accurate. However, comparison of data and methodologies with the independent data source FAOSTAT is useful tool for the QA activities. It can be assumed from this exercise that the activity data used in inventory of the Agriculture sector is in a good consistency and accuracy.

**Table 5.6: Comparison of fertilisers in different databases**

YEAR	SVK NIR 2025	FAOSTAT 2025	EUROSTAT 2025	IFASTAT 2025
	kg/year			
1993	64 852 000	64 883 000	NA	NA
1994	68 669 000	68 656 000	NA	68 700 000
1995	69 587 000	72 029 000	NA	72 000 000
1996	74 464 000	77 644 000	NA	77 600 000
1997	88 017 000	72 500 000	NA	72 500 000
1998	81 842 000	82 814 000	NA	82 800 000
1999	65 392 000	65 357 000	NA	65 400 000
2000	84 609 000	82 100 000	84 609 000	82 100 000
2001	102 423 000	81 345 000	102 423 000	85 000 000
2002	111 507 000	81 300 000	111 507 000	81 000 000
2003	97 727 000	79 911 000	97 727 000	93 000 000
2004	97 151 000	81 317 000	97 151 000	90 000 000
2005	99 760 000	78 681 000	99 760 000	90 000 000
2006	97 023 000	88 935 000	97 023 000	100 000 000
2007	113 298 000	87 737 000	113 298 000	105 000 000
2008	121 435 000	77 058 000	121 435 000	94 000 000
2009	96 334 000	86 873 000	96 334 000	83 000 000
2010	106 513 000	92 969 000	106 513 000	96 000 000

YEAR	SVK NIR 2025	FAOSTAT 2025	EUROSTAT 2025	IFASTAT 2025
	<i>kg/year</i>			
2011	120 555 000	101 004 000	120 555 000	113 000 000
2012	101 004 000	113 581 000	101 004 000	112 000 000
2013	113 581 390	113 581 000	113 581 000	118 000 000
2014	119 036 050	119 036 000	119 036 000	121 000 000
2015	114 773 000	114 773 000	114 773 000	133 300 000
2016	126 235 769	126 236 000	126 236 000	140 900 000
2017	122 541 152	122 541 152	122 541 000	125 900 000
2018	128 976 885	128 976 885	128 977 000	155 400 000
2019	128 532 971	128 532 970	128 533 000	138 200 000
2020	127 676 520	127 676 519	127 676 520	149 800 000
2021	127 494 597	127 494 600	127 495 000	151 900 000
2022	115 346 776	115 346 000	115 347 000	115 800 000
2023	107 607 314	NA	107 671 000	NA



**Table 5.7: Comparison of national data and the FAOSTAT in livestock population (heads) for the time series 1993 – 2023**

YEAR	DAIRY CATTLE		NON-DIARY CATTLE		GOATS		SHEEP		HORSES		SWINE		POULTRY	
	SVK NIR 2023	FAOSTAT 2023	SVK NIR 2023	FAOSTAT 2023	SVK NIR 2023	FAOSTAT 2023	SVK NIR 2023	FAOSTAT 2023	SVK NIR 2023	FAOSTAT 2023	SVK NIR 2023	FAOSTAT 2023	SVK NIR 2023	FAOSTAT 2023
heads														
1996	245 833	355 199	646 158	573 507	26 147	25 046	418 823	427 844	9 722	10 109	1 985 223	2 076 439	14 147 177	13 214 000
1997	299 614	335 381	503 784	556 610	26 778	26 147	417 337	418 823	9 533	9 722	1 809 868	1 985 223	14 221 713	13 985 000
1998	267 282	299 614	437 510	503 784	50 905	26 778	326 200	417 337	9 550	9 533	1 592 599	1 809 868	13 116 796	14 071 000
1999	250 974	283 895	414 081	420 897	51 075	50 905	340 346	326 199	9 342	9 550	1 562 106	1 592 599	12 247 440	13 027 000
2000	242 496	250 974	403 652	414 081	51 419	51 075	347 983	340 346	9 516	9 342	1 488 441	1 562 105	13 580 042	12 160 000
2001	230 379	242 496	394 811	403 652	40 386	51 419	316 302	347 983	7 883	9 516	1 517 291	1 488 441	15 590 404	13 482 000
2002	230 182	230 379	377 653	394 811	40 194	40 386	316 028	316 302	8 122	7 883	1 553 880	1 517 291	13 959 404	15 352 000
2003	214 467	230 182	378 715	377 653	39 225	40 194	325 521	316 028	8 114	8 122	1 443 013	1 553 880	14 216 798	13 817 000
2004	201 725	214 467	338 421	378 715	39 012	39 225	321 227	325 521	8 209	8 114	1 149 282	1 443 013	13 713 239	14 052 000
2005	198 580	201 725	329 309	338 421	39 566	39 012	320 487	321 227	8 328	8 209	1 108 265	1 149 282	14 084 079	13 565 000
2006	184 950	198 580	322 870	329 309	38 352	39 566	332 571	320 487	8 222	8 328	1 104 829	1 108 265	13 038 303	13 932 000
2007	180 207	184 950	321 610	322 870	37 873	38 352	347 179	332 571	8 017	8 222	951 934	1 104 829	12 880 124	12 882 000
2008	173 854	180 207	314 527	321 610	37 088	37 873	361 634	347 179	8 421	8 017	748 515	951 934	11 228 140	12 718 000
2009	162 504	173 854	309 461	314 527	35 686	37 088	376 978	361 634	7 199	8 421	740 862	748 515	13 583 284	11 081 000
2010	159 260	162 504	307 865	309 461	35 292	35 686	394 175	376 978	7 111	7 199	687 260	740 862	12 991 916	13 438 000
2011	154 105	159 260	309 253	307 865	34 053	35 292	393 927	394 175	6 937	7 111	580 393	687 260	11 375 603	12 846 000
2012	150 272	154 105	320 819	309 253	34 823	34 053	409 569	393 927	7 249	6 937	631 464	580 393	11 849 818	11 252 000
2013	144 875	150 272	322 945	320 819	35 457	34 823	399 908	409 569	7 161	7 249	637 167	631 464	10 968 918	11 693 000
2014	143 083	144 875	322 460	322 945	35 178	35 457	391 151	399 908	6 828	7 161	641 827	637 167	12 494 074	10 786 000
2015	139 229	143 083	318 357	322 460	36 324	35 178	381 724	391 151	6 866	6 828	633 116	641 827	12 836 224	13 084 000
2016	132 610	139 229	313 502	318 357	36 355	36 324	368 896	381 724	6 407	6 866	585 843	633 116	12 130 501	12 057 000
2017	129 863	132 610	309 963	313 502	37 067	36 355	365 344	368 896	6 145	6 407	614 384	585 843	13 353 837	13 133 000
2018	127 871	129 863	310 984	309 963	36 907	37 067	351 122	365 344	7 102	6 145	627 022	614 384	14 056 914	13 354 000
2019	125 848	125 850	306 405	306 405	35 594	35 590	320 555	320 560	6 960	6 960	589 228	589 230	13 131 941	13 132 000
2020	122 049	122 050	320 240	320 240	10 589	35 600	294 252	294 252	6 099	6 857	538 310	538 310	10 603 624	10 572 000
2021	120 068	120 070	314 021	31 402	10 434	32 000	290 918	290 918	6 738	6 044	453 076	453 080	10 364 509	10 297 703
2022	116 910	116 910	316 265	31 627	11 008	20 500	301 131	301 130	7 044	NA	380 895	380 900	9 340 713	9 275 000
2023	114 896	NA	314 825	NA	10 719	NA	289 849	NA	7 367	NA	403 037	NA	9 669 376	NA

## 5.4. Category-specific Recalculations

Recalculations developed in the Agriculture sector were provided and implemented in line with the strategy for improvement and prioritisation in 2024, reflecting recommendations received during previous reviews and the sectoral expert’s proposals. **Table 5.8** shows an overview of these recalculations and corrections implemented in 2025 submission.


**Table 5.8: Overview of recalculations and implemented improvements in the Agriculture sector**

NUMBER	CATEGORY	DESCRIPTION	REFERENCE
1.	3.A Enteric fermentation 3.B.4 Manure management	Addition of new emission sources from rabbits in the categories 3.A.4.h.i. Rabbit and CH4 and N2O from 3.B.4.h.i Rabbits.	5.2
2.	3.B.1.Manure management	Non-dairy cattle: Fixed inconsistency for the activity data in the share of pasture on total AWMS for the animal subcategory (heifers) of Non-dairy cattle.	5.4
3.	3.B.3.Manure management	Update of the method for the calculation of nitrogen excretion rate for the swine categories.	5.4
4.	3.D Agricultural soils	Recalculation based on the recommendation during the review on identified irregularities between nitrogen volatilized as NH <sub>3</sub> and NO <sub>x</sub> . It will have impact in the 3.D.1.b Organic N fertilizers and 3.D.2 Indirect N <sub>2</sub> O emissions from managed soils categories.	5.2
5.	3.D Agricultural soils	Recalculation based on the implementation of updated data on the N content in different types of fertilizers included in the category 3.D.1.b.iii Other organic fertilizers.	5.4

The overall impact of recalculations developed in the Agriculture sector resulted in 9.33% (184.5 kt of CO<sub>2</sub> eq) increase of emissions in 2025 compared to previous submission (2024). The Agriculture sector is specific sector regarding the recalculations process. Change in one category caused changes also in other categories across sector, due to methodology based on nitrogen and methane balance.


**Figure 5.5** shows overall trend of recalculated emissions and comparison of 2024 and 2025 submissions.

**Figure 5.5: Comparison of 2024 and 2025 submissions in the Agriculture sector (in Gg of CO<sub>2</sub> eq.)**




**Figure 5.6** shows overall trend of recalculated emissions and comparison of 2024 and 2025 submissions for 3.A category.

**Figure 5.6:** Comparison of 2024 and 2025 submissions for 3.A category (in Gg of CO<sub>2</sub> eq.)




**Figure 5.7** shows overall trend of recalculated emissions and comparison of 2024 and 2025 submissions for 3.B category.

**Figure 5.7:** Comparison of 2024 and 2025 submissions for 3.B category (in Gg of CO<sub>2</sub> eq.)



**Figure 5.8** shows overall trend of recalculated emissions and comparison of 2024 and 2025 submissions for 3.D category.

**Figure 5.8:** Comparison of 2024 and 2025 submissions for 3.D category (in Gg of CO<sub>2</sub> eq.)



During interinstitutional QA/QC process with Central Controlling and Testing Institute in Agriculture (ÚKSÚP) responsible for providing data on fertilizer use were identified inconsistencies in N content of different types of fertilizers accounted in the category 3.D.1.b.iii Other organic fertilizers.

More information on the method of calculation is available in the [Chapter 5.12.4](#). Data with corrected values are presented in the [Table 5.9](#).

**Table 5.9:** Revised N contents of other organic fertilizers applied 1990 – 2023

Year	Fugate		Compost		Natural harmony		Straw		Vitahum		Green fertilizers	
	OW	NC	OW	NC	OW	NC	OW	NC	OW	NC	OW	NC
	<i>t</i>											
1990	NO	NO	33 429.50	234.01	NO	NO	NO	NO	28 290.00	198.03	12 013.24	60.07
1991	NO	NO	34 303.00	336.17	NO	NO	NO	NO	26 501.00	185.51	11 752.48	58.76
1992	NO	NO	35 176.50	246.24	NO	NO	NO	NO	24 713.00	172.99	11 492.72	57.46
1993	NO	NO	36 050.00	252.35	NO	NO	NO	NO	22 924.00	160.47	11 230.96	56.15
1994	NO	NO	36 923.50	361.85	NO	NO	NO	NO	21 136.00	147.95	10 970.02	54.85
1995	NO	NO	37 797.00	264.58	NO	NO	NO	NO	19 348.00	135.44	10 709.44	53.55
1996	NO	NO	38 670.50	270.69	NO	NO	NO	NO	17 559.00	122.91	10 448.68	52.24
1997	NO	NO	39 544.00	387.53	NO	NO	NO	NO	15 771.00	110.4	10 187.92	50.94
1998	NO	NO	40 417.50	282.92	NO	NO	NO	NO	13 982.00	97.87	9 927.16	49.64
1999	NO	NO	41 291.00	289.04	NO	NO	NO	NO	12 194.00	85.36	9 666.40	48.33
2000	NO	NO	74 922.60	524	NO	NO	NO	NO	50 641.00	354.3	10 245.00	51.23
2001	NO	NO	40 885.09	286.2	NO	NO	NO	NO	54 338.00	380.37	18 285.00	91.43
2002	NO	NO	36 422.30	254.96	NO	NO	NO	NO	42 810.00	299.67	10 920.00	54.6
2003	NO	NO	34 225.20	239.58	NO	NO	NO	NO	9 321.00	65.25	6 206.00	31.03
2004	NO	NO	42 904.07	300.33	NO	NO	NO	NO	2 845.00	19.92	18 989.50	94.95
2005	NO	NO	7 005.62	49.04	NO	NO	NO	NO	3 552.00	24.86	5 905.00	29.53
2006	NO	NO	13 877.53	97.14	NO	NO	NO	NO	10 828.00	75.78	7 006.00	35.03

Year	Fugate		Compost		Natural harmony		Straw		Vitahum		Green fertilizers	
	OW	NC	OW	NC	OW	NC	OW	NC	OW	NC	OW	NC
	<i>t</i>											
2007	NO	NO	21 761.87	152.33	NO	NO	8 867.66	72.71	8 758.00	61.31	3 540.44	17.7
2008	NO	NO	21 317.38	149.22	NO	NO	90 976.84	746.01	7 185.00	50.3	13 533.88	67.67
2009	NO	NO	25 363.76	177.55	NO	NO	68 636.58	568.82	195	1.37	16 642.30	83.21
2010	NO	NO	40 096.76	280.68	NO	NO	36 773.65	301.54	4 999.00	34.99	11 955.63	59.78
2011	NO	NO	50 582.67	354.08	11 107.00	266.57	66 704.06	546.97	2 261.00	15.83	25 836.98	129.18
2012	108 181.00	995	18 291.36	128.04	205.2	4.92	25 019.69	205.16	NO	NO	1 401.27	7.01
2013	301 580.00	2 775.00	63 145.17	442.02	395.41	9.47	30 697.63	251.72	500	3.5	2 546.69	12.73
2014	382 111.00	3 515.00	85 906.56	601.35	17 819.00	427.66	40 911.78	335.48	NO	NO	6 374.80	31.87
2015	543 489.00	5 000.00	90 967.08	636.77	46 392.00	1 089.00	26 554.08	217.74	1 015.00	7.11	4 036.14	20.18
2016	391 789.00	2 651.00	46 700.66	318.06	48 703.00	1 137.00	NO	NO	NO	NO	NO	NO
2017	732 884.00	2 842.00	46 649.05	326.54	25 389.00	583	NO	NO	17 928.00	125.5	NO	NO
2018	720 233.00	2 790.00	43 256.57	410.75	30 791.00	731	NO	NO	NO	NO	NO	NO
2019	776 427.00	3 057.00	37 617.52	300.05	41 518.00	994	NO	NO	4 500.00	31.5	NO	NO
2020	800 393.00	2 936.00	43 556.51	249.87	NO	NO	NO	NO	NO	NO	NO	NO
2021	796 945.00	3 347.00	60 046.78	401.39	NO	NO	NO	NO	NO	NO	NO	NO
2022	793 655.00	2 812.00	56 181.00	375	NO	NO	NO	NO	NO	NO	NO	NO
2023	897 094.64	3 090.61	20 499.75	169.84	NO	NO	NO	NO	NO	NO	NO	NO

### Ad 3. Nitrogen excretion for the swine categories

Changes in the method of the NEX estimation were motivated by the effort to apply Tier 2 method revised last year to implement 2019 RF IPCC in the time series consistently. NEX calculation using feed composition data (one off expert judgement in 2017) was replaced by the quantification of NEX based on the national feeding system Petrikovič et al. (2002): *Nutrient requirements of pigs* for the entire time series. In this system, feed intake and crude protein content of the feed is estimated to cover animal requirements for production and maintenance. Animal requirements needs to be determined based on average weight of animal in the category or average daily weight gain (for Market swine). In the current absence of these data for the time series (back to 1990), this approach resulted in the flat value of NEX.

**Table 5.10** depicts all animal subcategories and their parameters used for the estimation of the NEX for the entire timeseries and all eight regions.

**Table 5.10: Swine subcategories and parameters of N balance in the time series**

Breeding swine subcategories	CP	N-intake	N <sub>EX</sub>	Market swine subcategories - FATTENING	CP	N-intake	N <sub>EX</sub>
	(%)	kg N animal/day	kg N/animal/year		(%)	kg N animal/day	kg N/animal/year
SOWS	14.8	0.07	21.36	PIGS UP to 20 kg	26.3	0.026	4.84
GILTS PREGNANT	14.7	0.05	12.5	PIGS 21-50 kg	24.7	0.045	9.43
GILTS UNPREGNANT	15.0	0.06	15.1	PIGS 50 -80 KG	22.16	0.061	13.8
HOGS	32.7	0.11	27.1	PIGS 80 – 110 KG	19.89	0.071	17.7
PIGS 21 – 50 KG	20.6	0.037	8.07	PIGS FROM 110 KG	18.3	0.073	18.7

The limitation in the data is planned to be tackled by ongoing investment in research project and capacity building exercise.

## 5.5. National Circumstances and Time-series Consistency

Slovak farmers have been adapted to changes in agriculture after 1990. They invested in the development of their farms to avoid the bankruptcy and to be self-competitive in this sector. The EÚ policy supported the used tools as the base of transformation. The EÚ policy and measures were transformed into the Slovak legal system. Farmers had to follow new strict criteria like changing of housing systems, a decrease of pasture time, new storage capacity for organic waste, which was supported by the Decree No 389/2005 Coll. and Nitrates Directive. These measures are well advanced and copy the practices used in the Western European countries. Therefore, default parameters for the Western Europe are used in inventory. The most significant animals in regard of emissions in Slovakia are cattle and swine.

Cattle breeding in the Slovak Republic is comparable with the Western European countries, which is documented by a high milk yield of dairy cattle and high daily weight gains of non-dairy cattle. To maintain a high milk yield and high daily gains, food rich on proteins and cereals is important. Dairy cows in three Slovak regions (Bratislava, Trnava and Nitra) produce 29-26 litres/day. In other regions, milk productivity is 18-23 litres/day. Lower milk production relates to feeding. In this case, pasture is included in the feeding ratio. It is typical for semi-intensive farming in regions Košice, Prešov, Banská Bystrica or Žilina. These circumstances are documented on [Figures 5.9](#) and [5.10](#). Highly productive dairy cows (milked 25 -30 litres/day) need to be fed by approximately 8 kg of cereals with excellent digestibility and high nutrition. Annual increase in milk productivity is the evidence of increasing productivity of animal production. Balanced and sustainable farming in Slovakia has an impact on the high value of AGEI (304 MJ/head/day) ([Table 5.36](#)).


**Table 5.11:** The comparison of the Slovak milk yield with other regions in 2023

DAIRY COWS	SLOVAKIA <sup>1</sup>	WESTERN EUROPE <sup>2</sup> (AVERAGE)	EASTERN EUROPE <sup>2</sup> (AVERAGE)	NORTH AMERICA <sup>6</sup> (AVERAGE)
	<i>kg/year/head</i>			
Milk yield	8 408	7 255	5 478	10 863

<sup>1</sup> The animal production, sales of primary production and crop balance (in Slovak) [www.statistics.sk](http://www.statistics.sk)


<sup>2</sup> Producing Animals (Slaughtered), Milk Production <http://www.fao.org/faostat/en/#data/QL>

**Figure 5.9:** Trend in average gross energy intake (MJ/day) in different Slovak regions




The number of dairy cows decreased according to data from the ŠÚ SR by 71% in 2022 compared to 1990 (**Figure 5.10**). Milk production increased up to 228% in 2022 (**Figure 5.11**) compared to the 1990 and by 3.8% compared to previous year despite the continuously decreasing number of the dairy cows. The main reason of this trend is the increase in an average performance. The high-performance average is the result of good animal husbandry, breeding conditions, new synergy with technologies and animal genetics. All factors contribute together to achieving milk yields of up to 10 000 kg of milk per head per year.

**Figure 5.10:** Correlation of milk production (kg/day/head) and nitrogen excretion rate (kg N/year/head)




**Figure 5.11: Trend in dairy cattle population and dairy milk production (kg/head/day)**



The pig farming system in the Slovak Republic is divided into two types - breeding and fattening pigs. Breeding pigs are bred for reproduction purposes. Fattening pigs are bred mainly for the production of pork meat and fat. Pigs are housed in the Slovak conditions for the whole year. Housing technology and diet can significantly affect the production of greenhouse gases. Stall conditions can be very variable. Pigs are bred in intensive farming on rosette floors, which is one of the low emission technics. Another part of pigs, mainly in semi-intensive farming, are reared on straw. Deep bedding is used mostly at micro and small farms. Diet has a significant impact on emissions production. The main component of the feeding is cereals (barley, triticale, wheat about 80-90%). Complementary feed ingredients are soybean meal, rapeseed meal, and brewing malt. The resultant feeding rations have a high nutritional value and are easily digestible ([Figure 5.12](#)). After 1990, the digestibility of feeding dose increased significantly due to the increase of cereals, vitamins, dietary fibre, crude proteins and amino acids. These changes affect the increase in pig performance. In 2021, visible increase of digestibility of feeding doses occurred. This value was estimated by VÚŽV and correlated with increase of pig performance in that year. The opposite trend is visible in the last 4 years mainly in breeding pigs. The decrease in crude proteins, cereals had an impact on the decrease of monitored parameters. Pig breeding in Slovakia has problems mainly due to risk of persistent morbidity - African swine fever and other economic reasons, which lead to decreasing numbers of pigs.



**Figure 5.12:** Trend of feed digestibility, nitrogen excretion rate and gross energy intake of swine in the Slovak Republic



## 5.6. Uncertainties

Uncertainty estimates of emissions were performed using tier 2 approach based on the Monte Carlo simulation. The simulation is done using the Python language. The following chapter gives preliminary overview of uncertainty estimates for the CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> emissions from Agriculture for the year 2020. These results have not been officially published yet.

Monte Carlo simulations are used in the modelling of the probability of different outcomes in a process that cannot easily be predicted due to the intervention of random variables. It is a technique using for understanding of the impact of risk and uncertainty in prediction and forecasting models. The Monte Carlo analysis was prepared on regional level. The uncertainties of livestock population for 2020 are presented in [Table 5.12](#). Uncertainties were estimated according to an assessment of the SHMÚ team while no information was provided by the ŠÚ SR in this area. The uncertainty analysis was performed by the coefficient of variation. The coefficient of variation is a statistical measure of the dispersion of data points in a data series around the mean. The coefficient of variation represents the ratio of the standard deviation to the mean, and it is a useful statistic value for comparing the degree of variation from one data series to another, even if the means are drastically different from one another. The overall weighted mean of the uncertainties in the livestock population is  $\pm 6.09\%$ .

**Table 5.12:** Uncertainty of animal population data for 2023

CATEGORY	UNIT	AGREGATED UNCERTAINTY OF NUMBER OF LIVESTOCK
Dairy cattle	head	$\pm 2.77\%$
Non-dairy cattle	heads	$\pm 1.94\%$
Sheep	heads	$\pm 2.08\%$
Goats	heads	$\pm 12.94\%$
Horses	heads	$\pm 2.48\%$
Swine	heads	$\pm 3.94\%$
Poultry	heads	$\pm 6.83\%$
Overall (weighted mean)	heads	$\pm 6.09\%$

The highest uncertainty increment to the total uncertainty of Agriculture sector represents N<sub>2</sub>O emissions from agricultural soils, particularly uncertainties of used emission factors. The overall sectoral uncertainty is strongly influenced by uncertainties and distribution among the EF<sub>1</sub>, EF<sub>4</sub> and EF<sub>5</sub>

emission factors. However, the partial uncertainties on category level were calculated, overall uncertainty of the sector is still not estimated and will be provided in next submission.

**Enteric Fermentation (CRF 3.A):** Results of the Monte Carlo simulation for methane emissions in the category 3.A – Enteric Fermentation were estimated at 35.52 Gg of CH<sub>4</sub> (38.65 Gg of CH<sub>4</sub> were estimated in inventory) with uncertainty (-15%, +15%), which represent 95% confidence interval in 2020. A probability distribution function for category 3.A is shown on [Figure 5.10](#). In this uncertainty simulation, symmetric confidence intervals is parametrized by Gaussian normal probability density function.

Uncertainties of EFs (CH<sub>4</sub>) from enteric fermentation for dairy cattle, non-dairy cattle and sheep were based on uncertainties of milk production, wool production and weight listed in [Tables 5.13-5.16](#). Data on milk production, weight of animals is readily available while the GE is checked against cattle feeding requirements arising from the biology of ruminants (e.g. ratio of crude protein, dry matter intake and proportion of silage in the diet).

**Table 5.13: Uncertainties of parameters used in enteric fermentation**

PARAMETER*	UNIT	UNCERTAINTY						
		Dairy cows	Calves	Heifers un-pregnant milk breed	Heifers pregnant milk breed	Fattening	Oxen	Breeding bulls
Body weight	%	±10	±10	±10	±10	±10	±10	±10
Milk yield	%	±2	-	-	-	-	-	-
DE of feed	%	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96
Ym	%	±3	±3	±3	±3	±3	±3	±3
Maintenance NE <sub>m</sub>	MJ/day	±2.46	±0.021	± 0.314	±0.243	±0.110	±9.66	±5.228
Activity NE <sub>a</sub>	MJ/day	±0.161		±0.408	±0.165	-	-	±0.838
Lactation NE <sub>l</sub>	MJ/day	±2.869	-	-	-	-	-	-
Work		-	-	-	-	-	±12.94	-
Growth NE <sub>g</sub>	MJ/day	-	±0.000	±1.570	±1.147	±0.562	-	-
Pregnancy NE <sub>p</sub>	MJ/day	±0.321	-	-	±0.165	-	-	-
REM		±0.019	±0.030	±0.057	±0.034	±0.019	±0.18	±0.056
REG		±0.011	±0.019	±0.032	±0.020	±0.011	±0.11	±0.031
Gross energy	MJ/head/day	±16.808	±1.812	±11.344	±9.951	±3.351	±116.613	±32.752
EFs	kg/head/year	±45.747	±3.915	±13.010	±18.211	±11.469	±62.545	±31.818

**Table 5.14: Uncertainties of parameters used in enteric fermentation**

PARAMETER*	UNIT	UNCERTAINTY						
		Suckling cows	Calves	Heifers un-pregnant milk breed	Heifers pregnant milk breed	Fattening	Oxen	Breeding bulls
Body weight	%	±10	±25	±25	±25	±25	±25	±25
Milk yield	%	±2	-	-	-	-	-	-
DE of feed	%	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96
Ym	%	±3	±3	±3	±3	±3	±3	±3
Maintenance NE <sub>m</sub>	MJ/day	±0.012	±0.052	±0.054	±0.109	±0.110	±7.198	±5.270
Activity NE <sub>a</sub>	MJ/day	±0.419	±0.243	±0.673	±0.922	-	-	-
Lactation NE <sub>l</sub>	MJ/day	±0.964	-	-	-	-	-	-
Work		-	-	-	-	-	±7.088	-
Growth NE <sub>g</sub>	MJ/day		±0.0	±1.242	±1.433	±1.848		-
Pregnancy NE <sub>p</sub>	MJ/day	±0.209			±0.465	-	-	-
REM		±0.023	±0.061	±0.055	±0.054	±0.058	±0.097	±0.083
REG		±0.013	±0.036	±0.031	±0.029	±0.032	±0.054	±0.046
Gross energy	MJ/head/day	±12.223	±5.065	±13.151	±20.077	±12.006	±68.701	±38.873

PARAMETER*	UNIT	UNCERTAINTY						
		Suckling cows	Calves	Heifers un-pregnant milk breed	Heifers pregnant milk breed	Fattening	Oxen	Breeding bulls
EFs	kg/head/year	±33.153	±5.715	±15.52	±24.317	±13.764	±47.698	±31.634

**Table 5.15:** Uncertainties of emission factors in non-key categories in enteric fermentation


Year	Uncertainty	Animals	Emission factor
2020	4.96%	Swine	1.5 kg/head
2020	4.55%	Horse	18 kg/head
2020	9.60%	Goats	5 kg/head

**Table 5.16:** Uncertainties of parameters calculated in enteric fermentation

PARAMETER*	UNIT	DAIRY SHEEP				BEEFSHEEP			
		A	B	C	D	E	F	G	H
DE of feed	%	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96	±4.96
Y <sub>m</sub>	%	±3	±3	±3	±3	±3	±3	±3	±3
Maintenance NE <sub>m</sub>	MJ/day	±0.169	±0.630	±0.518	±1.092	±0.169	±0.578	±0.544	±1.180
Activity NE <sub>a</sub>	MJ/day	±0.064	±0.168	±0.176	±0.154	±0.115	±0.201	±0.273	±0.167
Lactation NE <sub>l</sub>	MJ/day	±0.071				±0.111			
Wool production Ne <sub>wool</sub>	MJ/day	±0.005	±0.006	±0.009	±0.006	±0.011	±0.008	±0.021	±0.011
Growth NE <sub>g</sub>	MJ/day		±0.078	±0.227			±0.176	±0.226	
Pregnancy NE <sub>p</sub>	MJ/day	±0.229		±0.645		±0.425		±1.047	
REM		±0.019	±0.032	±0.033	±0.020	±0.037	±0.057	±0.072	±0.038
REG		±0.010	±0.016	±0.016	±0.011	±0.019	±0.029	±0.036	±0.019
Gross energy	MJ/head/day	±2.138	±2.816	±5.166	±4.564	±4.146	±4.036	±8.774	±5.425
EFs	kg/head/year	±5.847	±2.584	±6.518	±5.767	±6.644	±3.676	±7.389	±5.170

A: Mature ewes, B: Growing lambs, C: Growing lambs pregnant, D: Other mature sheep, E: Mature ewes, F: Growing lambs, G: Growing lambs pregnant, H: Other mature sheep, \* weighted average

**Figure 5.13:** Probability distribution function for the category 3.A (x-axis in Gg of CH<sub>4</sub>)



**Manure Management (CRF 3.B.1):** Results of the Monte Carlo simulation for methane emissions in the category 3.B.1 – Manure Management were calculated on the value 3.04 Gg of CH<sub>4</sub> (3.06 Gg of CH<sub>4</sub> were estimated in inventory) with uncertainty (-14.91%, +14.91%) which represent 95% confidence interval in 2020. A probability distribution function for category 3.B.1 is shown on **Figure 5.14**. In this uncertainty simulation, symmetric confidence intervals is parametrized by Gaussian normal probability density function.

Uncertainties of EFs (CH<sub>4</sub>) from manure management for dairy cattle, non-dairy cattle and sheep were based on uncertainties of storage of solid and liquid manure management systems from breeding animals listed in **Tables 5.17-5.19**. Data on storage systems and number of livestock is readily available.

**Table 5.17: Uncertainties of parameters used in manure management for cattle and sheep in 2020**

PARAMETERS	UNIT	DAIRY CATTLE	NON-DAIRY CATTLE	MATURE EWES	GROWING LAMBS	OTHER MATURE SHEEP
B <sub>o</sub> *	%	±15%	±15%	±15%	±15%	±15%
Ash content	%	±20%	±20%	±20%	±20%	±20%

**Table 5.18: Uncertainties of parameters used in manure management for market swine in 2020**

PARAMETERS	UNIT	A	B	C	D	E
B <sub>o</sub> *	%	±15%	±15%	±15%	±15%	±15%
Ash content	%	±20%	±20%	±20%	±20%	±20%


A: Fattening pigs up to 20 kg; B: Fattening pigs 21-50 kg; C: Fattening pigs 50-80 kg, D: Fattening pigs 80-110 kg; E: Fattening pigs over 110 kg

**Table 5.19: Uncertainties of parameters used in manure management for breeding swine in 2020**

PARAMETERS	UNIT	A	B	C	D	E	F
B <sub>o</sub> *	%	±15%	±15%	±15%	±15%	±15%	±15%
Ash content	%	±20%	±20%	±20%	±20%	±20%	±20%

A: Sows, B: Gilts non-pregnant, C: Gilts pregnant, D: Hogs; E: Piglets up to 20 kg; F: Piglets 21-50 kg; \*Bo for Western Europe was chosen

**Figure 5.14: Probability distribution function for the category 3.B.1 (x-axis in Gg of CH<sub>4</sub>)**



**Table 5.20: Uncertainties of parameters calculated in manure management for sheep in 2020**

ANIMAL		EFs	VSs
		kg VS/day	kg/head
DAIRY SHEEP	Mature ewes	±0.032	±0.046
	Growing lambs	±0.044	±0.064
	Growing lambs (pregnant)	±0.081	±0.117
	Other mature sheep	±0.096	±0.103
BEEF SHEEP	Mature ewes	±0.060	±0.089
	Growing lambs	±0.062	±0.091
	Growing lambs (pregnant)	±0.134	±0.199
	Other mature sheep	±0.115	±0.124

**Table 5.21: Uncertainties of parameters calculated in manure management for cattle in 2020**

ANIMAL		VSs	EFs
		kg/head	kg VS/day
MILK TYPE	Dairy cows	±0.258	±0.501
	Calves in 6. month	±0.021	±0.018
	Heifers	±0.193	±0.167
	Heifers (pregnant)	±0.167	±0.145
	Fattening	±0.054	±0.067
	Oxen	±1.643	±1.446
	Breeding bull	±0.549	±0.424
BEEF TYPE	Suckler cows	±0.239	±0.153
	Calves in 6. month	±0.070	±0.049
	Heifer	±0.252	±0.161
	Heifer (pregnant)	±0.253	±0.396
	Fattening	±0.228	±0.362
	Oxen	±1.110	±1.260
	Breeding bull	±0.684	±0.528


**Table 5.22: Uncertainties of parameters calculated in manure management for swine in 2020**

ANIMAL	VSs	GE	ME	EFs
	kg/head	MJ/day	MJ/day	kg/head
Sows	±0.279	±0.279	±0.28	±5.709
Gilts non-pregnant	±0.314	±0.314	±0.314	±4.872
Gilts pregnant	±0.390	±0.390	±0.390	±3.924
Hogs	±0.390	±0.390	±0.390	±3.924
Piglets 20 kg	±1.258	±1.258	±1.180	±1.217
Piglets 21-50kg	±0.670	±0.670	±0.649	±2.287
Fattening to 20 kg	±1.178	±1.178	±0.062	±0.923
Fattening to 21-50 kg	±0.649	±0.649	±0.649	±1.674
Fattening to 50-80 kg	±0.445	±0.445	±0.445	±2.453
Fattening to 80-100 kg	±0.355	±0.355	±0.355	±4.232
Fattening from 110 kg	±0.317	±0.317	±0.317	±4.721

**Manure Management (CRF 3.B.2):** Results of the Monte Carlo simulation for N<sub>2</sub>O emissions in the category 3.B.2 – Manure Management were calculated on the value 0.34 Gg of N<sub>2</sub>O (0.24 Gg of N<sub>2</sub>O were estimated in inventory) with uncertainty (±248.3), which represent 95% confidence interval in 2020. A probability distribution function for category 3.B.2 is shown on [Figure 5.15](#). In this uncertainty simulation, symmetric confidence intervals is parametrized by Gaussian normal probability density function.

Uncertainties of N<sub>2</sub>O emissions relating to the N excretion for cattle are  $\pm 0.015$  Gg and for swine  $\pm 0.0038$  Gg. Uncertainties of other animals' species as poultry are  $\pm 0.020$  Gg. The uncertainty of the manure management system usage (MST, S) are  $\pm 25\%$ , what is in accordance with the default value provided by 2006 IPCC Guidelines. The uncertainty of the EFs is  $\pm 2.6\%$ , therefore the lower combined uncertainty ( $\pm 12.17\%$ ) of the activity data and emission factor from manure management are estimated.

**Figure 5.15:** Probability distribution function for the category 3.B.2 (x-axis in Gg of N<sub>2</sub>O)




**Direct N<sub>2</sub>O Emissions from Managed Soils (CRF 3.D.1):** Results of the Monte Carlo simulation for N<sub>2</sub>O emissions in the category 3.D.1 – Direct N<sub>2</sub>O Emissions from Managed Soils were 3.986 Gg of N<sub>2</sub>O (3.73 Gg of N<sub>2</sub>O were estimated in inventory) with uncertainty (-7.90%, +7.90%), which represent 95% confidence interval in 2020. A probability distribution function for category 3.D.1 is shown on [Figure 5.16](#). In this uncertainty simulation, symmetric confidence intervals is parametrized by Gaussian normal probability density function.

The overall uncertainty of N<sub>2</sub>O emissions from agricultural soils was estimated based on information of nitrogen inputs into the soils, used emission factors and their uncertainties ([Table 5.23](#)). During the preparation of overall uncertainty, the lack of information on the uncertainty of activity data was identified by the ŠÚ SR and UKSÚP. The uncertainty analysis was performed by the coefficient of variation. Information on animal waste management systems and number of livestock were taken into consideration in emission estimation and uncertainties. The resulted uncertainty for activity data for category 3.D is  $\pm 9.50\%$  and the uncertainty in the emission factor is  $\pm 6.34\%$ .

**Table 5.23:** Uncertainties of activity data in 3.D - Agricultural Soils

N <sub>2</sub> O DIRECT/INDIRECT EMISSION FROM MANAGED SOILS	UNITS	UNCERTAINTIES
Animal Manure Applied to Soils	%	$\pm 39.32$
Urine and Dung deposited by grazing animals	%	$\pm 5.27$
Crop residues	%	$\pm 88.55$
Mineralization or Immobilization Associated with Loss or Gain of Soil Organic Matter	%	$\pm 92.44$
Inorganic N Fertilizers	%	$\pm 91.51$
Atmospheric Deposition	%	$\pm 131.40$
Nitrogen Leaching and Run-off	%	$\pm 109.50$

**Figure 5.16:** Probability distribution function for 3.D.1 (x-axis in Gg of N<sub>2</sub>O)




**Indirect N<sub>2</sub>O Emissions from managed soils (CRF 3.D.2):** Results of the Monte Carlo simulation for N<sub>2</sub>O emissions in the category 3.D.2 – Indirect N<sub>2</sub>O Emissions from Managed Soils were calculated on the value 0.48 Gg of N<sub>2</sub>O (0.64 Gg of N<sub>2</sub>O were estimated in inventory) with uncertainty (-103.3%, +103.3%), which represent 95% confidence interval in 2020.

The uncertainty in 3.D.2.1 category of indirect N<sub>2</sub>O emissions was estimated based on partial uncertainties in emission factors. These uncertainties were combined with the uncertainties in the Fra<sub>C<sub>GAS</sub>F</sub> (0.03-0.3) and Fra<sub>C<sub>GAS</sub>M</sub> (0.05-0.5). Uncertainties of emission factors in indirect N<sub>2</sub>O emissions from soils were calculated at a level of ±133.24%, which represent 95% confidence interval.


**Liming (3.G):** Results of the Monte Carlo simulation for CO<sub>2</sub> emissions in the category 3.G – Liming were 8.45 Gg of CO<sub>2</sub> (8.45 Gg of CO<sub>2</sub> were estimated in inventory) with uncertainty (-3.04%, +3.04%), which represent 95% confidence interval in 2020. A probability distribution function for the category 3.G is shown on **Figure 5.17**. In this uncertainty simulation, symmetric confidence intervals is parametrized by Gaussian normal probability density function.

**Figure 5.17:** Probability distribution function for the category 3.G (x-axis in Gg of CO<sub>2</sub>)



**Urea Application (3.H):** Results of the Monte Carlo simulation for CO<sub>2</sub> emissions in the category 3.H – Urea Application were calculated on the value 63.67 Gg of CO<sub>2</sub> (63.63 Gg of CO<sub>2</sub> were estimated in inventory) with uncertainty (-3.93%, +3.93%), which represent 95% confidence interval in 2020. A probability distribution function for category 3.H is shown on **Figure 5.18**. In this uncertainty simulation, symmetric confidence intervals is parametrized by Gaussian normal probability density function.

**Figure 5.18:** Probability distribution function for the category 3.H (x-axis in Gg of CO<sub>2</sub>)



**Agriculture sector:** Preliminary summary results of calculated uncertainties across categories in the sector are provided in the following table:

**Table 5.24:** Uncertainties of activity data, emission factors and emissions for key and particularly significant categories in agriculture identified by Monte Carlo approach

3 AGRICULTURE	GHG	UNCERTAINTY OF ACTIVITY DATA	UNCERTAINTY OF EMISSION FACTOR	UNCERTAINTY OF EMISSIONS
3.A Enteric Fermentation	CH <sub>4</sub>	±13.10%	±14.91%	±19.85%
3.B.1 Manure Management	CH <sub>4</sub>	±6.50%	±9.41%	±11.43%
3.B.2 Manure Management	N <sub>2</sub> O	±6.50%	±248.03%	±248.11%
3.D.1 Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	±59.3%7	±36.22%	±69.54%
3.D.2 Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	±77.10%	±103.30%	±128.90%
3.G Liming	CO <sub>2</sub>	±3.04%	±50.00%	±50.09%
3.H Urea Application	CO <sub>2</sub>	±3.93%	±50.00%	±50.15%

## 5.7. Enteric Fermentation (CRF 3.A)

*Emitted gas: CH<sub>4</sub>*

*Methods: T1 and T2*

*Emissions factors: D, CS*

*Key sources: Yes*

*Significant subcategories: Cattle*

The cattle is the most important producer of methane due to its digestive tract, weight and a relatively high number compared to other population of livestock in the Slovak Republic. Therefore, the trends in total CH<sub>4</sub> emissions reflect a number of animals and milk yield in this category. The number of dairy



cattle further decreased in 2023 in comparison with 2022 (-2%), non-dairy cattle decreased in 2023 in comparison with 2022 (-0,46%). Except for the population of domestic livestock, the amount of emitted methane is influenced by other parameters like age or weight of animals, amount of feed dry matter intake and its quality and the consumption of energy for basal metabolisms, milk production per day and fat content, the average amount of physical activity performed, wool growth and feed digestibility.


The decline in the number of all species of livestock since the base year is significant mostly in swine and cattle categories in the Slovak Republic. The highest decrease was observed in swine (-84%), cattle (-73%) and sheep (-52%) categories compared to the base year. The number of swine (-6%), horses (5%), poultry (4%) were increased compare to the previously year. Number of cattle (-0.8%), sheep (-4%), goats (-3%) decrease in 2023 compared to 2022.

Methane emissions from enteric fermentation have the major share on GHG emissions in agriculture. The cattle represent 89% of these emissions; from that dairy cattle 39% share. Other categories of domestic livestock provide 11% of emissions. Intensification of animal husbandry also increased methane emissions. Methane emissions from enteric fermentation of dairy and non-dairy cattle are key categories according to level and trend assessment for the base year and 2023. Total methane emissions from enteric fermentation decreased from 111.14 Gg in 1990 to 36.68 Gg in 2023 (-67%) and had decreased by nearly 0,19% compared to the previous year. More information is available in [Table 5.25](#) and on [Figure 5.19](#).

**Table 5.25: Methane emissions from enteric fermentation according to livestock in particular years**

YEAR	DAIRY CATTLE	NON-DAIRY CATTLE	SHEEP	GOAT	HORSES	SWINE
	CH <sub>4</sub> in Gg					
1990	34.548	65.246	7.272	0.052	0.245	3.781
1995	24.779	37.623	5.030	0.125	0.182	3.115
2000	24.676	21.454	3.787	0.257	0.171	2.233
2005	22.166	18.011	3.559	0.198	0.150	1.662
2010	18.060	17.653	4.357	0.176	0.128	1.031
2011	17.714	17.829	4.329	0.170	0.125	0.871
2012	17.466	18.452	4.521	0.174	0.130	0.947
2013	16.788	18.858	4.406	0.177	0.129	0.956
2014	16.254	18.488	4.326	0.176	0.123	0.963
2015	15.979	18.903	4.316	0.182	0.124	0.950
2016	15.610	18.489	4.066	0.182	0.115	0.879
2017	15.363	18.709	3.997	0.185	0.111	0.922
2018	15.041	18.837	3.901	0.185	0.128	0.941
2019	14.958	18.477	3.532	0.178	0.125	0.884
2020	14.343	19.377	3.268	0.053	0.110	0.807
2021	14.481	18.125	3.245	0.052	0.121	0.680
2022	14.277	18.207	3.510	0.055	0.127	0.571
2023	14.148	18.369	3.373	0.054	0.133	0.605


**Figure 5.19:** Trend in methane emissions (Gg) by animals in enteric fermentation in 1990 – 2023



Methane emissions from dairy and non-dairy cattle represent the significant share of emissions in enteric fermentation (38,6% and 50%). More than 9% belongs to sheep methane emissions. These animals are significant in this category and were estimated by tier 2 approach. Other animal categories were determined by tier 1 approach. The share of emissions in animal categories in enteric fermentation is shown on [Figure 5.20](#).

The trend of methane emissions from enteric fermentation correlates with the number of livestock, especially the key categories, where tier 2 approach was implemented. The correlations between important parameters (milk yield, weight gain, percentage digestibility of feed ratio) in the particular years were verified. In 1995, the number of animals increased in the two key categories: sheep (+7.5%) and other cattle (+3.5%). Emissions increased at a comparable rate. A similar situation is visible in 2012, where the number of non-dairy cattle increased by 2% and sheep by 4%. A completely different situation is visible in the partial year 2015, where the increase is mainly due to increases in the number of goats (+3%), and horses (+0.6%). The development of number of livestock had a significant impact on reducing emissions in this sector since 1990.

**Figure 5.20:** The share of aggregated emissions by categories within enteric fermentation in 2023



### 5.7.1. Methodological Issues – Methods

The cooperation with the NPPC-VÚŽV continues. Changes and improvements are entirely in accordance with tier 2 for key categories of animal categories (cattle and sheep). For other non-key categories of animals (goats, horses, and swine), tier 1 was used ([Table 5.33](#)). The overview is provided in [Tables 5.26-5.33](#). Used methodology is based on detailed national data about animals' number (more advanced livestock characteristics and better structured number of livestock). Data on animal numbers were provided by the ŠÚ SR.

The regional input data about feeding, weight, milk production, and wool production were provided by the ŠÚ SR. Other parameters for dairy cattle, non-dairy cattle and sheep categories (significant animal categories in Slovakia) were provided by the NPPC-VÚŽV.

**Cattle** – due to increase of transparency in methodology and used activity data, emissions estimation was completed by the parameters for average animal weight (597.76 kg), share of pregnancy (68.96%) and share of digestibility of feed (72.76%). Typical feeding for cattle is maize and alfalfa silage, cereal, hay and pasture in the Slovak Republic.

Total methane emissions from enteric fermentation of cattle were estimated based on the detailed classification of animals into the following categories: dairy cattle, high producing dairy cows in the 3.A.1.1 sub-category and other non-dairy cattle in the 3.A.1.2 sub-category (suckler cows, calves six months, heifer, pregnancy heifer, breeding bull, oxen, fattening). Slovak country specific approach is based on the particular division of non-dairy cattle. Part of non-dairy cattle is divided into milk type and beef type. The primary differences are in different breeding conditions and feeding doses. The feeding doses of the beef non-dairy cattle is mostly pasture and hay. Cereal and silage are added mainly into the feeding ration in milk type of non-dairy cattle. Different feeding rations are desirable during muscle mass formation (beef non-dairy cattle need to have higher daily muscle mass gain than milk type of non-dairy cattle). Milk type of non-dairy cattle is bred similarly as dairy cows. On the contrary, beef cattle is bred principally as slaughter. The country specific EFs for dairy and non-dairy cattle are estimated as weighted average of regions based AGEI and other parameters specific for each category.

**Table 5.26:** The overview of used country specific parameters for dairy cattle and suckler cows in 2023

PARAMETER*	UNIT	DAIRY COWS	SUCKLER COWS	SOURCES OF PARAMETERS**
Body weight	kg	597.76	594.99	NPPC-VÚŽV
Milk yield	l/day	22.31	4.50	Parameter from the ŠÚ SR
Milk yield	kg/day	23.00	4.64	Calculated parameter
Fat milk	%	3.94	4.00	Parameter from the ŠÚ SR
DE	%	72.76	64.83	Calculated parameter – based on feeding statistics
Ym	%	6.10	0.070	Default value from IPCC 2019 RF
Maintenance NEm	MJ/day	45.50	42.11	Calculated parameter eq. 10.3 (IPCC 2019 RF)
Activity NEa	MJ/day	0.94		Calculated parameter eq. 10.4 (IPCC 2019 RF)
Lactation Ni	MJ/day	70.19	8.31	Calculated parameter eq. 10.8 (IPCC 2019 RF)
Pregnancy NEp	MJ/day	3.14	3.48	Calculated parameter eq. 10.13 (IPCC 2019 RF)
Ratio of net energy REM		0.54	0.51	Calculated parameter eq. 10.14 (IPCC 2019 RF)
Ratio of net energy REG		0.34	0.31	Calculated parameter eq. 10.15 (IPCC 2019 RF)
Gross energy	MJ/head/day	306.80	204.75	Calculated parameter eq. 10.16 (IPCC 2019 RF)
EFs	kg/head/year	123.13	94.00	Calculated parameter eq.10.21 (IPCC 2019 RF)

**Table 5.27:** The overview of used country specific parameters for non-dairy cattle milk type in 2023

PARAMETER*	UNIT	CALVES 6 MONTHS	HEIFER	HEIFER PREGNANT	FATTENING	OXEN	BREEDING BULL
Body weight	kg	115.52	295.74	497.92	349.97	700.00	800.00
Daily gain	kg	0.840	0.67	0.68	0.78	0.63	-
DE	%	81.44	70.22	70.86	72.36	72.04	68.80
Ym	%	3.2	0.06	0.06	0.06	0.06	0.06
Maintenance NE <sub>m</sub>	MJ/day	11.34	22.96	33.93	29.93	50.35	55.66
Activity NE <sub>a</sub>	MJ/day	-	1.56	0.88	-	-	-
Growth NE <sub>g</sub>	MJ/day	12.47	13.29	14.89	10.93	11.53	-
NE <sub>p</sub>	MJ/day	-	-	3.39	-	-	-
REM		0.55	0.53	0.53	0.53	0.53	0.53
REG		0.37	0.33	0.34	0.34	0.34	0.33
Gross energy	MJ/head/day	66.48	122.73	164.07	121.42	177.88	153.92
EFs	kg/head/year	13.74	50.71	67.80	50.17	73.50	63.60

**Table 5.28:** The overview of used country specific parameters for non-dairy cattle beef type in 2023

PARAMETER*	UNIT	CALVES 6 MONTHS	HEIFER	HEIFER PREGNANT	FATTENING	OXEN	BREEDING BULL
Body weight	kg	127.19	374.34	602.55	349.29	700.00	800.00
Daily gain	kg	0.92	0.50	0.50	0.74	0.66	-
DE	%	76.28	65.60	64.49	65.91	65.03	68.80
Ym	%	0.03	0.07	0.07	0.063	0.063	0.06
Maintenance NE <sub>m</sub>	MJ/day	12.19	27.40	39.15	29.89	50.35	55.66
Activity NE <sub>a</sub>	MJ/day	1.39	5.40	7.72	-	-	4.94
Growth NE <sub>g</sub>	MJ/day	13.57	9.31	11.21	10.77	12.24	-
NE <sub>p</sub>	MJ/day	-	-	3.24	-	-	-
REM		0.54	0.52	0.51	0.52	0.51	0.53
REG		0.36	0.31	0.31	0.31	0.31	0.33
Gross energy	MJ/head/day	82.71	142.49	208.61	139.91	211.53	167.58
EFs	kg/head/year	17.97	65.42	95.78	57.81	87.41	69.25

\*weighted average \*\*sources of parameters are the same for dairy and non-dairy cattle

Average weight of cattle was calculated based on breed structure in the Slovak Republic. Breed structure of cattle is divided on the heavy (Slovak spoken, Holsteins, Braunvieh) and light breed (Pinzgauer and others). Average weight of heavy breed is 600 kg and average body weight of light breed is 500 kg. Different annual share of breed in cattle herd caused differences of body weight. Data about breed structure was taken from the PLIS – [Information System about Breeds](#).

Milk production is taken from the Statistical Yearbook. Digestibility of feed (DE) is calculated as a weighted average of calculated values from the feed ration and provided by the NPPC-VÚŽV. The methane conversion factor is in line with the default values provided in the IPCC 2019 RF. Gross energy is the sum of energies calculated by formulas referred to the IPCC 2019 RF with using typical national breed conditions. National emission factors were calculated by this approach for cattle (dairy and non-dairy).

Following formula was used for EFs calculation: 
$$EF = \left[ \frac{GE * \left( \frac{Y_m}{100} \right) * 365}{55.65} \right]$$

Where: **EF** = emission factor in kg CH<sub>4</sub>/head, **GE** = gross energy intake in MJ/head/day, **Y<sub>m</sub>** = methane conversion factor in percent of gross energy in feed converted to methane, **factor 55.65** = the energy content of methane in MJ/kg CH<sub>4</sub>.

**Table 5.29: Activity data, EFs and methane emissions for dairy cattle in particular years**

YEAR	POPULATION	MILK YIELD	AGEI	EFs	CH <sub>4</sub> EMISSIONS
	1 000 heads	kg/day	MJ/head/day	kg/head	Gg
1990	401.123	6.963	199.829	86.128	34.548
1995	262.664	8.829	218.873	94.336	24.779
2000	242.496	12.236	237.352	101.759	24.676
2005	198.580	15.180	260.901	111.624	22.166
2010	159.260	15.619	265.090	113.397	18.060
2011	154.105	16.347	268.789	114.945	17.714
2012	150.272	17.220	276.584	116.230	17.466
2013	144.875	17.342	274.689	115.879	16.788
2014	143.083	17.743	270.931	113.602	16.254
2015	139.229	17.846	271.465	114.770	15.979
2016	132.610	19.411	283.013	117.717	15.610
2017	129.863	19.557	283.271	118.303	15.363
2018	127.871	19.891	283.863	117.626	15.041
2019	125.848	20.217	287.557	118.857	14.958
2020	122.049	20.363	287.557	117.520	14.343
2021	120.068	22.019	300.454	120.611	14.481
2022	116.910	22.861	304.038	122.121	14.277
2023	114.896	22.997	306.800	123.134	14.148

**Table 5.30: Activity data, EFs and methane emissions for non-dairy cattle in particular years**

YEAR	POPULATION	AGEI	EFs	CH <sub>4</sub> EMISSIONS
	1 000 heads	MJ/head/day	kg/head	Gg
1990	1 161.947	135.726	56.153	65.246
1995	666.042	136.305	56.488	37.623
2000	403.652	130.629	53.151	21.454
2005	329.309	133.978	54.693	18.011
2010	307.865	139.992	57.340	17.653
2011	309.253	140.512	57.653	17.829
2012	320.819	142.876	57.516	18.452
2013	322.945	144.375	58.392	18.858
2014	322.460	135.853	57.335	18.488

YEAR	POPULATION	AGEI	EFs	CH <sub>4</sub> EMISSIONS
	1 000 heads	MJ/head/day	kg/head	Gg
2015	318.357	147.630	59.377	18.903
2016	313.502	146.920	58.974	18.489
2017	309.963	147.728	60.357	18.709
2018	310.984	148.156	60.573	18.837
2019	306.405	146.280	60.301	18.477
2020	320.240	146.666	60.507	19.377
2021	314.021	141.875	57.719	18.125
2022	316.265	141.498	57.568	18.207
2023	314.825	142.283	58.347	18.369

**Sheep** – total methane emissions from enteric fermentation of sheep were estimated based on the detailed classification of animals into two categories: milk sheep (ewes, ewe lambs, mated yearlings, rams) and beef sheep (ewes, ewe lambs, mated yearlings, rams). The emission factors are calculated as a weighted average from these four categories based on gross energy intake (milk productivity, wool productivity, average methane conversion rate) and other country specific information. Presented calculation approach and parameters were published.<sup>3</sup>

**Table 5.31: The overview of used country specific parameters for sheep in 2023**

PARAMETER*	UNIT	DAIRY SHEEP				BEEFSHEEP			
		A	B	C	D	E	F	G	H
Body weight	kg	60.00	32.50	55.00	80.00	70.00	47.50	65.00	90.00
Milk yield	l/day	0.475	-	-	-	0.266	-	-	-
Milk yield	kg/day	0.489	-	-	-	0.274	-	-	-
DE of feed	%	60.8	59.0	59.0	58.8	60.82	58.7	58.7	58.6
Y <sub>m</sub>	%	6.50	6.50	6.50	6.50	6.50	6.50	6.50	6.50
Maintenance NE <sub>m</sub>	MJ/day	4.68	3.13	4.38	6.68	5.25	4.16	4.97	7.29
Activity NE <sub>a</sub>	MJ/day	1.01	0.33	0.93	0.77	1.21	0.48	1.14	0.86
Lactation NE <sub>l</sub>	MJ/day	2.17	-	-	-	1.26	-	-	-
Wool production Ne <sub>wool</sub>	MJ/day	0.11	-	0.11	0.11	0.11	-	0.11	0.11
Growth NE <sub>g</sub>	MJ/day	-	1.20	1.79	-	-	1.64	2.09	-
Pregnancy NE <sub>p</sub>	MJ/day	0.44	-	0.36	-	0.49	-	0.47	-
REM		0.50	0.49	0.49	0.49	0.50	0.49	0.49	0.49
REG		0.28	0.27	0.27	0.27	0.28	0.27	0.27	0.27
Gross energy	MJ/head/day	28.31	19.48	31.50	26.56	27.74	26.65	36.99	29.22
EFs	kg/head/year	12.07	8.31	13.43	11.32	11.83	11.36	15.77	12.46

A: Mature ewes, B: Growing lambs, C: Growing lambs pregnant, D: Other mature sheep, E: Mature ewes, F: Growing lambs, G: Growing lambs pregnant, H: Other mature sheep, \*weighted average

Activity data for sheep is available in individual categories (mature ewes, growing lambs and other mature sheep) on regional level provided by the ŠÚ SR for 1990 – 2023. Data were provided including the input parameters (the wool production and the amount of milk for categories ewes). Milk production is taken from the Statistical Yearbook. Digestibility of feed (DE) is calculated as a weighted average of calculated values from the feed ration and provided by the NPPC-VÚŽV. Emission factors for sheep were estimated based on milk production, wool production, and average gross energy intake. These parameters are country specific. Methane emissions from enteric fermentation of mature sheep reflect milk production for the period 1997 – 2023. The extrapolation (linear function) was used for

<sup>3</sup> Differences in amounts of methane emissions from enteric fermentation from Slovak ewe farming between 2015 and 2016

reconstruction of milk production at regional level back to the base year. The net energy required for pregnancy (NEp) was calculated according to the Equation 10.13 p.10.28 of the IPCC 2019 RF. Pregnancy coefficient (Cp) for mature ewes and pregnant growing lambs was taken from Table 10.7 of the IPCC 2019 RF. Values reported in 2022 were 100% in pregnant growing lambs and 95.7% in mature ewes.

**Table 5.32: Activity data, EFs and methane emissions for sheep in particular years**

YEAR	POPULATION	AGEI	EFs	CH <sub>4</sub> EMISSIONS
	1 000 heads	MJ/head/day	kg/head	Gg
1990	600.426	28.40856912	12.111	7.272
1995	427.844	27.80407474	11.756	5.030
2000	347.983	27.75008348	10.881	3.787
2005	320.487	27.66105292	11.104	3.559
2010	394.175	25.92468317	11.052	4.357
2011	393.927	25.77605394	10.989	4.329
2012	409.569	25.89362208	11.039	4.521
2013	399.908	25.84443194	11.018	4.406
2014	391.151	25.94166289	11.060	4.326
2015	381.724	25.94558663	11.306	4.316
2016	368.896	25.85076758	11.021	4.066
2017	365.344	25.66299388	10.941	3,997
2018	351.122	26.05744793	11.109	3.901
2019	320.555	25.84433204	11.018	3.532
2020	294.252	26.05266711	11.107	3.268
2021	290.918	26.16653407	11.155	3.245
2022	301.131	27.34053693	11.656	3.510
2023	288.736		11.683	3.373

Goats, horses, and swine – emission factors for goats, horses and swine in enteric fermentation are default (IPCC 2019 RF) constantly used for whole time series. EF for goats is 5 kg/head/year (low productivity system, EF for horses is 18 kg/head/year and EF for swine is 1.5 kg/head/year (High productivity system) ([Table 5.33](#)). According to our long term improvements plans, tier 2 approach in the swine category will be developed in the future submissions. Implementation of tier 2 approach for swine is not processed due to lack of reliable data on methane conversion factor (YM). Other categories are insignificant sources of emissions.

**Table 5.33: Activity data, EFs and methane emissions for other animals in particular years**

YEAR	GOATS			HORSES			SWINE		
	HEADS	EFs	CH <sub>4</sub>	HEADS	EFs	CH <sub>4</sub>	HEADS	EFs	CH <sub>4</sub>
	1 000	kg/head	Gg	1 000	kg/head	Gg	1 000	kg/head	Gg
1990	10.322	5.000	0.052	13.595	18.000	0.245	2 520.524	1.500	3.781
1995	25.046	5.000	0.125	10.109	18.000	0.182	2 076.439	1.500	3.115
2000	51.419	5.000	0.257	9.516	18.000	0.171	1 488.441	1.500	2.233
2005	39.566	5.000	0.198	8.328	18.000	0.150	1 108.265	1.500	1.662
2010	35.292	5.000	0.176	7.111	18.000	0.128	687.260	1.500	1.031
2011	34.053	5.000	0.170	6.937	18.000	0.125	580.393	1.500	0.871
2012	34.823	5.000	0.174	7.249	18.000	0.130	631.464	1.500	0.947
2013	35.457	5.000	0.177	7.161	18.000	0.129	637.167	1.500	0.956
2014	35.178	5.000	0.176	6.828	18.000	0.123	641.827	1.500	0.963
2015	36.324	5.000	0.182	6.866	18.000	0.124	633.116	1.500	0.950
2016	36.355	5.000	0.182	6.407	18.000	0.115	585.843	1.500	0.879
2017	37.067	5.000	0.185	6.145	18.000	0.111	614.384	1.500	0.922

YEAR	GOATS			HORSES			SWINE		
	HEADS	EFs	CH <sub>4</sub>	HEADS	EFs	CH <sub>4</sub>	HEADS	EFs	CH <sub>4</sub>
	<i>1 000</i>	<i>kg/head</i>	<i>Gg</i>	<i>1 000</i>	<i>kg/head</i>	<i>Gg</i>	<i>1 000</i>	<i>kg/head</i>	<i>Gg</i>
2018	36.907	5.000	0.185	7.102	18.000	0.128	627.022	1.500	0.941
2019	35.594	5.000	0.178	6.960	18.000	0.125	589.228	1.500	0.884
2020	10.589	5.000	0.053	6.099	18.000	0.110	538.310	1.500	0.807
2021	10.434	5.000	0.052	6.738	18.000	0.121	453.076	1.500	0.680
2022	11.008	5.000	0.055	7.044	18.000	0.127	380.895	1.500	0.571
2023	10.719	5.000	0.054	7.367	18.000	0.133	403,037	1.500	0.605

### 5.7.2. Activity Data

Primary data sources used for the emissions evaluations were published in the Census of Sowing Areas of Field Crops in the Slovak Republic, the Annual Census of Domestic Livestock in the Slovak Republic, the Statistical Yearbooks 1990 – 2022 and the research results from projects and studies provided by several organizations inside the NPPC-VÚŽV.

Activity data for dairy, non-dairy cattle, sheep and swine are based on bottom-up statistical information at the regional level. The used input parameters were calculated as weighted averages. The ŠÚ SR provides annual livestock numbers at a detailed regional level in Livestock Census annually on 31<sup>st</sup> December.

Due to a different regionalisation of Slovakia in years 1990 – 1996 (only three regions: Západoslovenský, Stredoslovenský, and Východoslovenský), it was not possible to use time series immediately. The reallocation of older data into new regions (8 districts after 1997) was necessary. Reallocation was based on the following assumptions:

Západoslovenský region (1990 – 1996) is equal to Bratislavský, Nitriansky, Trnavský, Trenčiansky regions (1997 – present);

- Stredoslovenský region (1990 – 1996) is similar to Banskobystrický and Žilinský regions (1997 – present);
- Východoslovenský region (1990 – 1996) is similar to Prešovský and Košický regions (1997 – present).

A reallocation was prepared by using the linear extrapolation tools to reach statistical totals as reported by the ŠÚ SR and time series was extrapolated back to the base year. The ŠÚ SR and the SHMÚ use a standard statistical approach for data extrapolations. Good statistical practice is described in the [EUROSTAT Guidance](#). After 2017 submission, extrapolated number of swine was reported. The SHMÚ filled the data gap by using a standard statistical approach for extrapolation (linear extrapolation in spreadsheets). In 2017 submission, the ŠÚ SR provided complete time-series of official data, which is consistent with the EUROSTAT and the FAOSTAT ([Chapter 5.3.1](#)). In addition, time series 1997 – 2020 of the milk production, wool production and daily gain for cattle and sheep at regional level was provided by the ŠÚ SR in 2016. Activity data used for methane emissions estimation is summarized in [Table 5.34](#). Detailed statistical information is available at the regional level and emissions are estimated by bottom-up method (tier 2). The NPPC-VÚŽV implemented the results of a questionnaire farm survey where a better classification and disaggregation of cattle categories were used. Based on survey data, cattle were divided into dairy and non-dairy. Dairy cattle are estimated separately from non-dairy cattle. Dairy cattle are defined as cows that produce milk only for human consumption (highly productive cows). Suckler cows are defined as cows that are farmed for nutrition of calves (low productive cows). Suckler cows are included in non-dairy cattle category. In addition, non-dairy cattle includes breeding bull, oxen, calves, heifer pregnant, un-pregnant heifers and fattening bulls. This categorization is consistent in whole time series. The number of livestock decreased compared to the previous year in all species. The



highest declines were recorded in the swine category (-84%) compared to 1990. The main reason for this decrease is the data gap on self-sufficiency - small household's farmers and morbidity of animals. The same reason was the cause of the decline of poultry (-41%) and horses (-46%).

Between 2005 and 2023, the production of most agricultural crops showed an increasing trend. The production of potatoes increased by +59%, soya +49%, oil plants +47%, wheat +42%, meadows +35%, rye +19%, clover +16% and leguminous plant +4% during the given period. The decrease was recorded for tobacco by -100% and for beans by -100%.

Since 2005, livestock numbers have decreased for all farmed species. Between 2005 and 2023, the number of pigs decreased by -42%, dairy cattle by -64%, poultry by -31% and sheep by -10%.

**Table 5.34: Animal population (heads) according to categories at regional level for the year 2023**

REGION		A	B	C	D	E	F	G	H
DAIRY CATTLE		4 790	19 065	13 729	17 993	20 163	14 419	17 300	7 437
NON-DAIRY CATTLE	Suckling cows	1 658	2 024	4 558	1 807	9 466	18 602	23 351	11 980
	Calves in 6 month (milk sort)	1 787	9 491	5 526	8 489	6 634	4 984	5 640	2 406
	Heifer (milk sort)	971	5 077	4 502	5 875	8 575	5 091	6 281	2 765
	Heifer (pregnant) (milk sort)	1 434	4 906	3 508	6 743	4 804	3 037	3 742	1 427
	Fattening (milk sort)	481	9 981	4 164	7 243	3 815	3 829	3 886	1 864
	Oxen (milk sort)	4	4	7	9	213	42	35	8
	Breeding bull (milk sort)	21	134	68	109	291	387	455	316
	Calves in 6 month (beef sort)	619	1 008	1 835	852	3 114	6 431	7 612	3 877
	Heifer (beef sort)	336	539	1 494	590	4 026	6 569	8 478	4 454
	Heifer (pregnant) (beef sort)	496	521	1 165	677	2 256	3 919	5 052	2 299
	Fattening (beef sort)	167	1 060	1 383	727	1 791	4 940	5 246	3 003
	Oxen (beef sort)	1	0	2	1	100	55	48	13
	Breeding bull (beef sort)	41	267	135	218	583	775	911	632
SHEEP	Mature ewes	1 923	1 437	20 039	5 038	49 401	59 887	38 614	15 622
	Growing lambs	664	776	6 867	1 754	16 443	17 717	11 137	4 180
	Growing lambs (pregnant)	411	96	3 962	1 021	8 307	8 404	5 871	2 144
	Other mature sheep	58	38	601	152	1 443	1 743	1 112	444
SWINE	Breeding swine	5 559	17 822	4 976	8 118	132	6 498	248	695
	Fattening swine	13 323	163 666	29 884	105 216	1 644	26 519	5 155	13 582
HORSES	Horses (0-1year)	10	24	89	47	32	19	37	69
	Horses (1-3year)	33	74	206	267	82	87	106	172
	Stallions	50	61	107	61	81	86	110	32
	Mares	323	210	470	457	440	583	425	420
	Castrated stallions	297	100	266	244	347	369	225	226

REGION		A	B	C	D	E	F	G	H
GOATS	Mature goats	245	250	739	360	1 721	1 623	1 144	898
	Growing goats (pregnant)	98	9	41	67	598	46	169	158
	Other mature goats	120	96	328	108	245	469	533	654
POULTRY	Laying hens and roosters	547 615	113 768	228 502	905 606	406 618	440 161	14 808	207 962
	Breeding broilers	188 490	77 275	89 879	320 840	77 412	133 222	326	324 263
	Fattening broilers	25 856	1 002 652	1 162 082	1 183 866	300 933	1 304 888	89 499	343 605
	Turkeys	6	5 987	207	109 129	31 465	19	0	4
	Ducks	5	31 442	9	262	35	83	1	27
	Geese	0	56	5	387	2	20	4	93

REGIONS: A: Bratislava, B: Trnava, C: Trenčín, D: Nitra, E: Žilina, F: Banská Bystrica, G: Prešov, H: Košice

## 5.8. Manure Management (CRF 3.B.1) – CH<sub>4</sub> Emissions

*Emitted gas: CH<sub>4</sub>*

*Methods: Tier 1 and Tier 2*

*Emission factors: CS*

*Key sources: Yes*


*Particular significant subcategories: cattle and swine*

Methane can also be emitted in anaerobic conditions due to the decomposition of manure. These conditions can be found in large-scale farms (farms for cattle, fattening pigs and poultry). Methane emissions from manure management are the emissions depending on animal husbandry and the number of animals. Methane from manure management can be better mitigated (proper storage, digesters use) compared to methane originated from enteric fermentation. Mitigation measures possible in enteric fermentation have several limitations. Therefore it can be predicted, that manure management will emit less methane emissions in the future than enteric fermentation.

Methane emissions in manure management decreased from 25.58 Gg in 1990 to 3.62 Gg in 2023 due to decrease in livestock number of all categories except horses. The extreme reduction of animals was recorded in swine and cattle due to economic reasons. This situation consequently influenced methane emissions from the manure management. Emissions decreased by 86% compared to the base year. However, swine is a key category by trend assessment, tier 2 category was used for this category. Methane emissions in manure management increased in comparison with the previous year by 0,22%, caused by small increased number of swine, poultry and horses. [Figure 5.21](#) and [Table 5.35](#) summarize the overall situation. Methane emissions produced in manure management for cattle (dairy and non-dairy), swine and sheep were estimated using tier 2 and country specific emissions factors and parameters.

This estimation was provided in line with the emissions estimation in enteric fermentation based on regional data. In the previous years, the Slovak Republic was constantly developing a new approach of methane emissions estimation from swine. The NPPC-VÚŽV prepared the new country specific parameters, which were used in implementation of tier 2 approach. Swine are divided into two separate categories – market swine (fattening pigs) and breeding swine (sows, piglet's hogs for breeding purpose).

**Figure 5.21: Trend in CH<sub>4</sub> emissions (Gg) by categories within manure management in 1990 – 2023**




**Table 5.35: CH<sub>4</sub> emissions from manure management according to the animals in particular years**

YEAR	DAIRY CATTLE	NON-DAIRY CATTLE	SWINE	SHEEP	HORSES	POULTRY	GOATS
	CH <sub>4</sub> in Gg						
1990	3.957	3.000	17.780	0.212	0.040	0.584	0.002
1995	2.757	1.650	9.791	0.146	0.029	0.494	0.005
2000	2.542	0.960	6.865	0.111	0.028	0.462	0.010
2005	2.176	0.763	5.108	0.104	0.023	0.470	0.008
2010	1.919	0.690	3.321	0.127	0.019	0.452	0.007
2011	1.891	0.690	2.722	0.126	0.019	0.410	0.007
2012	1.867	0.731	2.914	0.132	0.020	0.425	0.007
2013	1.832	0.760	2.550	0.128	0.020	0.393	0.007
2014	1.594	0.683	2.882	0.126	0.019	0.428	0.007
2015	1.437	0.707	2.771	0.123	0.019	0.445	0.007
2016	1.398	0.692	2.386	0.119	0.018	0.429	0.007
2017	1.421	0.710	2.562	0.116	0.017	0.455	0.007
2018	1.416	0.720	2.750	0.113	0.020	0.479	0.007
2019	1.356	0.690	2.492	0.103	0.020	0.442	0.007
2020	1.324	0.731	1.922	0.095	0.017	0.332	0.002
2021	1.273	0.684	1.472	0.094	0.019	0.324	0.002
2022	1.248	0.686	1.247	0.102	0.020	0.302	0.002
2023	1.238	0.692	1.264	0.098	0.021	0.301	0.002

Figure 5.22 shows the share of individual categories on the production of manure methane emissions. Significant share is represented by cattle (53,3%). The important animal category is also swine 35%.

Methane emissions are calculated by the same IPCC methodology as used enteric fermentation. Emissions estimation in 3.A and 3.B are estimated with using the common parameters. Anyway, the key category of manure management methane emissions is 3.B.1.3 Swine category with high impact on emission trend in 3.B.1 category. In partial years number of swine decreased by -7% (1992), about 12% in 1998, and increased 6% in 2023. In 1992 and 1998, the number of all animal species decreased, except goats. Goats and horses categories have not significant effects on methane level in manure management. The number declined due to the economic situation at that time.

**Figure 5.22:** The share of methane emissions by animals within manure management in 2023



### 5.8.1. Methodological Issues – Methods

Cattle, sheep, swine, poultry - tier 2 approach based on national data was applied for methane emissions estimation in manure management for cattle, sheep and swine categories. Country specific parameters were introduced into estimation. The national approach is based on the number of animals divided by subcategories per region, the calculation of volatile solid excretion (VS), which is calculated from the gross energy intake, digestibility of the feed, ash urinary energy and methane conversion factor (MCF), expressed as inputs to the equation for the estimation of national EFs (Tables 5.36 - 5.42).

$$EF = (VS * 365) * \left[ B_o * \frac{0.67 \text{ kg}}{\text{m}^3} * \sum \frac{\text{MCF}}{100} * \text{MS} \right]$$

Where: **VS** = daily volatile solid excreted for livestock category, kg DM animal/day, **365** = annual VS production in days/year, **B<sub>o</sub>** = maximum methane producing capacity for manure by livestock category in m<sup>3</sup> CH<sub>4</sub>/kg of VS excreted, **0.67** = conversion factor of m<sup>3</sup> CH<sub>4</sub> to kilogram CH<sub>4</sub>, **MCF** = methane conversion factors for each manure management system S by climate region (%), **MS** = fraction of livestock category manure handled using manure management system S in climate region (cool).

The VS calculation is consistent with the equation 10.23, p 10.64 (IPCC 2019 RF).

Emission factors for cattle, swine and sheep are calculated as weighted average (region and animals). Values of maximum methane production capacity and emission factors for dairy cattle are shown in Table 5.43 for non-dairy cattle in Tables 5.44 and 5.45. Data for sheep is in Tables 5.46 and 5.57.

**Table 5.36:** Overview of country specific parameters used for cattle and sheep in 1990

PARAMETERS	UNIT	DAIRY CATTLE	NON-DAIRY CATTLE	MATURE EWES	GROWING LAMBS	OTHER MATURE SHEEP
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.24	0.18	0.19	0.19	0.19
Typical animal mass average	kg	589.41	330.08	64.50	53.85	84.61
Ash content	%	8	8	8	8	8
VS daily excretion	kg dm/head/day	3.59	2.51	0.57	0.72	0.62
Liquid system		26	26	NO	NO	NO

PARAMETERS	UNIT	DAIRY CATTLE	NON-DAIRY CATTLE	MATURE EWES	GROWING LAMBS	OTHER MATURE SHEEP
Solid storage and dry lot		2	2	2	2	2
PRP		0.47	0.47	0.47	0.47	NO
Digesters*		NO	NO	NO	NO	NO

**Table 5.37:** Overview of country specific parameters used for cattle and sheep in 2023

PARAMETERS	UNIT	DAIRY CATTLE	NON-DAIRY CATTLE	MATURE EWES	GROWING LAMBS	OTHER MATURE SHEEP
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.24	0.18	0.19	0.19	0.19
Typical animal mass average	kg	597.76	386.28	63.65	44.37	83.65
Ash content	%	8	8	8	8	8
VS daily excretion**	kg dm/head/day	4.77	2.52	0.60	0.58	0.62
Liquid system		26	26	NO	NO	NO
Solid storage and dry lot		2	2	2	2	2
PRP		0.47	0.47	0.47	0.47	0.47
Digesters		NO	NO	NO	NO	NO

**Table 5.38:** Overview of country specific parameters used for breeding swine in 1990

PARAMETERS	UNIT	A	B	C	D	E
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.45	0.45	0.45	0.45	0.45
Typical animal mass average	kg	200	85	140	145	35.5
Ash content	%	10	10	10	10	10
VS daily excretion**	kg dm/head/day	0.62	0.46	0.38	0.41	0.22
Liquid system		16	16	16	16	16
Solid storage and dry lot		2	2	2	2	2

A: Sows, B: Gilts non-pregnant, C: Gilts pregnant, D: Hogs; E: Piglets 21-50 kg

**Table 5.39:** Overview of country specific parameters used for breeding swine in 2023

PARAMETERS	UNIT	A	B	C	D	E
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.45	0.45	0.45	0.45	0.45
Typical animal mass average	kg	200	85	140	145	35.5
Ash content	%	10	10	10	10	10
VS daily excretion**	kg dm/head/day	0.64	0.50	0.41	0.41	0.17
Liquid system		16	16	16	16	16
Solid storage and dry lot		2	2	2	2	2

A: Sows, B: Gilts non-pregnant, C: Gilts pregnant, D: Hogs; E: Piglets 21-50 kg

**Table 5.40:** Overview of country specific parameters used for market swine in 1990

PARAMETERS	UNIT	A	B	C	D	E
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.45	0.45	0.45	0.45	0.45
Typical animal mass average	kg	10.60	35.50	65	95	110
Ash content	%	10	10	10	10	10
VS daily excretion**	kg dm/head/day	0.21	0.38	0.56	0.71	0.79
Liquid system		16	16	16	16	16
Solid storage and dry lot		2	2	2	2	2
Deep bedding		26	26	26	26	26

A: Fattening pigs up to 20 kg; B: Fattening pigs 21-50 kg; C: Fattening pigs 50-80 kg; D: Fattening pigs 80-110 kg; E: Fattening pigs over 110 kg

**Table 5.41: Overview of country specific parameters used for market swine in 2023**

PARAMETERS	UNIT	A	B	C	D	E
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.45	0.45	0.45	0.45	0.45
Typical animal mass average	kg	10.60	35.50	65	95	110
Ash content	%	10	10	10	10	10
VS daily excretion**	kg dm/head/day	0.10	0.18	0.26	0.33	0.36
Liquid system		16	16	16	16	16
Solid storage and dry lot		2	2	2	2	2
Deep bedding		26	26	26	26	26

A: Fattening pigs up to 20 kg; B: Fattening pigs 21-50 kg; C: Fattening pigs 50-80 kg; D: Fattening pigs 80-110 kg; E: Fattening pigs over 110 kg; \*B<sub>o</sub> for Western Europe was chosen; \*\*VS daily excretion were taken from table 10A-4 in the IPCC 2019 RF

**Table 5.42: Overview of country specific parameters used for poultry in 2023**

PARAMETERS	UNIT	A	B	C	D	E	F
B <sub>o</sub> *	m <sup>3</sup> /kg VS	0.39	0.36	0.36	0.36	0.36	0.36
Typical animal mass average	kg	2	1.02	1.02	3.28	1.53	2.05
Ash content	%	0.058	0.027	0.028	0.032	0.025	0.027
VS daily excretion**	kg dm/head/day	0.031	0.019	0.018	0.046	0.043	0.037
Poultry manure without bedding		1.5	-	-	-	-	-
Poultry manure with bedding		1.5	1.5	1.5	1.5	1.5	1.5
Pasture		-	-	-	-	0.47	0.47

A: Laying hens and cocks, B: Fattening broilers, C: Breeding broilers, D: Turkeys, E: Geese, F: Ducks

**Swine** – Due to the lack of specific methodology for GE calculation in the IPCC 2019 RF in swine category, the country specific methodology was implemented in 2020 submission. The VS calculation is consistent with the equation 10.23, p 10.64 (IPCC 2019 RF).

Methodological approach introduces more accurate country specific data such as gross energy intake (GE in MJ/head/day), digestibility of feed (DE in %) and new ash content. Digestibility of feed (DE in %) provided by the NPPC-VÚŽV, Department of Animal Feed, is calculated as a weighted average of calculated values from the feed ration. Digestibility was estimated based on each supplemented feeding ration. Metabolizable energy (ME) was taken from publication *Sommer and Petrikovič – Nutrition for Pigs*<sup>4</sup>. Ash content for pigs was taken from publication the *Strauch, Baader, Tietjen – Waste from agricultural production*<sup>5</sup>. Gross energy intake was calculated according to publication *Sommer and Petrikovič – Nutrition for Pigs*. The calculated values are in MJ per day.

ME was estimated by “Factorial method.” This method is based on estimated demand of metabolizable energy for the physiological functions such as maintenance, the growth of muscles, growth, and function of internal bodies, lactation and pregnancy. The sum of energies forms the total energy need for the farm animals. Incorporation of proteins (PR, kg/day) and fats (LR, kg/day) in the body is based on energy

<sup>4</sup> Petrikovič, P., Heger, J., Sommer A., 2005, Nutrition for Pigs, The Research Institute of animal production, ISBN 80-88872-45-6 in Slovak

<sup>5</sup> Strauch, D., Baader, W., Tietjen, C., 1995 Waste from agricultural production, Ulmer Eugen Verlag, ISBN-978-3800143283 in German

estimate. These values are default and are special for each pig subcategory for each day from birth up to 300 days of animal based on the equations below (derived from the Gompertz function):  
 $PR = B * P * \ln\left(\frac{P_{MAT}}{p}\right)$ ;  $LR = B * L * \ln\left(\frac{L_{MAT}}{p}\right)$

Where: **B** = growth parameter, **P** and **L** = protein content, fat in the body in kg/day, **P<sub>MAT</sub>**, **L<sub>MAT</sub>** = values of protein content and fat in adult animal body 's, **PR** = storing proteins in the body (kg/day), **LR** = storing fat in the body (kg/day).

Incorporation of proteins and fat can be characterized as potential growth abilities of pigs' genotype, assumed that the growth parameter (B) is the same value in all genotype.

$$ME_m = 1.02 * H^{0.6}$$

$$ME_p = PR * 37$$

$$ME_L = LR * 47.7$$

Where: **H** = body weight in kg, **PR** = storing proteins in the body (kg/day), **LR** =storing fat in the body (kg/day), **37** = energy storage costs for storing of proteins 37 MJ/kg, **47.7** = energy storage costs for storing of fat 47.7 MJ/kg.

Total demand of metabolized energy is the sum of energy for maintenance ( $ME_m$ ), energy for protein storage ( $ME_p$ ), energy for fat storage ( $ME_L$ ) (Noblet at al.):  $ME = ME_m + ME_p + ME_L$

Where: **ME<sub>m</sub>** = energy for maintenance in MJ/head/day, **ME<sub>p</sub>** = energy for protein storage in MJ/head/kg, **ME<sub>L</sub>** = energy for fat storage in MJ/head/kg, **ME** = metabolizable energy in MJ/head/kg.

ME is the difference between the digestible energy (DE) and the loss of energy in the form of urine and methane gas released by rumen and hind–gut microbes. ME is approximately 96% of DE in pigs, which means that approximately 4% of DE is lost as urine dung and energy. The 4% loss of DE approximates the energy losses, mainly via methane, urinary compounds and heat production by microorganisms in the rumen.

Percentage methane losses from non–ruminants are relatively low, and differences between DE and ME are therefore much smaller:  $DE = \frac{ME}{0.96}$

Where: **ME** = metabolizable energy in MJ/head/kg, **DE** = digestible energy, **0.96** = lost as faeces

Gross energy intake was calculated from digestibility energy and feed. Nutrition data were derived based on estimated daily feed intake:  $GE = \frac{DE}{\%DE}$

Where: **GE** = gross energy intake in MJ/kg/head, **DE** = metabolizable energy in MJ/head/kg, **%DE** = digestibility of feed in %.

**Table 5.43: The overview of used VS and EFs for dairy cattle in 2023**

PARAMETERS	UNIT	Regions							
		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
VS excretion per day on a dry organic matter base	kg VS/day	5.01	5.04	4.95	4.67	4.70	4.55	4.66	4.66
EFs	kg/head	5.86	11.22	10.55	16.24	7.96	13.70	8.71	6.77

**Table 5.44: The overview of used emission factors (kg/head) for non-dairy cattle in 2023**

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
MILK TYPE	Calves in 6. month	0.58	0.63	0.62	0.64	0.72	0.72	0.73	0.54
	Heifers	1.69	1.68	1.87	1.62	1.96	1.70	1.85	1.73

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
	Heifers (pregnant)	2.26	2.35	2.18	2.50	2.45	2.35	2.14	2.25
	Fattening	3.42	4.11	3.99	3.55	3.47	3.94	3.81	3.82
	Oxen	2.35	2.73	2.73	2.45	2.40	2.76	2.68	2.67
	Breeding bull	2.38	2.38	2.38	2.38	2.38	2.38	2.38	2.38
BEEF TYPE	Suckler cows	2.15	2.08	2.02	1.95	2.09	2.09	2.03	1.97
	Calves in 6. month	0.33	0.32	0.27	0.26	0.30	0.30	0.28	0.24
	Heifer	1.45	1.43	1.37	1.28	1.43	1.42	1.39	1.34
	Heifer (pregnant)	2.11	2.07	2.07	1.92	2.15	2.11	2.14	2.04
	Fattening	7.14	8.74	8.18	7.63	8.04	7.89	8.19	7.35
	Oxen	3.10	3.27	3.26	3.06	3.28	3.18	3.34	3.02
	Breeding bull	2.10	2.10	2.10	2.10	2.10	2.10	2.10	2.10

**Tables 5.45:** The overview of used VSs (kg VS/day) for non-dairy cattle in 2023

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
MILK TYPE	Calves in 6. month	0.65	0.72	0.70	0.72	0.82	0.82	0.83	0.62
	Heifers	1.96	1.94	2.16	1.87	2.27	1.96	2.14	2.00
	Heifers (pregnant)	2.61	2.72	2.53	2.89	2.84	2.72	2.47	2.60
	Fattening	1.77	2.12	2.06	1.83	1.79	2.04	1.97	1.97
	Oxen	2.67	3.10	3.10	2.78	2.73	3.14	3.04	3.04
	Breeding bull	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70
BEEF TYPE	Suckler cows	4.20	4.06	3.94	3.81	4.09	4.09	3.97	3.85
	Calves in 6. month	1.29	1.29	1.07	1.13	1.20	1.19	1.15	0.98
	Heifer	2.83	2.79	2.69	2.51	2.80	2.78	2.73	2.61
	Heifer (pregnant)	4.13	4.04	4.05	3.76	4.20	4.12	4.18	3.99
	Fattening	2.39	2.92	2.73	2.55	2.69	2.63	2.73	2.46
	Oxen	3.52	3.72	3.70	3.48	3.72	3.62	3.79	3.43
	Breeding bull	2.94	2.94	2.94	2.94	2.94	2.94	2.94	2.94

**Tables 5.46:** The overview of used emission factors (kg/head) for sheep in 2023

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
DAIRY SHEEP	Mature ewes	0.33	0.34	0.36	0.34	0.35	0.36	0.33	0.32
	Growing lambs	0.00	0.26	0.24	0.24	0.26	0.27	0.23	0.23
	Growing lambs (pregnant)	0.04	0.42	0.39	0.40	0.42	0.45	0.37	0.37
	Other mature sheep	0.50	0.47	0.46	0.55	0.50	0.51	0.46	0.45
BEEF SHEEP	Mature ewes	0.35	0.33	0.30	0.31	0.33	0.35	0.29	0.29
	Growing lambs	0.35	0.33	0.30	0.31	0.33	0.35	0.29	0.29
	Growing lambs (pregnant)	0.49	0.45	0.42	0.43	0.46	0.48	0.40	0.40
	Other mature sheep	0.55	0.51	0.50	0.60	0.55	0.56	0.50	0.49

Due to better disaggregation of sheep based on national data into following subcategories: other mature sheep (VS=0.62 kg dm/head/year), growing lambs (VS=0.58 kg dm/head/year) and mature ewes (VS=0,60 kg dm/head/year), VS can be calculated separately. Values of maximum methane production capacity according to the sheep subcategories are 0.19 m<sup>3</sup>/kg VS. MCF for manure management



systems in cool climate condition (Table 10.21 of the IPCC 2019 RF) was used. Allocation of animals into AWMS is described in [Chapter 5.9.1](#).

**Tables 5.47:** The overview of used VSs (kg VS/day) for sheep in 2023

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
DAIRY SHEEP	Mature ewes	0.59	0.60	0.64	0.60	0.62	0.63	0.58	0.55
	Growing lambs	0.00	0.45	0.41	0.42	0.45	0.48	0.39	0.40
	Growing lambs (pregnant)	0.08	0.74	0.68	0.69	0.74	0.79	0.65	0.65
	Other mature sheep	0.62	0.58	0.57	0.68	0.61	0.63	0.57	0.55
BEEF SHEEP	Mature ewes	0.60	0.59	0.60	0.61	0.60	0.60	0.60	0.59
	Growing lambs	0.65	0.61	0.56	0.57	0.61	0.65	0.53	0.53
	Growing lambs (pregnant)	0.91	0.84	0.77	0.79	0.85	0.90	0.74	0.74
	Other mature sheep	0.67	0.63	0.62	0.74	0.67	0.69	0.62	0.60

Values of maximum methane production capacity and emission factors for swine are provided in [Tables 5.55 and 5.56](#).

**Tables 5.48:** The overview of used emissions factors for swine in 2023

REGION	Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
<b>Sows</b>	8.47	8.86	8.47	8.06	6.81	8.43	6.94	7.51
<b>Gilts non-pregnant</b>	6.69	6.69	6.69	6.69	6.69	6.69	6.69	6.69
<b>Gilts pregnant</b>	5.39	5.39	5.39	5.39	5.39	5.39	5.39	5.39
<b>Hogs</b>	5.39	5.39	5.39	5.39	5.39	5.39	5.39	5.39
<b>Piglets 21-50kg</b>	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28
<b>Fattening to 20 kg</b>	1.43	1.43	1.43	1.43	1.43	1.43	1.43	1.43
<b>Fattening to 21-50 kg</b>	2.59	2.59	2.59	2.59	2.59	2.59	2.59	2.59
<b>Fattening to 50-80 kg</b>	3.79	3.79	3.79	3.79	3.79	3.79	3.79	3.79
<b>Fattening to 80-100 kg</b>	4.76	4.76	4.76	4.76	4.76	4.76	4.76	4.76
<b>Fattening from 110 kg</b>	5.31	5.31	5.31	5.31	5.31	5.31	5.31	5.31

**Tables 5.49:** The overview of used VSs (kg VS/day) for swine in 2023

REGION	Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
Sows	0.64	0.67	0.64	0.61	0.51	0.63	0.52	0.56
Gilts non-pregnant	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50

REGION	Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
Gilts pregnant	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41
Hogs	0.41	0.41	0.41	0.41	0.41	0.41	0.41	0.41
Piglets 21-50kg	0.17	0.17	0.17	0.17	0.17	0.17	0.17	0.17
Fattening to 20 kg	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Fattening to 21-50 kg	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
Fattening to 50-80 kg	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26
Fattening to 80-100 kg	0.33	0.33	0.33	0.33	0.33	0.33	0.33	0.33
Fattening form 110 kg	0.36	0.36	0.36	0.36	0.36	0.36	0.36	0.36

**Tables 5.50:** The overview of used VSs (kg VS/day) for poultry in 2023

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
POULTRY	Laying hens	0.031	0.031	0.031	0.031	0.031	0.031	0.031	0.031
	Fattening broilers	0.019	0.019	0.019	0.019	0.019	0.019	0.019	0.019
	Breeding broilers	0.018	0.018	0.018	0.018	0.018	0.018	0.018	0.018
	Turkeys	0.046	0.046	0.046	0.046	0.046	0.046	0.046	0.046
	Geese	0.043	0.043	0.043	0.043	0.043	0.043	0.043	0.043
	Ducks	0.037	0.037	0.037	0.037	0.037	0.037	0.037	0.037

**Tables 5.51:** The overview of used emission factors (kg/head) for poultry in 2023

REGION		Bratislava	Trnava	Trenčín	Nitra	Žilina	Banská Bystrica	Prešov	Košice
POULTRY	Laying hens	0.044	0.044	0.044	0.044	0.044	0.044	0.044	0.044
	Fattening broilers	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
	Breeding broilers	0.024	0.024	0.024	0.024	0.024	0.024	0.024	0.024
	Turkeys	0.061	0.061	0.061	0.061	0.061	0.061	0.061	0.061
	Geese	0.037	0.037	0.037	0.037	0.037	0.037	0.037	0.037
	Ducks	0.032	0.032	0.032	0.032	0.032	0.032	0.032	0.032

**Other animals** – methodology used for the methane emissions estimation in manure management is based on tier 1 using the default EFs according to the IPCC 2019 RF. Emissions factors are summarized in **Table 5.52**.

**Table 5.52:** Emission factors used for the estimation of CH<sub>4</sub> emissions from manure management

MANURE MANAGEMENT SYSTEMS	Implied EFs in kg CH <sub>4</sub> /year/head
Goats	0.17
Horses	2.90

## 5.8.2. Activity Data

The number of animals is consistent with the number of animals described in the **Chapter 5.3.1 (Table 5.7)**.

# 5.9. Manure Management (CRF 3.B.2) – N<sub>2</sub>O Emissions

EMITTED GAS: N<sub>2</sub>O

METHODS: TIER 1 and TIER 2

EMISSION FACTORS: CS

KEY SOURCES: YES


PARTICULARLY SIGNIFICANT SUBCATEGORIES: CATTLE AND SWINE

Manure nitrogen (N) from cattle production facilities can lead to negative environmental effects, such as contribution to greenhouse gas emissions, leaching and runoff to aqueous ecosystems leading to eutrophication, and acid rain. To mitigate these effects and to improve the efficiency of N use, accurate prediction of N excretion and secretions is required.

Domestic livestock produces different kinds of nitrogen inputs (liquid, solid and deep bedding, litter) into the ecosystem, also the structure of domestic livestock is important (the ratio of different categories of domestic livestock) from direct emissions as well as the emissions from the AWMS. Except for it, the production of nitrogen per head per year also plays a specific role.

Solid and liquid systems are the most common types of excreta storage in manure management (especially for cattle and swine) in the Slovak Republic. The pasture range in some periods of the year (200 days per year on average) is a specific management system for sheep, horses, and goats (partly for non-dairy cattle). The input of nitrogen oxide from manure management was 0.38 Gg of N<sub>2</sub>O in 2023 and the total decrease was 71% compared to the base year (*Figure 5.23* and *Table 5.53*). *Figure 5.24* shows the share of individual categories on the production of nitrogen from manure. A dominant share represents dairy cattle (45%), non-dairy cattle (37%) and swine (9%).


**Figure 5.23:** Trend in N<sub>2</sub>O emissions (Gg) by categories within manure management in 1990 – 2023



**Table 5.53:** *N<sub>2</sub>O emissions (Gg) in manure management according to the animals in particular years*

YEAR	DAIRY CATTLE	NON-DAIRY CATTLE	SHEEP	GOATS	HORSES	SWINE	POULTRY
1990	0.388	0.600	0.037	0.001	0.008	0.248	0.022
1995	0.277	0.334	0.027	0.002	0.006	0.186	0.018
2000	0.292	0.225	0.020	0.003	0.005	0.134	0.017
2005	0.267	0.184	0.020	0.003	0.004	0.096	0.018
2010	0.213	0.157	0.024	0.002	0.004	0.062	0.017
2011	0.211	0.158	0.024	0.002	0.004	0.051	0.015
2012	0.211	0.161	0.024	0.002	0.004	0.057	0.015
2013	0.203	0.161	0.024	0.002	0.004	0.047	0.014
2014	0.201	0.160	0.023	0.002	0.004	0.051	0.016
2015	0.194	0.154	0.023	0.002	0.004	0.050	0.016
2016	0.194	0.148	0.022	0.002	0.003	0.042	0.015
2017	0.184	0.145	0.022	0.002	0.003	0.044	0.017
2018	0.188	0.152	0.021	0.002	0.004	0.044	0.018
2019	0.186	0.155	0.019	0.002	0.004	0.042	0.017
2020	0.181	0.160	0.018	0.001	0.003	0.037	0.014
2021	0.175	0.148	0.017	0.001	0.004	0.031	0.013
2022	0.174	0.146	0.018	0.001	0.004	0.034	0.012
2023	0.170	0.141	0.017	0.001	0.004	0.034	0.012

**Figure 5.24:** *The share of N<sub>2</sub>O emissions by animals within manure management in 2023*



### 5.9.1. Methodological Issues – Methods

**Animal waste management systems (AWMS)** – allocation of manure into AWMS is based on survey on manure management practices used. A questionnaire survey in farms was performed in the cooperation with the NPPC-VÚŽV and other research institutions during the year 2014. Farmers reported the total produced amount of solid and liquid manure and amount of manure, which was processed in anaerobic digesters by regions. This survey defined more accurately numbers of days on pasture for cattle, sheep, goats and horses. Manure left on pasture was estimated based on this data. Time-series was completed by extrapolation. Animal waste management systems will be revised in the next submissions, due to lack of accurate information on abatements in manure management systems.

Allocation according to the climatic conditions is 100% for cool temperate dry climate for all animals based on the IPCC 2019 RF and climate data for the Slovak Republic.

Western Europe default value for nitrogen excretion was used, more information is in the [Chapter 5.5](#).

**Nitrogen excretion rate for cattle** – a country specific nitrogen excretion rate based on tier 2 approach was used. This was implemented for each subcategory of cattle based on statistical inputs - milk yield, weight and daily gain of the animal. The average annual requirements of crude protein for the maintenance, lactation, pregnancy and daily gain were estimated. Milk yield, daily gain and share of proteins in milk at the regional level, were taken from the ŠÚ SR statistics. Average body weights were estimated using the country specific method documented in the [Chapter 5.7.1](#). While the same activity data was used, the calculation model is in line with enteric fermentation model. This methodology was developed in the cooperation with the NPPC-VÚŽV. Additional information regarding maintenance and pregnancy was taken into account. Country specific parameters are documented in [Table 5.54](#).

**Table 5.54: Additional parameters for estimation of nitrogen excretion rate**

NAME OF PARAMETER	PARAMETERS WITH UNITS*	SOURCE
Crude protein per litter of milk	85 g per litter	P. Petrikovič – A. Sommer: Nutrition for Cattle
Share of protein in calf meat	21.5%	J. Keresteš at all.: Biotechnology nutrition and health
Usability for maintenance	2%	P. Petrikovič – A. Sommer: Nutrition for Cattle
Usability for pregnancy	20%	P. Petrikovič – A. Sommer: Nutrition for Cattle
Nitrogen overage -dairy cattle	25%	Expert judgement
Nitrogen overage - other cattle	20%	Expert judgement
Share of protein in beef meat	21%	J. Keresteš at all.: Biotechnology nutrition and health
Conversion factor from CP to N	6.25	IPCC 2019 RF p.10.58
Time without milking	60 days	<a href="https://www.plis.sk/">https://www.plis.sk/</a>
Crude protein for pregnancy begin part of pregnancy	680 g/day	P. Petrikovič – A. Sommer: Nutrition for Cattle
Crude protein for pregnancy begin part of pregnancy	765 g/day	P. Petrikovič – A. Sommer: Nutrition for Cattle

\*consistent in all time-series

The nitrogen excretion rate was determined for the whole time-series with methods according to the publication *P. Petrikovič – A. Sommer: Nutrition for Cattle*.<sup>6</sup> The complex of crude protein contains amount of protein nitrogen and non-protein nitrogen estimated with the Kjeldahl method. Crude protein is multiplied by a conversion factor of 6.25 to dietary nitrogen. The calculation method is based on a reverse estimation of nitrogen excretion from the average parameters of animal production (milk yield and daily gain, body weight) of the cattle. Parameters are multiplied with tabular values of crude protein from individual physiological activities. Subsequently, the partial crude protein from activities is summed to the total crude protein. Total crude protein was recalculated to the nitrogen.

Dairy cattle:

$$CP_{m-Total} = \left[ (4.93 * H^{0.75} * U_m) - \left( \frac{CP_m}{100} * U_m \right) \right]$$

$$CP_{l-Total} = \left[ (MY * CP_l) - \left( \frac{MY * 1000}{100 * SP_l} \right) \right]$$

$$CP_{p-Total} = \frac{C_{p1} + C_{p2}}{100} * U_p$$

<sup>6</sup> Perikovič, P., Sommer, A., 2002, [Nutrition for Cattle](#), The Research Institute for Animal Production, ISBN: 80-88872-21-9

$$\text{Total}_{\text{CP}} = \frac{(\text{CP}_{\text{m-Total}} + \text{CP}_{\text{l-Total}}) * \text{lactation period}}{1000} + \frac{(\text{CP}_{\text{m-Total}} + \text{CP}_{\text{p-Total}}) * \text{time without milking}}{1000} * 365$$

$$N_{\text{intake (T)}} = \left( \frac{\text{Total}_{\text{CP}}}{\frac{100}{6.25}} \right)$$

$$\text{NEX}_{(\text{T})} = N_{\text{intake (T)}} + (N_{\text{intake (T)}} * O_{\text{N}})$$

### Non-dairy cattle:

$$\text{CP}_{\text{m-Total}} = \left[ (4.93 * H^{0.75} * U_{\text{m}}) - \left( \frac{\text{CP}_{\text{m}}}{100} * U_{\text{m}} \right) \right]$$

$$\text{CP}_{\text{dg-Total}} = [(200 + (4.43 * H^{0.75})) * \text{dg}] * \text{SP}_{\text{m}}$$

$$\text{Total}_{\text{CP}} = \frac{(\text{CP}_{\text{m-Total}} + \text{CP}_{\text{dg-Total}})}{1000} * 365$$

$$N_{\text{intake (T)}} = \left( \frac{\text{Total}_{\text{CP}}}{\frac{100}{6.25}} \right)$$

$$\text{NEX}_{(\text{T})} = N_{\text{intake (T)}} + (N_{\text{intake (T)}} * O_{\text{N}})$$

Where: **CP<sub>m-Total</sub>** = crude protein for maintenance in g per day, **H<sup>0.75</sup>** = metabolic body size, **H** = average body weight in kg, **U<sub>m</sub>** = Usability for maintenance in %, **MY** = milk yield in kg/day **CP<sub>l-Total</sub>** = crude protein for lactation g per day, **CP<sub>p-Total</sub>** = crude protein for pregnancy in g per day, **CP<sub>dg-Total</sub>** = crude protein for daily gain in g per day, **dg** = daily gain of animal in kg, **4.93** factor for maintenance, **4.43** factor crude protein per daily gain, **SP<sub>l</sub>** = share of proteins in milk in %, **SP<sub>m</sub>** = share of proteins in meat in %, **lactation period** = period of milk production in days, **intervening period** = is figure indicating the time elapsed between two calves in days, **Total<sub>CP</sub>** = total calculated crude protein in kg, **NEX<sub>(T)</sub>** = annual N excretion rates, kg N animal<sup>-1</sup> year<sup>-1</sup>, **6.25** = conversion from kg of dietary protein to kg dietary N, kg feed protein (kg N)<sup>-1</sup>, **O<sub>N</sub>** = share of overage of nitrogen in N, **N<sub>INTAKE (T)</sub>** = daily N consumed per animal of category T, **CP<sub>p1</sub>** = crude protein for pregnancy begin part of pregnancy **CP<sub>p2</sub>** = crude protein for pregnancy final part of pregnancy

**Nitrogen excretion rate for swine** – a country specific nitrogen excretion rate was used for swine category, based on the tier 2 method from the IPCC 2019 RF. The nitrogen excretion rates were developed based on the nutrient requirements of pigs supported by feed database developed at the NPPC-VÚŽV and expert judgment on the average daily weight gain of particular swine subcategories.

The value of gross energy intake is consistent with the value used in the category 3.B.1.3. Data on gross energy intake were calculated according to publication *Petrikovič et al. (2002): Nutrient requirements of pigs*. The nitrogen intakes were determined from the crude protein estimated to cover requirements for each of subcategories of swine and their gross energy intake.

$$N_{\text{intake (T)}} = \frac{\text{GE}}{18.45} * \left( \frac{\text{CP \%}}{\frac{100}{6.25}} \right)$$

Where: **N<sub>INTAKE (T)</sub>** = daily N consumed per animal of category T, kg N/head/day, **GE** = gross energy intake from feeding ration MJ/animal/day, **18.45** = conversion factor for dietary GE/kg of dry matter MJ/kg, **CP** = percent crude protein in diet %, **6.25** = conversion from kg of dietary protein to kg of dietary N, kg feed protein (kg/N).

The values of the annual nitrogen excretions, rates of N retention, and estimated crude protein content are summarized in **Table 5.55**. The results for poultry for 2023 were presented in **Table 5.56**. Sheep are also significant contributors to emissions, but data about crude protein were unavailable. The N-excretion rates were calculated according to Equation 10.321 of the IPCC 2019 RF:

$$\text{NEX}_{(\text{T})} = N_{\text{intake (T)}} * (1 - N_{\text{retention}})$$

Where:  $NEX_{(T)}$  = annual N excretion rates in kg N/head/yr,  $N_{INTAKE (T)}$  = the annual N intake per head of animal of species/category T, kg N/head/yr,  $N_{RETENTION (T)}$  = fraction of annual N intake that is retained by animal of species (according to Table 10.20 of the IPCC 2019 RF).

**Table 5.55: Country specific regional parameters for poultry for in 2023**

2022		Bratislava	Trnava	Trenčín	Nitra	Banská Bystrica	Žilina	Prešov	Košice
BREEDING BROILERS	N retention (%)	0.300	0.300	0.300	0.300	0.300	0.300	0.300	0.300
	<i>N-intake (kg N animal/day)</i>	0.004	0.004	0.004	0.004	0.004	0.004	0.004	0.004
	<i>N<sub>EX</sub> (kg N/animal/year)</i>	0.898	0.898	0.898	0.898	0.898	0.898	0.898	0.898
DUCKS	N retention (%)	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
	<i>N-intake (kg N animal/day)</i>	0.006	0.006	0.006	0.006	0.006	0.006	0.006	0.006
	<i>N<sub>EX</sub> (kg N/animal/year)</i>	1.295	1.295	1.295	1.295	1.295	1.295	1.295	1.295
FATTENING BROILERS	N retention (%)	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
	<i>N-intake (kg N animal/day)</i>	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003
	<i>N<sub>EX</sub> (kg N/animal/year)</i>	0.739	0.739	0.739	0.739	0.739	0.739	0.739	0.739
GEESE	N retention (%)	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
	<i>N-intake (kg N animal/day)</i>	0.007	0.007	0.007	0.007	0.007	0.007	0.007	0.007
	<i>N<sub>EX</sub> (kg N/animal/year)</i>	1.739	1.739	1.739	1.739	1.739	1.739	1.739	1.739
LAYING HENS INCLUDING COCKS	N retention (%)	0.300	0.300	0.300	0.300	0.300	0.300	0.300	0.300
	<i>N-intake (kg N animal/day)</i>	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003
	<i>N<sub>EX</sub> (kg N/animal/year)</i>	0.794	0.794	0.794	0.794	0.794	0.794	0.794	0.794
TURKEYS	N retention (%)	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003
	<i>N-intake (kg N animal/day)</i>	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010
	<i>N<sub>EX</sub> (kg N/animal/year)</i>	3.529	3.529	3.529	3.529	3.529	3.529	3.529	3.529

**Other animals** – the calculation is based on the determination of body weight. All animals have their specific body weight. This parameter was estimated and is country specific. The body weight parameter is consistent across the time-series and specific for animal species. The NPPC-VÚŽV provided specific

body mass for animals. Annual nitrogen excretion rates were calculated for sheep, goats, horses and poultry. N-excretion rates were calculated based on the IPCC 2019 RF, Equation 10.30:

$$NEX_T = N_{rate(T)} * \frac{TAM}{1000} * 365$$

Where: **N<sub>EXT</sub>** = annual N-excretion for each livestock species respectively category in kg N per animal; **N<sub>RATE(T)</sub>** = default N-excretion rate in kg N (100 kg/animal mass)/day (IPCC 2019 RF), **TAM** = country specific animal mass for each livestock species/category in kg per animal

Direct emissions from manure management systems were estimated according to the following equation:

$$N_2O_{EM} = \left[ \sum \left[ \sum (N * N_{EX} * AWMS) \right] * EF \right] * \frac{44}{28}$$

Where: **N<sub>2O<sub>EM</sub></sub>** = direct N<sub>2</sub>O emissions from manure management in kg N<sub>2</sub>O; **N** = number of livestock species respectively category, **N<sub>EX</sub>** = annual average N-excretion/head of species respectively category in kg N/animal, **AWMS** = percentage of total annual nitrogen excretion for each livestock category, that is managed in manure management systems in the country, **EF** = default emission factor for direct N<sub>2</sub>O emissions from manure management system in kg N<sub>2</sub>O-N/kg N in manure management system, **44/28** = conversion of N<sub>2</sub>O-N emissions to N<sub>2</sub>O emissions

**Table 5.56: Country specific regional parameters for dairy cattle in 1990**

CATEGORIES	N <sub>EX</sub>	Body mass	Liquid	Solid	Pasture	Anaerobic digester
	kg N head/year	kg	%			
Dairy cows Bratislava region	82.63	589	42.85	56.86	0.29	NO
Dairy cows Trnava region	78.69	589	18.57	79.79	1.64	NO
Dairy cows Trenčín region	74.60	589	7.12	86.92	5.97	NO
Dairy cows Nitra region	75.84	589	16.56	82.62	0.82	NO
Dairy cows Žilina region	66.07	589	5.93	75.34	18.73	NO
Dairy cows Banská Bystrica region	71.66	589	10.67	77.88	11.44	NO
Dairy cows Prešov region	62.65	589	4.06	80.43	15.51	NO
Dairy cows Košice region	69.36	589	2.41	86.29	11.30	NO

**Table 5.57: Country specific regional parameters for dairy cattle in 2023**

CATEGORIES	N <sub>EX</sub>	Body mass	Liquid	Solid	Pasture	Anaerobic digester
	kg N head/year	kg	%			
Dairy cows Bratislava region	132.14	600	0.00	99.52	0.48	0.00
Dairy cows Trnava region	131.65	600	8.11	77.00	1.34	13.55
Dairy cows Trenčín region	126.63	600	7.58	77.29	6.18	8.95
Dairy cows Nitra region	128.43	600	16.49	80.47	0.64	2.40
Dairy cows Žilina region	111.66	595	5.89	57.31	30.11	6.70
Dairy cows Banská Bystrica region	113.17	599	13.95	69.60	10.74	5.70
Dairy cows Prešov region	109.43	593	6.35	70.74	20.25	2.67
Dairy cows Košice region	108.72	597	3.04	77.64	10.78	8.55



**Table 5.58: Country specific parameters for poultry in 2023**

CATEGORIES	N <sub>EX</sub>	Body mass	Pasture	Manure poultry without litter	Poultry manure with litter
	kg N head/year	kg	%		
Laying hens including cocks	0.79	2	-	75%	25%
Fattening broilers	0.74	1.02	-	-	100%
Breeding broilers	0.90	1.02	-	-	100%
Turkeys	2.41	3.28	-	-	100%
Geese	1.74	1.53	50%	50%	50%
Ducks	1.30	2.05	50%	50%	50%

**Table 5.59: N<sub>EX</sub> and share (%) for different domestic livestock and share in AWMS in 2023**

CATEGORIES		N <sub>EX</sub>	LIQUID	SOLID	PASTURE	OTHER (LITTER)
		N kg/head	%			
NON-DAIRY CATTLE	Suckler cows	40.35	-	45.21	54.79	-
	Calves in 6 month (milk type)	19.96	-	-	100.00	-
	Heifer (milk type)	38.42	-	97.55	2.45	-
	Heifer (pregnant) (milk type)	57.56	-	97.55	2.45	-
	Fattening (milk type)	46.48	10.00	90.00	-	-
	Oxen (milk type)	93.79	-	100.00	-	-
	Breeding bull (milk type)	66.21	-	100.00	-	-
	Calves in 6 month (beef type)	12.18	-	40.00	60.00	-
	Heifer (beef type)	38.29	-	45.21	54.79	-
	Heifer (pregnant) (beef type)	54.73	-	45.21	54.79	-
	Fattening (beef type)	42.61	20.00	80.00	-	-
	Oxen (beef type)	68.41	-	100.00	-	-
	Breeding bull (beef type)	43.35	-	75.34	24.66	-
	2023*	41.16	2,46	71,17	26,37	-
SHEEP	Mature ewes (milk type)	7.88	-	49.59	50.41	-
	Mature ewes (beef type)	9.20	-	45.20	54.80	-
	2023*	8.36	-	48,03	51,97	-
	Growing lambs (milk type)	4.27	-	49.59	50.41	-
	Growing lambs pregnant (milk type)	7.23	-	49.59	50.41	-
	Growing lambs (beef type)	6.24	-	45.21	54.79	-
	Growing lambs pregnant (beef type)	8.54	-	45.21	54.79	-
	2023*	5.90	-	48.08	51,92	-
	Rams (milk type)	10.51	-	100.00	-	-
	Rams (beef type)	11.83	-	100.00	-	-
	2023*	10.99	-	81.63	18.37	-
GOATS	Mature female goats	9.23	-	49.60	50.40	-
	Pregnant goats	7.98	-	49.60	50.40	-
	Other mature goats	3.61	-	49.60	50.40	-
	2023	7.82	-	49.60	50.40	-
HORSES	Young horses up to 1 year	17.32	70.00	-	30.00	-
	Young horses from 1 to 3 year	39.858	70.00	-	30.00	-
	Castrated horses	66.43	70.00	-	30.00	-
	Stallions	52.20	70.00	-	30.00	-
	Mares	47.45	70.00	-	30.00	-
	2023*	50.78	70.00	-	30.00	-

\*weighted average

The IPCC default emission factors for N<sub>2</sub>O emissions estimation per AWMS are based on the Table 10.21 of the IPCC 2019 RF ([Table 5.60](#)).

**Table 5.60: Emission factors for N<sub>2</sub>O emissions used in manure management in 2023**

MANURE MANAGEMENT SYSTEMS	EFs (N <sub>2</sub> O-N)
	kg N <sub>2</sub> O-N/kg N <sub>EX</sub>
Solid storage and dry lot	0.01
Liquid system	0.005
Anaerobic digesters	0.0006
Cattle and Swine deep bedding	0.01
Poultry manure with litter	0.001
Poultry manure without litter	0.001

### 5.9.2. Activity Data

The NPPC-VÚŽV is a data provider for animal housing, pasture, and production of manures and slurries. More information on animal numbers can be found in the previous chapters.

## 5.10. Indirect N<sub>2</sub>O Emissions from Manure Management (CRF 3.B.2.5)

### 5.10.1 Volatilisation from Manure Management Systems

Indirect emissions result from volatile nitrogen losses that occur primarily in the form of ammonia and NO<sub>x</sub>. The fraction of excreted organic nitrogen that is mineralized to ammonia nitrogen during manure storage depends mainly on time and temperature. Simple forms of organic nitrogen such as urea and uric acid are rapidly mineralized to ammonia nitrogen, which is highly volatile and easily diffused into the surrounding air. Nitrogen losses begin with excretion in housing and continue with on-site management in storage and treatment systems. Pasture losses are considered separately in emissions from managed soils.

#### Methodological Issues – Methods

Tier 1 approach of the IPCC 2019 RF for nitrogen estimation of N volatilization in forms of NH<sub>3</sub> and NO<sub>x</sub> from manure management systems is based on multiplication of the amount of nitrogen excreted from all livestock categories and managed in each manure management systems by a fraction of volatilized nitrogen. N losses were then summed from all manure management systems. Emission factor is 0.01 kg NH<sub>3</sub>-N and NO-N for N<sub>2</sub>O emissions from atmospheric deposition of nitrogen. The losses were calculated for all farm animals. Calculations were performed using the following equations:

$$N_{\text{volatilization-MMS}} = \sum_S \left[ \sum_T \left[ \left( (N_T * Nex_T) * AWMS_{T,S} \right) + N_{cdg(s)} \right] * FraC_{GasMS(T,S)} \right]$$

$$N_2O_{MM} = (N_{\text{volatilization-MMS}} * EF) * \frac{44}{28}$$

Where: **N<sub>T</sub>** = number of head of farm animals' species/category, **Nex<sub>T</sub>** = annual average N excretion per head of species respectively category in kg N per animal, **MS<sub>T,s</sub>** = fraction of total annual nitrogen excretion for each farm animals' species respectively category, that is managed in manure management systems, **Frac<sub>GasMS</sub>** = percent of managed manure nitrogen for livestock category T that volatilizes as NH<sub>3</sub> and NO<sub>x</sub> in the manure management systems S in %.

## Activity data

Volatilized nitrogen (NH<sub>3</sub> and NO<sub>x</sub>) from animal waste was 21 965,587 t of N, which represents 0.345 Gg of N<sub>2</sub>O in 2021. Activity data in this category are consistent with the activity data used in animal manure. **Table 5.61** shows the time series of input data and emissions.

**Table 5.61: Input parameters and EFs in category 3.B.2.5 - Atmospheric Deposition in particular years**

YEAR	VOLATILIZED N FROM ANIMAL MANURE	IEF	N <sub>2</sub> O EMISSIONS
	kg	kg N <sub>2</sub> O-N/kg N	Gg
1990	43 190 242.97	0.01	0.679
1995	21 805 137.56	0.01	0.343
2000	21 710 544.69	0.01	0.341
2005	22 276 783.88	0.01	0.350
2010	22 150 081.79	0.01	0.348
2011	23 472 237.71	0.01	0.369
2012	21 597 635.43	0.01	0.339
2013	23 262 404.75	0.01	0.366
2014	24 407 797.38	0.01	0.384
2015	23 254 916.50	0.01	0.365
2016	24 688 735.51	0.01	0.388
2017	24 248 249.73	0.01	0.381
2018	25 243 418.16	0.01	0.397
2019	25 101 736.14	0.01	0.394
2020	24 370 521.93	0.01	0.383
2021	24 254 243.29	0.01	0.381
2022	22 728 563.82	0.01	0.357
2023	21 965 587.31	0.01	0.345

## 5.10.2. Nitrogen Leaching and Run-off from Manure Management Systems

This category was included in the inventory for the first time this year based on the implementation of the IPCC2019 RF. The new methodological guidelines provides the default values of  $Frac_{LeachMS}$ . The default values were adopted and N<sub>2</sub>O emission was possible to estimate.

### Methodological Issues – Methods

Tier 1 approach of the IPCC 2019 RF for nitrogen estimation of N leaching and run-off from manure management systems is based on multiplication of the amount of nitrogen excreted from all livestock categories and managed in each manure management systems by a fraction of volatilized nitrogen. N losses were then summed from all manure management systems. Emission factor is 0.011 kg N<sub>2</sub>O–N (kg N leaching/runoff)<sup>-1</sup>. The losses were calculated for all farm animals.  $N_{cdg(s)}$  is defined as amount of nitrogen from co-digesters added to biogas plants such as food wastes or purpose grown crops. National data about this activity is missing, therefore value was neglected. Calculations were performed using the following equations:

$$N_{leaching-MMS} = \sum_S \left[ \sum_{T,P} \left[ (N_T * Nex_T * AWMS_{T,S}) + N_{cdg(s)} * Frac_{LeachMS(T,S)} \right] \right]$$

$$N_2O_{MM} = (N_{\text{leaching-MMS}} * EF_5) * \frac{44}{28}$$

Where:  $N_T$  = number of head of farm animals' species/category,  $N_{exT}$  = annual average N excretion per head of species respectively category in kg N per animal,  $AWMS_{T,s}$  = fraction of total annual nitrogen excretion for each farm animals' species respectively category, that is managed in manure management systems,  $Frac_{leachMS}$  = percent of managed manure nitrogen for livestock category T that volatilizes as  $NH_3$  and  $NO_x$  in the manure management systems S in %,  $N_{cdg(s)}$  = amount of nitrogen from co-digestates added to biogas plants such as food wastes or purpose grown crops,  $kg\ N\ yr^{-1}$

### Activity data

N lost through leaching and run-off from animal waste was 660,846 t of N, which represents 0.011 Gg of  $N_2O$  in 2023. Activity data in this category are consistent with the activity data used in animal manure.

**Table 5.62** shows the time series of input data and emissions.

**Table 5.62: Input parameters and EFs in category 3.B.2.5 - Nitrogen leaching and run-off in particular years**

YEAR	N LOST THROUGH LEACHING AND RUN-OFF	IEF	N <sub>2</sub> O EMISSIONS
	kg	kg N <sub>2</sub> O-N/kg N	Gg
1990	1 572 714.68	0.011	0.027
1995	1 083 647.03	0.011	0.019
2000	945 488.86	0.011	0.016
2005	853 634.74	0.011	0.015
2010	748 176.34	0.011	0.013
2011	741 600.21	0.011	0.013
2012	749 871.46	0.011	0.013
2013	730 650.19	0.011	0.013
2014	729 524.16	0.011	0.013
2015	714 108.59	0.011	0.012
2016	701 794.66	0.011	0.012
2017	687 485.67	0.011	0.012
2018	697 645.35	0.011	0.012
2019	698 571.26	0.011	0.012
2020	691 860.87	0.011	0.012
2021	665 886.84	0.011	0.012
2022	662 224.41	0.011	0.011
2023	660 846.60	0.011	0.011

## 5.11. Rice Cultivation (CRF 3.C)

No emissions from rise cultivation were estimated because this activity did not occur in the Slovak Republic in 1990 – 2023. Therefore, notation keys NO were used in all time-series.

## 5.12. Agricultural Soils (CRF 3.D)

EMITTED GAS: N<sub>2</sub>O

METHODS: TIER 1, TIER 2

EMISSION FACTORS: CS, D

KEY SOURCES: YES

PARTICULARLY SIGNIFICANT SUBCATEGORIES: SYNTHETIC FERTILIZERS

Direct emissions are the primary source of N<sub>2</sub>O in the Slovak inventory. In 2023, 66% of the national total N<sub>2</sub>O emissions originated from this category, which includes N inputs from synthetic N-fertilizer, organic manures as animal manure use, sewage sludge application and compost, emissions from urine and dung N deposited on pasture and crop residues. Trend of total N<sub>2</sub>O emissions from the **Agriculture sector** reflects trend of direct emissions from cultivated soil, emissions from applied manure and indirect emissions from leaching and deposition of ammonia and NO<sub>x</sub>. The productivity of different categories of domestic livestock varies significantly depending on the scale and the production level of farms in different regions. In the Slovak Republic, both the extensive and intensive farming systems in animal husbandry can be found. Nitrogen inputs can differ from the calculations in the range of ±10%.

Total N<sub>2</sub>O emissions from agricultural soils were 2,88 Gg of N<sub>2</sub>O in 2023. The emissions increased by 37% in comparison with 2022 and decreased by 46% in comparison with the base year 1990 (**Table 5.63**). The major reason for the overall decreasing trend is a sharp decrease in the use of synthetic fertilizers in early 90-ties and the continual decrease in the use of animal manure caused by the reduction in the number of animals (**Figure 5.25**). **Figure 5.25** shows, that since 1999 the trend is stable with the small fluctuations caused by changes in animal population and inter-annual differences in categories 3.D.1.4 - Crop Residues, 3.D.1 - Inorganic Nitrogen Fertilizers and 3.D.2 - Indirect N<sub>2</sub>O Emissions. No emissions are reported in the categories 3.D.1.6 - Cultivation of Organic Soils. More information is available in the **Chapter 5.12.8**.

**Table 5.63:** N<sub>2</sub>O emissions (Gg) in 3.D - Direct Soils according to the subcategories in particular years

YEAR	3.D.1 DIRECT N <sub>2</sub> O EMISSIONS FROM MANAGED SOIL					3.D.2 INDIRECT N <sub>2</sub> O EMISSIONS FROM MANAGED SOIL	
	3.D.1.1 Synthetic fertilizers	3.D.1.2. Organic N-fertilizers	3.D.1.3 Urine and dung deposited by grazing animal	3.D.1.4 Crop residues	3.D.1.5 N in mineral soils that is mineralized/ immobilized in association with loss of soil C	3.D.2.1 Atmospheric deposition	3.D.2.2 Nitrogen leaching and runoff
1990	1.746	0.572	0.091	0.617	0.0002	0.679	1.628
1995	0.547	0.423	0.073	0.517	0.0008	0.343	0.766
2000	0.665	0.378	0.056	0.320	0.0012	0.341	0.600
2005	0.784	0.338	0.056	0.457	0.0016	0.350	0.857
2010	0.837	0.299	0.060	0.346	0.0016	0.348	1.270
2011	0.947	0.291	0.060	0.456	0.0016	0.369	0.218
2012	0.794	0.298	0.062	0.389	0.0015	0.339	0.223
2013	0.892	0.308	0.063	0.441	0.0014	0.366	0.557
2014	0.935	0.326	0.065	0.569	0.0013	0.384	0.819
2015	0.902	0.300	0.065	0.473	0.0013	0.365	0.106
2016	0.992	0.307	0.065	0.593	0.0013	0.388	0.703
2017	0.963	0.307	0.063	0.462	0.0012	0.381	0.375
2018	1.013	0.314	0.067	0.510	0.0011	0.397	0.100
2019	1.010	0.312	0.066	0.518	0.0010	0.394	0.377

YEAR	3.D.1 DIRECT N <sub>2</sub> O EMISSIONS FROM MANAGED SOIL					3.D.2 INDIRECT N <sub>2</sub> O EMISSIONS FROM MANAGED SOIL	
	3.D.1.1 Synthetic fertilizers	3.D.1.2 Organic N-fertilizers	3.D.1.3 Urine and dung deposited by grazing animal	3.D.1.4 Crop residues	3.D.1.5 N in mineral soils that is mineralized/immobilized in association with loss of soil C	3.D.2.1 Atmospheric deposition	3.D.2.2 Nitrogen leaching and run-off
2020	1.003	0.289	0.066	0.552	0.0009	0.383	0.530
2021	1.002	0.283	0.067	0.524	0.0007	0.381	0.230
2022	0.906	0.276	0.068	0.436	0.0007	0.357	0.058
2023	0.845	0.280	0.067	0.545	0.0007	0.345	0.797

Figure 5.25: Trend in N<sub>2</sub>O emissions (kt) by subcategories within agricultural soils in 1990 – 2023





Figure 5.26 shows, that major share of emissions belongs to synthetic fertilizers use (29%), crop residues (19%), organic nitrogen fertilizers (9,7%) and indirect emissions from agricultural soils (39,7%).

Figure 5.26: The share of aggregated emissions by categories within agricultural soils in 2023



### 5.12.1. Inorganic Fertilizers (CRF 3.D.1.1)

The applied amounts of synthetic fertilizers into cultivated soils decreased in the last 15 years. Nowadays, the amount of synthetic fertilizers applied to the agricultural soils has increased again. This fact is the main driver in increasing emissions in the sector. The potential for the volatilization of ammonia emissions can vary in a very large range. The best information on NH<sub>3</sub> emissions from cultivated soils in the Slovak Republic is based on the applied nitrogen fertilizers. Emissions also depend on the type of fertilizers, soil parameters (pH), meteorological conditions, application technics and time of fertilizers application in relation to crop development. Information on applied nitrogen fertilizers was provided by the ŠÚ SR.

#### Methodological issues - method

Default emission factor was used from the IPCC 2019 RF (0.005 kg N<sub>2</sub>O–N/kg N). Total N<sub>2</sub>O emissions from using the synthetic fertilizers were 0.845 Gg in 2023. Tier 1 method was applied in combination with the default EF. According to the prioritization plan, Tier 2 approach will be implemented in 2025. Implementation is not processed yet due to missing geographical data on inorganic N fertilizer consumption (including Urea application).

#### Activity data

The Central Control and Testing Institute in Agriculture (UKSÚP) provided the data annually into the SHMÚ based on cooperation agreement between the both institutions. The UKSÚP collected data on farm level electronically. The farmers are obliged to report the amount of applied nitrogen into the UKSÚP each year. The UKSÚP as administrator of databases makes validation of data each year.

The consumption of synthetic fertilizers decreased during the last decade of the 20<sup>th</sup> century, from 222,3 kt in 1990 to 107,6 kt in 2023 (52%). On the other hand, consumption of the synthetic fertilizers increased by 8% in 2023 compared to 2005 and decreased by almost 7% in comparison with the year 2022. Decreasing numbers of domestic livestock caused the demand for inorganic nitrogen. Higher consumption of synthetic fertilizers compensates missing organic nitrogen in soils.

Activity data on N input from the application of inorganic fertilizers to agricultural soils is summarized in [Table 5.64](#).

**Table 5.64: Input parameters and EFs in 3.D.1.1 - Inorganic N-Fertilizers in particular years**

YEAR	N-INPUT IN FERTILIZERS	EFs	N <sub>2</sub> O EMISSIONS
	t	kg N <sub>2</sub> O-N/kg N	KT
1990	222 255.00	0.005	1.746
1995	69 587.00	0.005	0.547
2000	84 609.00	0.005	0.665
2005	99 760.00	0.005	0.784
2010	106 513.00	0.005	0.837
2011	120 555.00	0.005	0.947
2012	101 004.00	0.005	0.794
2013	113 581.39	0.005	0.892
2014	119 036.05	0.005	0.935
2015	114 773.00	0.005	0.902
2016	126 235.77	0.005	0.992
2017	122 541.15	0.005	0.963
2018	128 976.88	0.005	1.013
2019	128 532.97	0.005	1.010
2020	127 676.52	0.005	1.003
2021	127 494.60	0.005	1.002
2022	115 346.78	0.005	0.906

YEAR	N-INPUT IN FERTILIZERS	EFs	N <sub>2</sub> O EMISSIONS
	<i>t</i>	<i>kg N<sub>2</sub>O-N/kg N</i>	<i>KT</i>
2023	107 607.31	0.005	0.845

### 5.12.2. Animal Manure Applied to Soil (CRF 3.D.1.2.a)

As domestic livestock produces a different kind of nitrogen inputs (liquid or solid) into the ecosystem, also the structure of domestic livestock is important (the ratio of different categories of domestic livestock) as well as the emissions from the AWMS. In addition, the production of nitrogen per head per year also plays a certain role.

#### Methodological issues – method

Managed manure nitrogen, available for application to managed soil (NMMS\_Avb) was calculated based on the Equations 10.34(update), 10.34<sub>A</sub>, 10.34<sub>B</sub> (IPCC 2019 RF).

Losses are defined as losses of following gases N<sub>2</sub>, NH<sub>3</sub>, NO<sub>x</sub> and N<sub>2</sub>O. Losses are calculated according to the 2019 IPCC GL from the total amount of liquid, deep bedding and manure managed in anaerobic digesters. Losses as Frac<sub>lossMS</sub> used for managed manure as are calculated in line with 3.B.2.5 categories and fractions were calculated from these both categories (equation 10.34<sub>A</sub>). Fractions (Frac<sub>FEED</sub>, Frac<sub>FUEL</sub>, Frac<sub>CNST</sub>) in the Equation 11.4 (IPCC 2019 RF) are considered zero. Managed manure nitrogen available for application to managed soils (NMMS\_Avb) was calculated based on Equation 10.34 (IPCC 2019 RF). The case of straw-based systems N inputs with straw were also taken into account in the inventory according to the above mentioned equation. Straw N from pigs and poultry for deep litter was considered. The Hungarian value for poultry nitrogen content was used due to absent country specific study concerning of nitrogen content from bedding materials. The Hungary is neighbouring country with similar climatic and agricultural conditions.

**Table 5.65: Nitrogen in bedding materials by animal category and manure management systems**

ANIMAL CATEGORY	N-CONTENT OF BEDDING MATERIALS BY MANURE MANAGEMENT SYSTEMS (kg N/head)	SOURCES
	DEEP LITTER	
Market swine	1.6	p. 10.66 of the IPCC 2006 GL
Poultry*	0.022	<u>Expert judgement in accordance with Hungary inventory</u>

\*Poultry manure with bedding

The calculated amount of nitrogen input from animal waste applied to soil was 32 218,39 t/N/year when the default EF = 0.005 kg N<sub>2</sub>O-N/kg N was used. Total amount of N<sub>2</sub>O emissions from animal excreta applied to soil was 0.253 Gg in 2023.

**Table 5.66: Input parameters and EFs in the category 3.D.1.2.a - Animal Manure in particular years**

YEAR	Total nitrogen from MM	Fraction of leached N	Fraction of volatilized nitrogen	Nitrogen from bedding materials (pigs, poultry)	N input from manure applied to soils	EFs	Emissions
	<i>tons N/Year</i>	%	%	<i>tons N/Year</i>	<i>tons N/ Year</i>	<i>kg N<sub>2</sub>O-N/kg</i>	<i>Kt</i>
1990	119 824.42	0.013	0.348	415.531	71 987.36	0.005	0.566
1995	85 594.07	0.013	0.330	319.524	53 167.28	0.005	0.418
2000	74 201.55	0.013	0.316	309.934	47 137.97	0.005	0.370
2005	66 523.80	0.013	0.310	299.021	42 667.20	0.005	0.335
2010	57 544.89	0.013	0.302	233.091	37 286.06	0.005	0.293



YEAR	Total nitrogen from MM	Fraction of leached N	Fraction of volatilized nitrogen	Nitrogen from bedding materials (pigs, poultry)	N input from manure applied to soils	EFs	Emissions
	tons N/Year	%	%	tons N/Year	tons N/ Year	kg N <sub>2</sub> O-N/kg	Kt
2011	55 087.88	0.013	0.299	190.096	35 698.56	0.005	0.280
2012	56 388.78	0.013	0.301	202.753	36 511.78	0.005	0.287
2013	55 040.13	0.013	0.301	192.604	35 631.12	0.005	0.280
2014	56 566.76	0.013	0.304	226.830	36 565.43	0.005	0.287
2015	55 941.57	0.013	0.304	228.504	36 184.97	0.005	0.284
2016	53 940.91	0.013	0.302	207.206	34 955.42	0.005	0.275
2017	54 359.47	0.013	0.303	239.565	35 245.13	0.005	0.277
2018	55 684.09	0.013	0.305	253.338	36 013.22	0.005	0.283
2019	54 536.96	0.013	0.304	240.214	35 267.80	0.005	0.277
2020	51 811.87	0.013	0.300	221.479	33 605.72	0.005	0.264
2021	49 627.48	0.013	0.298	212.643	32 304.52	0.005	0.254
2022	48 902.61	0.014	0.296	181.063	31 877.62	0.005	0.250
2023	49 174.55	0.013	0.293	197.405	32 218.39	0.005	0.253

### Activity data

Livestock number ([Chapter 5.7.2](#)) and information on the AWMS are described in the [Chapter 5.9.1](#). Direct inputs of nitrogen slightly vary according to the applied methodology. Based on the IPCC 2019 RF, total nitrogen excretion per liquid (4 624 t/N/year), per digesters (1489 t/N/year) and solid system (21 291 t/N/year) in manure management were used for the estimation of total nitrogen input of manure applied to soil in 2023.

### 5.12.3. Sewage Sludge Applied to Soils (CRF 3.D.1.2.b)

Reduction of organic matter in the soil depends on the continuous decline of livestock production. The lack of organic fertilizers causes pressure to find alternative sources of organic fertilizers. Sewage sludge is one of the ways to resolve this issue. Sludge is a potential source of nutrients and organic matter. Sewage sludge must be stabilized and afterward applied to the soils. Sludge must be treated biologically, chemically or by heat, long-term storage or any other appropriate process. These processes cause a significant reduction in health risks and save the environment. Act No 188/2003 Coll. on application of sewage sludge and bottom sediments into soil regulates the application of sludge to agricultural soils. Sludge from domestic or urban treatment plants can be applied to agricultural soils.

#### Methodological issues – method

Tier 1 and default emission factor were used (0.005 kg N<sub>2</sub>O-N/kg N) for the estimation of direct N<sub>2</sub>O emissions from sewage sludge applied to soils.

The methodology is in accordance with the IPCC 2019 RF. Emissions were estimated by using these equations:

$$N_2O - N_{\text{sewage sludge}} = N_{\text{sewage sludge}} * P_N \text{ and } N_2O_{\text{sewage sludge}} = N_2O - N_{\text{sewage sludge}} * EF * \frac{44}{28}$$

Where: **N<sub>2</sub>O-N<sub>sewage sludge</sub>** = input of pure nitrogen from sewage sludge applied into the soil in kg, **N<sub>sewage sludge</sub>** = amount of sludge from wastewater treatment in kg, **P<sub>N</sub>** = weighted percentage of nitrogen from sewage sludge (3.31%), **EF** = default emission factor in kg N<sub>2</sub>O-N/kg N


**Table 5.67: Input parameters and EFs used in the category 3.D.1.2.b - Sewage Sludge in particular years**

YEAR	MUNICIPAL SLUDGE	INDUSTRIAL SLUDGE	INPUT INTO SOIL	N-INPUT FROM SEWAGE SLUDGE	N <sub>2</sub> O EMISSIONS
	t				Gg
1990	6 832	3 160	9 991.900	330.732	0.0025986
1995	4 043	2 251	6 294.400	208.345	0.0016370
2000	1 254	1 342	2 596.900	85.957	0.0006754
2005	5 870	2 231	8 101.027	268.144	0.0021068
2010	923	1 102	2 024.855	67.023	0.0005266
2011	358	685	1 043.384	34.536	0.0002714
2012	1 254	478	1 732.337	57.340	0.0004505
2013	518	627	1 145.022	37.900	0.0002978
2014	8	688	695.500	23.021	0.0001809
2015	0	813	812.649	26.899	0.0002113
2016	0	1 134	1 133.631	37.523	0.0002948
2017	0	362	362.151	11.987	0.0000942
2018	0	287	287.404	9.513	0.0000747
2019	0	49	48.937	1.620	0.0000127
2020	0	1	0.968	0.032	0.0000003
2021	0	1	1.000	0.033	0.0000003
2022	0	1	1.000	0.033	0.0000003
2023	NO	NO	NO	NO	NO

#### Activity data

Activity data on sewage sludge consumption in agriculture ([Table 5.70](#)) is based on the data provided by the Water Research Institute (WRP) (applied sludge from municipal wastewater treatment plants) and the Ministry of Environment of the Slovak Republic (Industrial sludge). In 2022 submission, industrial sludge was implemented into inventory for the first time. The WRP collects data on nitrogen inputs (bottom up approach) into the soils. The Water Research Institute informed, that municipal sewage sludge was not applied into agricultural soils in years 2015 – 2023, therefore notation key NO was used. Application of NO notation key was extended to other sources of sludge as there has been information considered from the WRP that there was no application of sewage sludge to agricultural soils in 2023. The data are consistent with the Waste sector. Missing data were extrapolated to enhance completeness before the year 2003 (municipal sewage sludge) and 2002 (Industrial sewage sludge), due to unavailable statistics. Percentage of pure nitrogen from sewage sludge was provided by the [Guidelines for the Sewage Sludge Application](#) by the Soil Science and Conservation Research Institute. According to the mentioned publication, the sludge contains 3.31% of the nitrogen.

**Figure 5.27:** The map of sensitive areas of the Slovak Republic, where application of sludge is prohibited according to the Nitrate directive



#### 5.12.4. Other Organic Fertilizers Applied to Soils (CRF 3.D.1.2.c)

Compost is organic matter that has been decomposed in a process called composting. This process recycles various organic materials otherwise regarded as waste products and produces a soil fertilizer. It is used, for example, in gardens, landscaping, horticulture, urban agriculture and organic farming. The compost is beneficial for the land in many ways, including as a soil fertilizer, addition of vital humus or humic acids, and as a natural pesticide for soil. In ecosystems, compost is useful for erosion control, land and stream reclamation, wetland construction, and as landfill cover.

##### Methodological issues – method

Tier 1 (IPCC 2019 RF) and default emission factor (0.005 kg N<sub>2</sub>O-N/kg N) were used for the estimation of direct N<sub>2</sub>O emissions from compost applied to soils. Emissions were estimated, by using these equations:

$$N_2O - N_{\text{compost}} = N_{\text{compost}} * P_N; N_2O_{\text{compost}} = N_2O - N_{\text{compost}} * EF * \frac{44}{28}$$

Where: **N<sub>2</sub>O-N<sub>compost</sub>** = input of pure nitrogen in compost applied in to the soil in kg, **N<sub>compost</sub>** = amount of compost from composting plant, **P<sub>N</sub>** = Share of nitrogen in organic waste, **EF** = 0.01 kg N<sub>2</sub>O-N/kg N (default).

**Table 5.68:** Share of pure nitrogen from other nitrogen fertilizers in %

TYPE OF FERTILIZERS	P <sub>N</sub>	SOURCES
	%	
Fugate	0.38	<a href="https://nasepole.sk/digestat-vo-vyzive-a-hnojeni-repy/">https://nasepole.sk/digestat-vo-vyzive-a-hnojeni-repy/</a>
Compost	0.7	ÚKSÚP
Natural harmony (organic waste from pharmaceutical production)	2.4	ÚKSÚP
Strae	0.82	<a href="https://nasepole.sk/dusikate-hnojenie-po-zbere-obilnin/">https://nasepole.sk/dusikate-hnojenie-po-zbere-obilnin/</a>
Vitahum (organic - humus fertilizer made from natural substances)	0.7	ÚKSÚP
Green fertilizers	0.5	ÚKSÚP

##### Activity data

Other organic fertilizers applied to soils include the composted waste, digested slurry from digesters, compost and vitahum, natural harmony and green fertilizers. The Consumption is provided with total amount of organic waste into soils (OW) and the data (**Table 5.11, Chapter 5.4**) is provided by the UKSÚP. The Data are converted into nitrogen content (NC).

Data is available from 2000 to 2023. Other organic nitrogen fertilizers were applied to the soil even before the year 2000, but there are no available statistics. Missing data was extrapolated by linear extrapolation in excel spreadsheets.

### 5.12.5. Urine and Dung Deposited by Grazing Animals (CRF 3.D.1.3)

Pasture is typical for some livestock categories. Animals as sheep, goats, horses and cattle (not dairy) are mainly grazed during spring, summer and autumn in the small farms. Animals are housed during the winter. In 2024 submission for the first time pasture from poultry was introduced. Geese and Ducks are grazed 183 days per year.

#### Methodological issues – method

The N<sub>2</sub>O estimation from pasture is based on default emission factors (0.004 kg N<sub>2</sub>O-N/kg N for cattle and poultry and 0.003 kg N<sub>2</sub>O-N/kg N for sheep and other animals). Nitrogen excretions per AWMS were estimated in manure management category. Total nitrogen from pasture was 12 640,72 tN/year in 2023. Total N<sub>2</sub>O emissions from pasture were 0.067 Gg of N<sub>2</sub>O in 2023. This category is estimated in conjunction with the category 3.B.2.

**Table 5.69: Input parameters and EFs in the category 3.D.1.3 - Urine and Dung Deposited by Grazing Animals in particular years**

YEAR	N-EXCRETION ON PASTURE	EFs	N <sub>2</sub> O EMISSIONS
	t N/year	kg N <sub>2</sub> O-N/kg N	Gg
1990	16 438.34	0.0035	0.091
1995	13 554.47	0.0034	0.073
2000	10 911.09	0.0033	0.056
2005	10 785.91	0.0033	0.056
2010	11 653.98	0.0033	0.060
2011	11 578.97	0.0033	0.060
2012	12 029.77	0.0033	0.062
2013	12 114.88	0.0033	0.063
2014	12 375.59	0.0033	0.065
2015	12 435.84	0.0033	0.065
2016	12 342.91	0.0033	0.065
2017	12 145.47	0.0033	0.063
2018	12 692.94	0.0034	0.067
2019	12 553.32	0.0034	0.066
2020	12 380.31	0.0034	0.066
2021	12 660.56	0.0034	0.067
2022	12 746.87	0.0034	0.068
2023	12 640.72	0.0034	0.067

#### Activity data

It is assumed that sheep, goats and horses can stay on pasture for 200 days, 41% of non-dairy cattle stays only for 150 days. The statistical research concerning the amount of pastoral biomass consumed by breeding animals is currently unavailable in Slovakia.

Results of the analysis of different AWMS were used for the calculation of nitrogen input from animal husbandry into N-cycle. This analysis was based on the results collected from questionnaires of the 222 agricultural subjects (21.3% of total subjects in Slovakia). These subjects cultivated 14.7% of total agricultural land and 15.2% of arable land. Duration of the grazing period can vary significantly depending on weather conditions and regions. Reliable data for statistical evaluation is not available, but significant differences can be found in this regard. N<sub>2</sub>O emissions from pasture were based on the proportion of the pasture for housing that was made by the NPPC-VÚŽV. The proportions of the pasture

are demonstrated in the [Chapter 5.9.1](#). Number of animals are summarized in [Table 5.7](#). Activity data in this category are consistent with the activity data used for estimation in category 3.B.2.

### 5.12.6. Crop Residue (CRF 3.D.1.4)

Directly after incorporation of the crop residues into the soil, the multilateral interactions between organic compounds and nutrients present in the residues with the mineral and organic components of soil take place. The knowledge of nutrient potential in crop residues by crop rotation are mostly actual in the present requirements of sustainable land use - greening in plant production. Incorporation of the crop residues into the soil is used as sustainable agricultural practice, due to high nutrition potential.

**Table 5.70: Input parameters and EFs in the category 3.D.1.4 - Crop Residue in particular years**

YEAR	HARVESTED AREA	CROP (т)	CROP RESIDUES	EFs	N <sub>2</sub> O EMISSIONS
	ha	kg d.m./ha	kg N/year	kg N <sub>2</sub> O-N/kg N	Gg
1990	2 147 737.00	67 462.00	78 466.26	0.005	0.617
1995	2 152 852.00	63 385.60	65 755.82	0.005	0.517
2000	2 080 004.00	45 812.10	40 680.82	0.005	0.320
2005	1 721 125.00	68 071.30	58 191.21	0.005	0.457
2010	1 617 786.00	54 869.50	44 086.33	0.005	0.346
2011	1 680 333.00	71 665.70	58 053.86	0.005	0.456
2012	1 703 613.00	63 316.20	49 566.28	0.005	0.389
2013	1 716 326.00	63 795.80	56 186.63	0.005	0.441
2014	1 745 299.00	79 312.00	72 472.34	0.005	0.569
2015	1 728 043.00	66 539.90	60 237.61	0.005	0.473
2016	1 717 480.00	85 742.50	75 505.66	0.005	0.593
2017	1 722 049.00	67 594.50	58 749.09	0.005	0.462
2018	1 725 424.00	76 862.98	64 962.87	0.005	0.510
2019	1 750 468.00	76 280.40	65 934.37	0.005	0.518
2020	1 736 499.00	81 025.73	70 272.93	0.005	0.552
2021	1 741 541.00	74 002.50	66 749.56	0.005	0.524
2022	1 733 440.00	62 542.82	55 418.40	0.005	0.436
2023	1 663 263.00	78 338.14	69 378.87	0.005	0.545

Total N<sub>2</sub>O emissions from crop residues represented 0.545 Gg of N<sub>2</sub>O from 69 378 870 kg of nitrogen in crop residues returned to soils in 2023. Total harvested area (wheat, ray, barley, oat, maize, potato, sugar beet, oil plants, tobacco, maize for silage, leguminous, fodder leguminous, soya, meadows) decreased in comparison with the previous year. In 2023, harvested area was 1 663.26 kha.

#### Methodological issues – method

According to the 2019 IPCC RF, nitrogen input from crop residues was estimated used by equation 11.6 p.11.16.

There is no comprehensive survey on the amount of crop residues burned as fuel in the Slovak Republic. Therefore, no removal from the burning of fuel was assumed. Also, data on fraction of above-ground residues of crop removed annually for a purpose such as feed bedding and construction is not available. The seams and leaves are usually utilized as a fodder of domestic livestock. Data on straw exported abroad are missing.

**Country specific nutrition potential:** The country specific value for sugar beet regarding potential nitrogen nutrition was considered instead of the IPCC default method which is not accurate for the Slovak conditions. According to the national publication [Postharvest residues of sugar beet and their role in the nutrient cycle by Stanislav Torma](#), 20 kg N/ha for sugar beet was taken as country specific value. The default values were considered for other crops. The values are presented in [Table 5.71](#).

**Table 5.71: Parameters used to estimate emissions from crop residues**

CROP TYPE	N <sub>(AG)</sub>	N <sub>(BG)</sub>	SLOPE	INTERCEPT	RS <sub>(T)a</sub>	DRY MATTER FRACTION OF HARVESTED PRODUCTS (DRY)	NUTRITION POTENTIAL IN CROP RESIDUES
	kg N (kg d.m.) <sup>-1</sup>				kg d.m. (kg d.m.) <sup>-1</sup>		kg N/ha
Wheat	0.006	0.009	1.510	0.520	0.230	0.890	-
Rye	0.005	0.011	1.090	0.880	0.220	0.880	-
Barley	0.007	0.014	0.980	0.590	0.220	0.890	-
Oat	0.007	0.008	0.910	0.890	0.250	0.890	-
Maize	0.006	0.007	1.030	0.610	0.220	0.870	-
Potato	0.019	0.014	0.100	1.060	0.200	0.220	-
Sugar beet	-	-	-	-	-	-	20
Oil plants	0.008	0.008	1.130	0.850	0.190	0.910	-
Tobacco	0.015	0.012	0.300	0.000	0.540	0.900	-
Maize for silage	0.015	0.007	0.000	0.000	0.540	0.900	-
Meadows	0.015	0.012	0.300	0.000	0.800	0.900	-
Peas	0.008	0.008	1.130	0.850	0.190	0.910	-
Lens	0.008	0.008	1.130	0.850	0.190	0.910	-
Beans	0.008	0.008	1.130	0.850	0.190	0.910	-
Other leguminous plants	0.027	0.022	0.300	0.000	0.400	0.900	-
Soya	0.008	0.008	0.930	1.350	0.190	0.910	-
Clover	0.025	0.016	0.300	0.000	0.800	0.900	-
Alfaalfa	0.027	0.019	0.290	0.000	0.400	0.900	-

**Country specific FRAC<sub>Renew</sub>:** Equation 11.6 (IPCC 2019 RF) requires use the fractions of the total area of crops, that is renewed annually. For annual crops, FraC<sub>Renew</sub> equals to 1 and FraC<sub>Renew</sub> equals to 0.2. These assumptions are for the forage/pasture five-years renewal frequency. The perennial forage such as alfalfa and clover grows in 4 and 3 rotations. The topic was discussed with experts from the National Agricultural and Food Centre – The Research Institute of Grassland and Mountain Farming. Information published in the article - *Growing and Utilization of Grassland and Clover grassland on Arable Land of Foothill and Mountain Areas* (in Slovak) by Mariana Jančová assumed clover rotation in 3-years cycle and alfalfa rotation in 4-years cycle. Clover and alfalfa are grown in monocultures for seed growing purpose. In addition, FraC<sub>Renew</sub> equal to 0.2 was assumed for the forage/pasture renewal, assuming five-year renewal frequency. These values were based on expert judgment.

**Country specific FRAC<sub>Remove</sub>:** Slovak inventory uses a N-flow approach to calculate the emissions from 3.B and 3.D, which is in line with the IPCC Guidelines, the N<sub>2</sub>O emissions from straw used for bedding is reported in CRF 3.D.a.2 Animal manure applied to soils, and this amount of N was taken into account in the value of FraC<sub>Remove</sub>. The value of FraC<sub>Remove</sub> was calculated for all year from the N content of straw used for bedding divided by the sum of the N content of the above-ground biomass of grain crops of which straw is used for bedding (wheat, barley, rye and oats). The amount of straw used as bedding material was taken from Articles: Livestock breeding by Vojtech Brestenský and Storage of agricultural fertilizers by Vojtech Brestenský (in Slovak) and Removal and storage of fertilizers by Vojtech Brestenský (in Slovak). Publications were provided litter requirements per species and categories per day in kilograms. Nitrogen input from straw was not available in presented publications. Nitrogen input from straw was taken from article Nitrogen fertilization after harvesting cereals by Štefan Gáborík (in Slovak). In aforementioned article, average nitrogen inputs from straw in selected cereals (wheat, barley) were estimated as 0.82%. FraC<sub>Remove</sub> parameter for silage maize was implemented while only

below-ground biomass was considered. It is assumed, that maize for silage is used for fodder purpose in Slovakia.

According to the ERT recommendation A.4 from the draft ARR 2022 (sent on 28<sup>th</sup> February 2023), the amount of forage consumed by livestock was removed from below-ground biomass in meadows. Maize for silage is using for biogas production in biogas stations. Based on expert judgement of ERT and country expert judgement  $Frac_{Remove}$  for maize is 1. According to the publication Guidelines for the support for selected non-projects measures, the farmer is obliged to maintain agricultural areas in a condition suitable for grazing or cultivation in accordance with § 5 of the SR Government Regulation no. 342/2014 Coll. Areas of permanent grassland or meadows must be managed in accordance with agro-technical practice. For areas of meadows, this means maintaining all areas by mowing, grazing and additionally by mulching according to altitude and in following terms: Mowing 4 times per year from 0-800 meters above sea level, grazing 4 times per year from 0-800 meters above sea level and mulching as well. Based on presented information, it was impossible to derivate share for  $Frac_{Remove}$ . The review analysis of inventories was done and only in Polish inventory  $Frac_{Remove}$  parameter was derivated. Poland is neighbouring country with similar agricultural conditions and value was taken into Slovak inventory. Used  $Frac_{Remove}$  and  $Frac_{Renew}$  values are presented in [Tables 5.72](#) and [5.73](#).

**Table 5.72: Parameters used to estimate emissions from crop residues**

TYPE OF CROP	$Frac_{Renew}$	$Frac_{Remove}$
WHEAT	1	0.17
RYE	1	0.17
BARLEY	1	0.17
OAT	1	0.17
MAIZE	1	0
POTATO	1	0
SUGAR BEET	1	0
OIL PLANTS	1	0
TOBACCO	1	0
MAIZE FOR SILAGE	1	1
MEADOWS	0.20	0.95
PEAS	1	0
LENS	1	0
BEANS	1	0
OTHER LEGUMINOUS PLANTS	1	0
SOYA	1	0
CLOVER	0.33	0
ALFALFA	0.25	0

**Table 5.73: Nitrogen in bedding materials and  $Frac_{Remove}$  in particular years**

YEAR	N INPUT FROM BEDDING MATERIALS	N CONTENT OF ABOVE-GROUND BIOMASS OF GRAIN CROPS USED AS BEDDING MATERIAL	$Frac_{Remove}$ (WHEAT, BARLEY, RYE, OAT)
	t		%
1990	415.531	6 167.765	6.74
1995	319.525	5 038.292	6.34
2000	309.934	3 677.159	8.43
2005	299.021	2 921.960	10.23
2010	233.091	1 961.914	11.88
2011	190.096	1 657.890	11.47
2012	202.753	1 789.406	11.33

YEAR	N INPUT FROM BEDDING MATERIALS	N CONTENT OF ABOVE-GROUND BIOMASS OF GRAIN CROPS USED AS BEDDING MATERIAL	Frac <sub>Remove</sub> (WHEAT, BARLEY, RYE, OAT)
	t		%
2013	192.604	1 779.498	10.82
2014	226.830	1 780.288	12.74
2015	228.504	1 818.204	12.57
2016	207.206	1 655.661	12.51
2017	239.565	1 742.469	13.75
2018	253.338	1 828.406	13.86
2019	240.214	1 704.719	14.09
2020	221.479	1 487.173	14.89
2021	221.643	1 316.377	16.15
2022	181.063	1 112.213	16.28
2023	197.405	1 162.191	16.99

### Activity data

Activity data on crop yields and annual area of harvested crops were taken from the ŠÚ SR. To estimate the N added to soils from crop residues and forage/pasture renewal, mainly default parameters from Table 11.2, 11.1A (IPCC 2019 RF) were used. Since yield statistics are reported as field-dry weight, a correction factor was applied to estimate dry matter yields in accordance with the Equation 11.7 IPCC 2019 RF):

$$\text{Crop}_{(T)} = \text{Yield Fresh}_{(T)} * \text{DRY}$$

Where: **Crop<sub>(T)</sub>** = harvested dry matter yield for crop T in kg d.m./ha, **Yield Fresh<sub>(T)</sub>** = kg of fresh weight per ha, **DRY** = dry matter fraction of harvested crop T in kg of d.m.

**Table 5.74: Growing areas and total nitrogen in crops and legumes in 2023**

CROP		HARVESTED AREA	HARVESTED ANNUAL CROP YIELD CROP <sub>(T)</sub>	ANNUAL AMOUNT OF N IN CROP RESIDUES
		ha	t d.m. ha <sup>-1</sup>	kg N yr <sup>-1</sup>
CEREALS	Wheat	406 762.14	5 449.46	28 535.46
	Ray	10 428.50	2 972.87	300.04
	Barley	114 186.55	4 700.09	6 393.18
	Oat	9 689.59	1 866.92	166.81
OTHER	Maize	138 448.29	6 947.89	8 952.49
	Potato	5 428.18	5 503.58	148.90
	Sugar beet	22 126.51	0.00	442.53
	Oil plants	272 568.38	2 833.84	9 485.95
	Tobacco	0.00	0.00	0.00
	Maize for silage	63 186.70	28 304.22	6 760.34
NITROGEN FIXING CROPS	Meadows	494 928.65	2 478.35	3 337.60
	Peas	5 386.34	1 931.90	127.80
	Lens	476.7	362.15	2.12
	Beans	0.00	0.00	0.00
	Other leguminous plants	12 142.31	1 688.91	400.71
	Soya	50 152.51	2 355.01	1 225.87
	Clover	10 792.38	5 103.98	718.66
	Alfalfa	46 560.13	5 839.17	2 795.12
2023 TOTAL		<b>1 663 263.22</b>	<b>78 338.14</b>	<b>69 378.87</b>



## 5.12.7. Mineralization or Immobilization Associated with Loss or Gain of Soil Organic Matter (CRF 3.D.1.5)

Emissions are reported in the categories 3.D.1.5 – Mineralization or immobilization associated with loss or gain of soil organic matter for the first time in 2021 submission.

### Methodological issues – method

$F_{SOM}$  refers to the amount of N mineralised from loss in soil organic C in mineral soils through land-use change or management practices. In order to estimate the N mineralised as consequence of this loss of soil carbon, the Equation 11.8 of 2006 IPCC Guidelines was applied:

$$F_{SOM} = \sum_{LU} \left[ \left( \Delta C_{Mineral,LU} * \frac{1}{R} \right) * 1000 \right]$$

$F_{SOM}$  = the net annual amount of N mineralized in mineral soils as a result of loss of soil carbon through change in land use or management, kg N,  $\Delta C_{Mineral, LU}$  = average annual loss of soil carbon for each land-use type (LU), tonnes,  $R$  = C:N ratio of the soil organic matter.  $LU$  = land-use and/or management system type

The  $N_2O$  estimation from mineralization and immobilization of nitrogen is based on default emission factors according to table 11.1 of the 2019 IPCC RF (0.005 kg  $N_2O$ -N/kg N). A default value of 15 for the C:N ratio (R) was applied according to the p.11.20 IPCC 2019 RF. Used activity data is consistent with the LULUCF sector category 4(III) – Direct  $N_2O$  emissions from N mineralization/immobilization.

### Activity data

The activity data was taken from the carbon loss from management changes under 4.B.1 - Cropland Remaining Cropland/mineral soils. These carbon losses calculated in the LULUCF sector based on the detailed land-use matrices were used as activity data to calculate the N-losses due to mineralization.

**Table 5.75: Activity data and emissions in the category 3.D.1.5 in 1990 – 2023**

YEAR	N IN MINERAL SOILS THAT IS MINERALIZED/IMMOBILIZED IN ASSOCIATION WITH LOSS OF SOIL C	3.D.1.5 - MINERALIZATION/IMMOBILIZATION ASSOCIATED WITH LOSS/GAIN OF SOIL ORGANIC MATTER
	kg/year	Gg
1990	30 760.0	0.0002
1995	102 840.0	0.0008
2000	157 710.0	0.0012
2005	206 870.0	0.0016
2010	208 010.0	0.0016
2011	202 690.0	0.0016
2012	188 440.0	0.0015
2013	176 690.0	0.0014
2014	168 230.0	0.0013
2015	167 030.0	0.0013
2016	160 670.0	0.0013
2017	158 000.0	0.0012
2018	143 750.0	0.0011
2019	121 420.0	0.0010
2020	116 130.0	0.0009
2021	92 440.0	0.0007
2022	83 590.0	0.0007
2023	82 990.0	0.0007

### 5.12.8. Cultivation of Organic Soils (CRF 3.D.1.6)

The area of histosols is very limited in the Slovak Republic. The area of histosols in agricultural area was 450 ha in 2023 and is constant in time series. Emissions from this source are below the threshold of significance for all years as documented in [Table 5.76](#). Therefore, notation key 'NE' is reported for the N<sub>2</sub>O emissions in CRF Table 3.D. Used activity data is consistent with the LULUCF sector.

**Table 5.76: Activity data, emission factors and emissions from histosols in particular years**

YEAR	AREA	EFs	N <sub>2</sub> O EMISSIONS
	ha	kg N <sub>2</sub> O-N/ha <sup>1</sup>	Gg
1990	450	0.029	0.00565714
1995	450	0.029	0.00565714
2000	450	0.029	0.00565714
2005	450	0.029	0.00565714
2010	450	0.029	0.00565714
2015	450	0.029	0.00565714
2016	450	0.029	0.00565714
2017	450	0.029	0.00565714
2018	450	0.029	0.00565714
2019	450	0.029	0.00565714
2020	450	0.029	0.00565714
2021	450	0.029	0.00565714
2022	450	0.029	0.00565714
2023	450	0.029	0.00565714

### 5.12.9. Atmospheric Deposition (CRF 3.D.2.1)

This part of N<sub>2</sub>O emissions resulted from the processes of atmospheric deposition of ammonia and NO<sub>x</sub>, as well as due to the transformation of nitrogen from leaching and runoff losses. Because of the decrease in direct nitrogen input to the soil, the indirect emissions decreased during the evaluated period, too. Total indirect emissions from atmospheric deposition were 0.345 Gg in 2023, which were -49% lower compared to 1990 and 3% lower compared to previous year.

#### Methodological issues – method

Tier 1 approach and default emission factor were used for estimation of indirect N<sub>2</sub>O emissions from atmospheric deposition. This category is estimated in conjunction with the category 3.B - Manure Management. Emissions were estimated following this equation:

$$N_2O_{(ATD)} = [(F_{SN} * Frac_{GASF}) + ((F_{ON} + F_{PRP}) * Frac_{GASM})] * EF_4 * \frac{44}{28}$$

Where: **N<sub>2</sub>O<sub>(ATD)</sub>** = annual amounts of N<sub>2</sub>O emissions from atmospheric deposition of N volatilised from managed soils in kg, **F<sub>SN</sub>** = annual N amount of synthetic fertilisers applied to soils in regions in kg, **F<sub>ON</sub>** = annual amount of managed animal manure and sewage sludge applied to soils in kg N, **F<sub>PRP</sub>** = annual amount of urine and dung N deposited by grazing animals in kg, **Frac<sub>GASF</sub>** = fraction of synthetic fertiliser N that volatilised as NH<sub>3</sub> and NO<sub>x</sub> kg volatilised in kg of N applied (added), **Frac<sub>GASM</sub>** = fraction of applied organic N fertilizer and urine & dung deposited by grazing animals in kg N volatilised as NH<sub>3</sub> and NO<sub>x</sub>, **EF<sub>4</sub>** = emission factor for N<sub>2</sub>O emissions from atmospheric deposition in kg N-N<sub>2</sub>O on soils and water surfaces (kg NH<sub>3</sub>-N + NO<sub>x</sub>-N volatilised)

The mean value for leaching of nitrogen varies in the range of 7-10 kg/ha/ year (7% of N-inputs) in national conditions (Bielek, 1998). The IPCC default emission factor (0.010 kg N<sub>2</sub>O-N/kg N) was used in time-series. It is assumed, that 10% of nitrogen input from synthetic fertilizers applied on soil volatilizes (NH<sub>3</sub> and NO<sub>x</sub>) and 20% of nitrogen from manure applied on soil volatilizes.

## Activity data

Activity data in this category is consistent with the activity data in the categories 3.D.1.1 – Synthetic Fertilizers and 3.D.1.2 – Animal Manure Applied to Soil. **Table 5.77** shows time series of activity data, emission factors and N<sub>2</sub>O emissions in this category.

**Table 5.77: Input parameters, EFs and N<sub>2</sub>O emissions in 3.D.2.1 - Atmospheric Deposition in particular years**

YEAR	TOTAL VOLATILIZED N	EFs	N <sub>2</sub> O EMISSIONS
	<i>t</i>	<i>kg N<sub>2</sub>O-N/kg N</i>	<i>Gg</i>
1990	43 190.24	0.01	0.679
1995	21 805.14	0.01	0.343
2000	21 710.54	0.01	0.341
2005	22 276.78	0.01	0.350
2010	22 150.08	0.01	0.348
2011	23 472.24	0.01	0.369
2012	21 597.64	0.01	0.339
2013	23 262.40	0.01	0.366
2014	24 407.80	0.01	0.384
2015	23 254.92	0.01	0.365
2016	24 688.74	0.01	0.388
2017	24 248.25	0.01	0.381
2018	25 243.42	0.01	0.397
2019	25 101.74	0.01	0.394
2020	24 370.52	0.01	0.383
2021	24 254.24	0.01	0.381
2022	22 728.56	0.01	0.357
2023	21 965.59	0.01	0.345

### 5.12.10. Nitrogen Leaching and Run-off (CRF 3.D.2.2)

Total losses in soils were 1.5% of nitrogen input due to leaching, runoff, and erosion in the Slovak Republic, which is country specific value. Country specific methodology for estimation of  $Frac_{Leach-National}$  was implemented into the inventory during 2022 submission according to continual improvement of emission estimation. In 2021, used methodology was published in the international publication *Atmosphere*<sup>7</sup>.

Total indirect emissions from nitrogen leaching and run-off were 0.797 Gg, which is more than 86% less than 1990 value and -71% compared to previous year. After 2005, the value of  $Frac_{Leach}(national)$  has a dynamic character due to the unstable trend of the wet area - alternation of very dry (low  $Frac_{Leach}(national)$ ) and very humid years (high  $Frac_{Leach}(national)$ ), which may be caused by changing climatic conditions in Slovakia over the last 30 years - floods (2010) and drought (2015,2022). Please see **Table 5.63**. For more information on national study, please see reference 9 with the link to the scientific article in *Atmosphere* (p. 311).

<sup>7</sup> [Estimation of N<sub>2</sub>O emissions from the agricultural soils and determination of nitrogen leakages](#). *Atmosphere. Land-Atmosphere Interactions: Biogeophysical and Biogeochemical Feedbacks*, 2020, Zv. 11

## Methodological issues – method

Tier 2 method and default emission factor were used for the estimation of indirect N<sub>2</sub>O emissions from nitrogen leaching and run-off. This category is estimated in conjunction with category 3.B.2. Emissions were estimated following the equation:

$$N_2O_{(L)} = (F_{SN} + F_{ON} + F_{PRP} + F_{CR} + F_{SOM}) * \text{Frac}_{LEACH-(H)} * EF_5 * \frac{44}{28}$$

Where: **N<sub>2</sub>O<sub>(L)</sub>** = annual amount of N<sub>2</sub>O emissions produced from leaching and run-off of N additions to managed soils in kg, **F<sub>SN</sub>** = annual amount of synthetic fertilizer N applied to soils in kg N, **F<sub>ON</sub>** = annual amount of managed animal manure, compost, sewage sludge and other organic N additions applied to soils, where leaching and run-off occurs in kg N, **F<sub>SOM</sub>** = 0, **F<sub>PRP</sub>** = annual amount of urine and dung N deposited by grazing animals where leaching and run-off occurs in kg N, **F<sub>CR</sub>** = amount of N in crop residues including N-fixing crops here leaching a run-off occurs in kg N, **Frac<sub>LEACH-(H)</sub>** = fraction of all N added in managed soils, where leaching run-off occurs, that is through leaching and run-off in kg of N additions, **EF<sub>5</sub>** = emission factor for N<sub>2</sub>O emissions from N leaching and run-off in kg N<sub>2</sub>O-N (kg N leached and run-off)

Default emission factor (0.011 kg N<sub>2</sub>O-N/kg N) was used for time series.

According to *Mosier et al*, the suggested value of **Frac<sub>LEACH</sub>** is 30%. Value is recommended for calculation of N<sub>2</sub>O emission through leaching in the 2019 IPCC GL where it is defined that for the areas with active irrigation and areas where the total precipitation is for a short time higher than evaporation, the value 30% of the proportion of nitrogen leached out of the utilized agricultural land (**Frac<sub>LEACH</sub>**) is used. For dryland regions, where precipitation and irrigation are lower than evapotranspiration throughout most of the year, leaching is unlikely to occur, **Frac<sub>LEACH</sub>** is equal to zero.

Inclusion of irrigated areas and humid areas modify the default nitrogen leached from arable land and grassland **Frac<sub>LEACH</sub>** to the country specific value according to the equation:

$$\text{Frac}_{LEACHNATIONAL} = (\text{Frac}_{IRR} + \text{Frac}_{WET}) * \text{Frac}_{LEACH}$$

Where: **Frac<sub>IRR</sub>** = the proportion of irrigated areas to the total agricultural land area, **Frac<sub>WET</sub>** = share of the humid area to the total area of arable land and grassland in %, **Frac<sub>LEACHNATIONAL</sub>** = the national value of the proportion of the leached nitrogen from the cultivated soil in %.

## Analysis of Irrigated Areas in Slovakia

The share of irrigated areas in Slovakia was derived from the official statistics published by the Hydromelioration, the state enterprise. Area for particular years 1990 – 2002 was not available, therefore, the data gap was modelled using linear extrapolation tool in Excel. Obtained data were compared with the EUROSTAT datasets. Identified data gaps and inconsistencies are shown in [Table 5.78](#). The total of the utilized agricultural area was taken from the official statistics of the Statistical Office of the Slovak Republic. For the correct determination of the proportion of irrigated areas, it was important to distinguish the type of irrigation. In the case of drip irrigation, water is gradually soaked into the soil, and no nitrogen leaching occurs. Therefore, drip irrigation areas were excluded from the analysis. From the statistics it is visible, that the proportion of irrigated areas in Slovakia is decreasing due to the obsolescence of the irrigation network, i. e. decrease by 86.4% in 2021 compared to 1990. Statistical data about irrigated areas could not be fully verified because only Hydromelioration publishes this type of data in its annual reports. The Statistical Office of the Slovak Republic did not publish such data type and EUROSTAT published only an incomplete data on proportion of irrigated area (proportions are available for 2006, 2008, 2011 and 2014).

In 2023, total irrigated area in Slovakia was 55 603 hectares, representing only 4.2% of agricultural land. The proportion of irrigated areas to the total utilized agricultural areas is listed in [Table 5.78](#).


**Table 5.78:** The proportion of irrigated areas to the total utilized agricultural areas

YEAR	TOTAL IRRIGATED AREAS	UTILIZED AGRICULTURAL AREAS	SHARE OF IRRIGATED AREAS TO THE TOTAL AREAS OF AGRICULTURAL USE FRAC <sub>IRR</sub>	SHARE OF IRRIGATED AREAS ACCORDING TO EUROSTAT
	ha		%	
1990	406 138	1 473 453	27.6%	
1995	348 888	1 487 714	23.5%	
2000	291 638	1 507 178	19.3%	
2001	280 188	1 502 051	18.7%	
2005	147 519	1 504 147	9.8 %	
2010	206 523	1 501 997	13.7 %	
2011	194 215	1 500 905	12.9 %	0.8 %
2012	187 574	1 499 568	12.5 %	
2013	168 277	1 498 986	11.2 %	
2014	154 698	1 498 119	10.3 %	1.3 %
2015	62 239	1 495 789	4.2 %	
2016	60 818	1 494 900	4.1 %	
2017	54 421	1 494 566	3.6 %	
2018	56 408	1 406 399	4.0%	
2019	54 952	1 348 919	4.1%	
2020	23 441	1 346 047	1.7%	-
2021	55 393	1 347 023	4.2%	-
2022	25 887	1 403 864	1.8%	-
2023	55 360	1 307 119	4.2%	-

### Estimation of humid areas in Slovakia

Climatic parameters, evapotranspiration and precipitation (*Figure 5.28*) were used to estimate humid areas in Slovakia. Detailed data were obtained from 41 professional meteorological stations operated by the SHMÚ. Data were analysed and aggregated to monthly and annual averages for purposes of the analysis.

**Figure 5.28:** Network of meteorological stations in Slovakia



The evaporation in agricultural areas occurs mainly through evapotranspiration ( $ET_0$ ) and depends on meteorological conditions, soil characteristics, farming practices and crop types. It means that evapotranspiration can vary within the country or in time and cannot be expressed by one single representative value. For purposes of this study, we assumed the appearance of vegetation during the

whole year, therefore we replaced evaporation. Evapotranspiration was estimated by SHMÚ experts for all 41 meteorological stations with the Penman-Monteith combined method. The equation uses standard climatological data of solar radiation (sunshine), air temperature, humidity and wind speed. The weather parameters' measurements should be made at 2 m (or converted to that height) above an extensive surface of green grass, completely shading the ground and with adequate humidity.

A climatic indicator of humidification is a climatological index used for regionalization of the climate in terms of humidification. It represents the relationship between the amount of water, which is possible to evaporate from the surface of sufficiently humidified soil and vegetation. The climatic indicator of humidification is calculated by the relationship:

$$\sum(P) + \sum(ET_0) > K$$

Where:  $ET_0$  = the sum of potential evapotranspiration,  $P$  = the precipitation total,  $K$  = the humidification of soils.

The rainy season has to be identified for the estimation of humid areas. The rainy season is defined as the period when precipitation is higher than evapotranspiration. Parameter of humidification of the soil is higher than 1, the equation adjusts to:

$$\frac{P}{ET_0} > 1$$


According to the definition of the  $Frac_{LEACH}$  in the 2006 IPCC Guidelines, the determination of 'rainy seasons' is based on precipitation and Pan Evaporation ( $E_{PAN}$ ) data. Rainy seasons are defined as periods when rainfall  $> 0.5 \cdot Pan$  Evaporation, then  $P/E_{PAN} > 0.5$ , where  $P$  is the monthly precipitation. In the case of this study, we use evapotranspiration  $\sum P/\sum ET_0 \geq 1$ . The share  $P/ET_0$  was analysed for 41 meteorological stations.

To cover the whole area of Slovakia, the presented meteorological data were interpolated. The interpolation was processed in the Geographic Information System (QGIS software) using the Inverse Distance Weighting Interpolation function. Interpolation parameters distance coefficient 2, number of columns 3 000 and number of rows 1 500 were applied. In the raster image ([Figure 5.29](#)), areas with a  $\sum P/\sum ET_0 \geq 1$  were extracted by using the contours function and used to trim the underlying layers by available geoprocessing tools. The highly accurate database called The Land Parcel Identification System (LPIS) was used as the underlying layer. Based on geoprocessing analysis, arable land data was revealed. Data on evapotranspiration and precipitation were available in 2023, therefore geoprocessing analysis was performed. Based on [Figure 5.29](#), decrease of humid areas will continue in 2023.

In 2022, the total humid area was 61 292,1 ha, which is 3% of the total agricultural area registered in LPIS ( $Frac_{WET}$ ). The total irrigated area ( $Frac_{IRR}$ ) in Slovakia 25 887 hectares, representing only 1.8% of agricultural land. To calculate the specific national value for nitrogen losses from agricultural land due to leaching ( $Frac_{LEACHNATIONAL}$ ) we used equation:

$$Frac_{LEACHNATIONAL} = (Frac_{irr} + Frac_{wet}) * Frac_{LEACH}$$

**Figure 5.29:** Grassland and arable land where  $\sum P/\sum ET \geq 1$  for 2005, 2010, 2015, 2021 and 2022



### Activity data

Activity data in this category is consistent with activity data in categories 3.D.1.1 - Synthetic Fertilizers and 3.D.1.2 – Animal Manure Applied to Soil. **Table 5.79** shows the time series of parameters, EFs and N<sub>2</sub>O emissions.

**Table 5.79:** Input parameters, EFs and N<sub>2</sub>O emissions in 3.D.2.2 - Nitrogen Leaching and Run-off in particular years

YEAR	TOTAL LOSS OF N	EFs	N <sub>2</sub> O EMISSIONS	THE FRACTION OF N INPUT TO MANAGED SOILS THAT IS LOST THROUGH LEACHING AND RUN-OFF
	<i>t</i>	<i>kg N<sub>2</sub>O-N/kg N</i>	<i>Gg</i>	<i>%</i>
1990	94 177.54	0.0110	1.628	24%
1995	44 335.98	0.0110	0.766	22%
2000	34 728.14	0.0110	0.600	19%
2005	49 549.73	0.0110	0.857	23%
2010	73 446.67	0.0110	1.270	37%
2011	12 617.09	0.0110	0.218	6%
2012	12 918.08	0.0110	0.223	6%
2013	32 196.59	0.0110	0.557	15%
2014	47 386.83	0.0110	0.819	19%
2015	6 145.46	0.0110	0.106	3%

YEAR	TOTAL LOSS OF N	EFs	N <sub>2</sub> O EMISSIONS	THE FRACTION OF N INPUT TO MANAGED SOILS THAT IS LOST THROUGH LEACHING AND RUN-OFF
	<i>t</i>	<i>kg N<sub>2</sub>O-N/kg N</i>	<i>Gg</i>	<i>%</i>
2016	40 666.08	0.0110	0.703	16%
2017	21 722.04	0.0110	0.375	9%
2018	5 800.28	0.0110	0.100	2%
2019	21 812.99	0.0110	0.377	9%
2020	30 671.60	0.0110	0.530	12%
2021	13 316.96	0.0110	0.230	5%
2022	3 358.90	0.0110	0.058	2%
2023	46 124.46	0.0110	0.797	20%

## 5.13. Prescribed Burning of Savannas (CRF 3.E)

The category 3.E Prescribed Burning of Savannas does not occur in the Slovak Republic. Therefore, notation key 'NO' is reported for CRF 3.E category.

## 5.14. Field Burning of Agricultural Residues (CRF 3.F)

This form of cultivation is strictly prohibited by the law in the Slovak Republic. No emissions from this category were estimated. Therefore, notation key 'NO' is reported for CRF 3.F category.

## 5.15. Liming (CRF 3.G)

The soil acidity causes deficient of calcium and magnesium in soils. The presence of the cations of hydrogen and aluminium in the sorption complex causes adverse effects for the growth of the root system of plants. The result is a decrease in the volume of soil and lack of water and nutrients for crops from the soils. The purpose of liming is a correction of soil acidity to normal value with limestone application.

### 5.15.1. Limestone Application (3.G.1)

#### Methodological issues – method

Emissions were calculated according to tier 1 method (IPCC 2006 GL). Due to missing geographical data on limestone consumption, Tier 2 approach is still not implemented. The CO<sub>2</sub> emissions from liming were calculated according to the equation:

$$\text{CO}_2 \text{ emissions} = M * \text{EF} * \frac{44}{12}$$

Where: **CO<sub>2</sub> emissions** = emissions from application of besides limestone and other materials, **M** = annual amount of limestone in tonnes, **EF** = default, a **carbon conversion factor (44/12)** = coefficient for conversion CO<sub>2</sub>-C to CO<sub>2</sub>

The default conversion factor (EF) used for limestone (CaCO<sub>3</sub>) is 0.12.



**Table 5.80: Activity data, EFs and estimated CO<sub>2</sub> emissions in 3.G.1 – Limestone Application in particular years**

YEAR	TOTAL AMOUNT OF CaCO <sub>3</sub>	CARBON CONVERSION FACTOR	CO <sub>2</sub> EMISSIONS
	t		Gg
1990	99 514.70	0.12	43.786
1995	82 398.20	0.12	36.255
2000	72 805.93	0.12	32.035
2005	20 086.88	0.12	8.838
2010	17 533.07	0.12	7.715
2011	32 130.27	0.12	14.137
2012	23 977.70	0.12	10.550
2013	25 362.49	0.12	11.159
2014	25 425.12	0.12	11.187
2015	26 321.44	0.12	11.581
2016	11 287.59	0.12	4.967
2017	4 471.34	0.12	1.967
2018	7 572.04	0.12	3.332
2019	8 248.01	0.12	3.629
2020	14 206.24	0.12	6.251
2021	8 944.16	0.12	3.935
2022	2 017.17	0.12	0.888
2023	21 017.00	0.12	9.25

### Activity data

The consumption of limestone increased in 2023 compared to 2022 by 1042% due to increase in consumption compared to the previous year (2022) and decrease compare to base year almost 79%. This was caused by the cancellation of subsidies for the purchase of limestone by agricultural enterprises and an increase in the purchase prices of dolomite and limestone.

Data on liming of agricultural soils (cropland) are provided by the ÚKSUP. For the years 1998 – 2023, activity data are based on summarization of records that were submitted by landowners/users to the ÚKSUP according to the Act No 136/2000 Coll. on fertilizers as amended by Act No 555/2004 Coll. For the years 1990 – 1998, only estimated values are available. Data was extrapolated with linear extrapolation tool in Excel sheet. Data contain only limestone or fertilizers containing limestone, which is a difference compared to previous submission. Other calcareous substances containing only Ca and CaO were subtracted from activity data.

## 5.15.2. Dolomite Consumption (CRF 3.G.2)

### Methodological issues – method

The CO<sub>2</sub> emissions from liming of dolomite were calculated according to the equation:

$$\text{CO}_2 \text{ emissions} = M * \text{EF} * \frac{44}{12},$$

Where: **CO<sub>2</sub> emissions** = emissions from application of besides components containing dolomite, **M** = annual amount of limestone in tonnes, **EF** = default, a **carbon conversion factor (44/12)** = coefficient for conversion CO<sub>2</sub>-C to CO<sub>2</sub>

The default conversion factor (EF) used for limestone (MgCO<sub>3</sub>) is 0.13.

### Activity data

The data on consumption of dolomite was provided by the UKSÚP. Consumption of dolomite increased in 2023 compared to 2022 by 114%. For the years 1998 – 2023, data are based on the summarization

of records that were submitted by landowners/users to the ÚKSUP according to the Act No 136/2000 Coll. on fertilizers as amended by Act No 555/2004 Coll. Data contain applied MgCO<sub>3</sub> substances put on soil annually. The total MgCO<sub>3</sub> amount was calculated. For the years 1990 – 1998, only estimated values are available. Data was extrapolated with linear extrapolation tool in Excel sheet. Data contain only dolomite or fertilizers containing dolomite, which is a difference compared to previous submission. Other dolomite substances containing only Mg and MgO were subtracted from activity data.

**Table 5.81: Activity data, EFs and estimated CO<sub>2</sub> emissions in 3. G.2 - Dolomite Application in particular years**

YEAR	TOTAL AMOUNT OF MgCO <sub>3</sub>	CARBON CONVERSION FACTOR	CO <sub>2</sub> EMISSIONS
	t		Gg
1990	4 076.22	0.13	1.943
1995	3 668.34	0.13	1.749
2000	4 840.07	0.13	2.307
2005	921.77	0.13	0.439
2010	1 083.43	0.13	0.516
2011	2 107.91	0.13	1.005
2012	1 579.02	0.13	0.753
2013	1 659.65	0.13	0.791
2014	1 625.76	0.13	0.775
2015	1 744.18	0.13	0.831
2016	3 791.40	0.13	1.807
2017	1 365.99	0.13	0.651
2018	1 844.65	0.13	0.879
2019	2 268.64	0.13	1.081
2020	4 614.54	0.13	2.200
2021	4 198.29	0.13	2.001
2022	6 994.77	0.13	3.334
2023	14 990.20	0.13	7.150

## 5.16. Urea Application (CRF 3.H)

In conditions of Slovakia, urea as fertilizer is applied mainly on medium heavy and heavy soils and less on light sandy soils because of its high solubility and possible loss of nitrogen without its uptake by plants. The urea is neither applied on very acid soils. Urea is not the primary source of nitrogen.

### 5.16.1. Methodological Issues – Method

Tier 1 method according to the Equation 11.13 (IPCC 2006 GL) was used for emissions estimation in this category. Default conversion factor (EF) used for urea is 0.20. Estimated emissions are shown in **Table 5.82**. CO<sub>2</sub> emissions from urea application were calculated as follows:

$$\text{CO}_2 \text{ emissions} = M_{\text{CO(NH}_2)_2} * \text{EF} * \frac{44}{12}$$

Where: **CO<sub>2</sub> emissions** = emissions from application of urea in tonnes of CO<sub>2</sub>, **M<sub>CO(NH<sub>2</sub>)<sub>2</sub></sub>** = annual amount of urea fertilizers in tonnes, **EF** = default, a **urea conversion factor (44/12)** = coefficient for conversion CO<sub>2</sub>-C to CO<sub>2</sub>

**Table 5.82: Activity data, EFs and estimated CO<sub>2</sub> emissions in 3.H - Urea Application in particular years**

YEAR	TOTAL AMOUNT OF UREA	UREA CONVERSION FACTOR	CO <sub>2</sub> EMISSIONS
	t		Gg
1990	20 846.74	0.20	15.288
1995	20 846.74	0.20	15.288
2000	16 500.69	0.20	12.101
2005	27 699.02	0.20	20.313
2010	42 189.25	0.20	30.939
2011	54 146.88	0.20	39.708
2012	61 934.09	0.20	45.418
2013	70 899.73	0.20	51.993
2014	79 009.80	0.20	57.941
2015	83 072.60	0.20	60.920
2016	86 006.26	0.20	63.071
2017	86 636.61	0.20	63.534
2018	89 953.97	0.20	65.966
2019	86 644.29	0.20	63.539
2020	86 817.95	0.20	63.666
2021	86 772.93	0.20	63.633
2022	77 207,45	0.20	56.619
2023	76 456,73	0.20	56.068

### 5.16.2. Activity Data

The ÚKSUP provides data on urea application on agricultural soils (cropland). For the years 1998 – 2022, the data was based on the summarization of recordings that had to be submitted by landowners/users to the ÚKSUP according to the national legislation. For the years 1990 – 1997, the data have been estimated as the average of three years' period (1998 – 2000). In the past, the three years' period of urea application was fluctuating with low, medium and higher doses.

## 5.17. Other Carbon – Containing Fertilizers (CRF 3.I)

This category is not estimated in the current submission. The category will be completed in future submissions. The used notation key is NO.

---

<b>CHAPTER 6. LULUCF (CRT 4)</b> .....	<b>309</b>
6.1. Overview of the LULUCF Sector .....	309
6.2. Category-specific QA/QC and Verification Process.....	314
6.3. Category-specific Recalculations.....	315
6.4. Category-specific Improvements and Implementation of Recommendations .....	317
6.5. Time-series Consistency and Uncertainties .....	317
6.6. Forest Land (CRT 4.A) .....	319
6.7. Cropland (CRT 4.B) .....	334
6.7.1. Grassland (CRT 4.C) .....	341
6.8. Wetlands (CRT 4.D) .....	345
6.9. Settlements (CRT 4.E).....	345
6.10. Other Land (CRT 4.F) .....	348
6.11. Direct and indirect nitrous oxide (N <sub>2</sub> O) emissions from nitrogen (N) inputs to managed soils (CRT 4(I)).....	351
6.12. Emissions and removals from drainage and rewetting and other management of organic and mineral soils (CRT 4(II)) .....	351
6.13. Direct and indirect nitrous oxide (N <sub>2</sub> O) emissions from nitrogen (N) mineralization/immobilization associated with loss/gain of soil organic matter resulting from change of land use or management of mineral soils (CRT 4(III)) .....	351
6.14. Biomass Burning (CRT 4(IV)) .....	353
6.15. Harvested Wood Products (HWP) (CRT4.Gs1-2).....	353
<b>Annex A6.1. Land-Use Matrix</b> .....	<b>356</b>
<b>Annex A6.2. Uncertainty Analyses in the LULUCF Sector</b> .....	<b>376</b>

## CHAPTER 6. LULUCF (CRT 4)

This chapter was prepared using GWP<sub>100</sub> taken from the 5<sup>th</sup> Assessment Report of the IPCC by the sectoral experts and institutions involved in the National Inventory System of the Slovak Republic:

INSTITUTE	CHAPTER	SECTORAL EXPERT
National Forest Centre – Forest Research Institute (NFC-FRI)	Chapter 6.1 – 6.6 Chapter 6.9 – 6.17 Annex A6.1 Annex A6.2	Tibor Priwitzer Ivan Barka Pavel Pavlenda
National Agriculture and Food Centre - Soil Science and Conservation Research Institute (NAFC-SSCRI)	Chapter 6.7	Michal Sviček
National Agriculture and Food Centre - Grassland and Mountain Agriculture Research Institute (NAFC-GMARI)	Chapter 6.8	Štefan Pollák

### 6.1. Overview of the LULUCF Sector


The Forestry and Land Use sector covers the wide range of biological and technical processes within the landscape, which are reflected in the GHG inventory. This sector includes all GHGs (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) and basic pollutants from forest fires (NO<sub>x</sub> and CO). Individual inventory of LULUCF categories are linked with all relevant processes related to all five carbon pools (living biomass – above and below ground, dead organic matter – dead wood and litter, soil carbon), as have been defined in the Marrakech Accords. In addition, wood products referred to as harvested wood products (HWP) are reported as an additional pool under LULUCF (CRT sector 4.G).

The inventory in LULUCF sector is based on the definition of representative types of land use categories – Forest Land (FL), Cropland (CL), Grassland (GL), Wetlands (W), Settlements (S) and Other Land (OL). In addition, their temporal changes are reported. The first three categories have the highest importance due to their relative coverage of Slovakia, representing more than 90% of the whole territory. The processes linked to the land use and land-use change are mostly related to CO<sub>2</sub> balance.

Biomass burning, which represents managed processes (i.e. burning of harvest residues) and unmanaged processes (i.e. forest fires), is a special category in the landscape. This category covers all three main GHGs and basic pollutants. The inventory covers also the estimation of CO<sub>2</sub> emissions from the agricultural lime application.

The LULUCF sector with net removals -7 483.72 Gg of CO<sub>2</sub> eq. in 2023 is very important sector and comprises several key categories. [Table 6.1](#) shows summary of total emissions according to the categories. Time series of emissions and removals are illustrated on [Figure 6.1](#) and summarised in [Table 6.2](#). This document uses the GWP 100 based on the IPCC Fifth Assessment Report for the year 2023.

**Figure 6.1: Emissions and removals (Gg of CO<sub>2</sub> eq.) according to the categories in 1990 – 2023**



**Table 6.1: Summary of total emissions and removals according to the categories in 2023**

Category	Net CO <sub>2</sub>		CH <sub>4</sub>	N <sub>2</sub> O	NO <sub>x</sub>	CO
	Emissions/Removals (Gg)		Emissions (kt)			
<b>4. LULUCF</b>	<b>NO</b>	<b>-7 525.91</b>	<b>0.54</b>	<b>0.10</b>	<b>0.35</b>	<b>12.31</b>
A. Forest Land	NO	-7 009.50	0.54	0.03	0.35	12.31
B. Cropland	NO	-654.30	NA, NO	0.03	NO	NO
C. Grassland	NO	-27.85	NO	0.00	NO	NO
D. Wetlands	NO	NO	NO	NO	NO	NO
E. Settlements	76.94	NO	NO	0.02	NO	NO
F. Other Land	88.78	NO	NO	0.02	NO	NO

**Table 6.2: Summary of GHG emissions and removals according to the categories in particular years**

YEAR	Forest land	Cropland	Grassland	Settlements	Other land	LULUCF (CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O)		
	Net CO <sub>2</sub> in Gg					Gg		
1990	-8 262.31	-484.43	-195.77	96.59	293.10	<b>-9 023.23</b>	0.44	0.45
1995	-8 598.32	-371.05	-257.78	61.11	104.09	<b>-9 120.72</b>	0.31	0.31
2000	-7 502.03	-443.24	-309.97	54.04	106.21	<b>-9 015.06</b>	1.08	0.23
2005	-1 876.28	-515.22	-200.82	62.40	186.46	<b>-4 339.92</b>	1.05	0.17
2010	-2 842.11	-559.71	-215.52	102.08	90.36	<b>-4 759.49</b>	0.80	0.12
2011	-3 370.79	-565.49	-274.67	70.27	81.26	<b>-5 128.28</b>	0.96	0.13
2012	-5 098.28	-643.36	-216.49	82.43	116.98	<b>-6 267.12</b>	1.84	0.18
2013	-6 032.18	-502.96	-203.83	97.13	96.93	<b>-6 985.34</b>	0.61	0.12
2014	-3 593.97	-503.93	-182.37	81.38	110.26	<b>-4 817.02</b>	0.90	0.14
2015	-3 940.37	-502.44	-190.85	85.70	185.73	<b>-5 302.92</b>	1.02	0.15
2016	-3 720.88	-590.73	-178.50	80.44	99.35	<b>-5 373.97</b>	0.84	0.14
2017	-3 601.61	-622.22	-164.81	100.99	95.42	<b>-5 269.27</b>	0.93	0.15
2018	-2 893.26	-627.83	-110.83	81.56	143.96	<b>-4 295.58</b>	0.92	0.15
2019	-3 833.66	-633.22	-118.02	84.23	80.62	<b>-5 064.97</b>	1.08	0.15
2020	-6 499.74	-578.04	-92.82	79.30	94.80	<b>-7 243.78</b>	0.98	0.14
2021	-6 330.47	-654.02	-55.23	86.36	72.29	<b>-7 263.27</b>	0.71	0.12
2022	-6 643.65	-649.74	-36.24	80.39	76.37	<b>-7 316.72</b>	1.64	0.17
2023	-7 009.50	-654.30	-27.85	76.94	88.78	<b>-7 525.91</b>	0.54	0.10

GHG Inventory submission 2025 of Slovakia reports carbon stock changes, as well as greenhouse gas emissions and removals from Forest Land (CRT 4.A), Cropland (CRT 4.B), Grassland (CRT 4.C), Settlements (CRT 4.E), Other Land (CRT 4.F) and Harvested Wood Products (CRT 4.G). In the category

4.A - FL, carbon stock change in living biomass, dead organic matter and mineral soils is reported. In the 4.B - CL, carbon stock change in living biomass is reported. The carbon stock changes in living biomass, dead organic matter and mineral soils are reported for CL, GL, S and OL converted from the FL. Direct N<sub>2</sub>O emissions from N fertilization of Forest Land and Others (CRT 4(I)) as well as non-CO<sub>2</sub> emissions from drainage of soils and wetlands (CRT 4(II)) are not reported. N<sub>2</sub>O emissions (direct and indirect) from N mineralization associated with conversion to Cropland, Grassland, Settlements and Other land are reported (CRT 4(III)). Emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from the Biomass Burning are reported in CRT Table 4(IV). Summary of all categories is described in [Table 6.3](#).

**Table 6.3: Reported emissions, methodological tiers and emission factors (EF) in LULUCF in 2023**

CATEGORY	CO <sub>2</sub>		CH <sub>4</sub>		N <sub>2</sub> O		
	method applied	EF	method applied	EF	method applied	EF	
4.A	FOREST LAND						
4.A.1	Forest Land Remaining Forest Land	T1,T2	CS,D				
4.A.1-4(IV)	Biomass Burning	T1,T2	CS,D	T2	CS,D	T2	CS,D
4.A.2	Land Converted to Forest Land	T1, T2	CS, D	T2	CS, D	T2	CS, D
4.A.2.1	Cropland Converted to Forest Land	T1, T2	CS				
4.A.2.2	Grassland Converted to Forest Land	T1, T2	CS				
4.A.2.5	Other Land Converted to Forest Land	T1, T2	CS				
4.A.2-4(IV)	Biomass Burning	T2	CS, D	T2	CS, D	T2	CS, D
4.B	CROPLAND						
4.B.1	Cropland remaining Cropland	T1, T2	CS, D				
4.B.2	Land Converted to Cropland	T1, T2	CS, D			T2	CS, D
4.B.2.1	Forest Land Converted to Cropland	T1, T2	CS, D				
4.B.2.2	Grassland Converted to Cropland	T1, T2	CS, D				
4.B.2.5	Other Land Converted to Cropland	T1, T2	CS, D				
4.B.2-4(III)	Direct and indirect N <sub>2</sub> O Emissions from N Mineralization/ Immobilization					T1, T2	CS, D
4.C	GRASSLAND						
4.C.1	Grassland remaining Grassland	T1					
4.C.2	Land Converted to Grassland	T1, T2	CS, D			T2	CS, D
4.C.2.1	Forestland Converted to Grassland	T1, T2	CS, D				
4.C.2.2	Cropland Converted to Grassland	T1, T2	CS, D				
4.C.2.5	Other Land Converted to Grassland	T1, T2	CS, D				
4.C.2-4(III)	Direct and indirect N <sub>2</sub> O Emissions from N Mineralization/Immobilization					T1, T2	CS, D
4.E	SETTLEMENTS						
4.E.2	Land Converted to Settlements	T1, T2	CS, D			T2	CS, D
4.E.2.1	Forest Land Converted to Settlements	T1, T2	CS, D				
4.E.2.2	Cropland Converted to Settlements	T1, T2	CS, D				
4.E.2.3	Grassland Converted to Settlements	T1, T2	CS, D				
4.E.2-4(III)	Direct and indirect N <sub>2</sub> O Emissions from N Mineralization/Immobilization					T1, T2	CS, D
4.F	OTHER LAND						
4.F.2	Land Converted to Other Land	T2	CS, D				
4.F.2.1	Forest Land Converted to Other Land	T2	CS, D				
4.F.2.2	Cropland Converted to Other Land	T2	CS, D				
4.F.2.3	Grassland Converted to Other Land	T2	CS, D				
4.F - 4(III)	Direct and indirect N <sub>2</sub> O Emissions from N Mineralization/ Immobilization					T1, T2	CS, D
4.G	HARVESTED WOOD PRODUCTS						
4.G	Harvested Wood Products	T2	CS, D				

---

The area of Forest Land in the Slovak Republic covers 41.4% of the territory and wood harvesting is historically an important economic activity. The LULUCF sector represents a sink of GHG since 1990. Historically stable trend was disrupted due to high wood extraction from damaged forests in 2005 after strong windstorm from the end of 2004, which consequently resulted in the decrease of total sinks to the half of previous volumes.

Slovakia provides further explanation of the climate domain and ecological zones of Slovakia. The entire territory of Slovak lies in the climatic reference region of Western and Central Europe according to IPCC climatic reference. According to the IPCC 2006 GL (Vol. 4, Chap. 3, Annex 3.A.5, "Default climate and soil classifications"), the Slovakian territory belongs to IPCC climate domain: "Cool Temperate Moist" and ecological zone: "Temperate continental forest".

The identification of the LULUCF categories is based on the data from the Geodesy, Cartography and Cadastre Authority of the Slovak Republic (GCCA), which represents a key data source for identification of spatial extent of individual categories. The GCCA annually issues the Statistical Yearbook of the Soil Resources in the Slovak Republic. It provides updated cadastral information of the LULUCF areas. Since 2007, this book is available on the website of the GCCA. The [GCCA database](#) distinguishes ten land categories, six of them belonging to the land utilized by agriculture (arable land, hop-fields, vineyards, gardens, orchards, grasslands) and the rest of them under other use (forest, water surfaces, built-up areas and courtyards, and other land). Six land-use categories have been selected – Forest Land, Cropland, Grassland, Wetlands, Settlements and Other Land as given in the IPCC 2006 GL, Volume 4, Agriculture, Forestry and Other Land Use. The Slovak Republic used the following LULUCF definitions for reporting of GHG emissions and removals in the categories:

**Forest Land** - this category includes the land covered by all tree species serving for the fulfilment of forest functions and the land on which the forest stands were temporarily removed with aim of their regeneration or establishment of forest nurseries or forest seed plantation. In the Permanent Forest Inventory and the Statistical Office databases, it is referred to as timberland.

**Cropland** - this category includes lands for growing cereals, root-crops, industrial crops, vegetables and other kinds of agricultural crops. Perennial woody crops are also included in this category. There are included lands temporarily overgrown with grass or used for growing of fodder lasting several years, as well as hotbeds and greenhouses if they are built up on the arable land. This category also includes fallow land, which is arable land left for regeneration for one growing season. During this period there were not sown specific crops or just crops for green manure, eventually it is covered by spontaneous vegetation, which would be ploughed in.

**Grassland** - this category includes permanent grasslands and meadows used for the pasture or hay production, which is not considered as cropland.

**Wetlands** - this category include artificial reservoirs and dam lakes, natural lakes, rivers and swamps.

**Settlements** - this category include all developed land, including transportation infrastructure and human settlements of any size.

**Other Land** - this category represented by bare soil, rock and all unmanaged land areas that do not fall into any of the other categories. Each of these categories is divided into land remaining in the given category during the inventory year, and land converted into the category from another one. The areas of six LULUCF categories remaining in the specific category are in [Table 6.4](#).

The increasing trend of FL is evident in the Slovak Republic since 1970. The opposite, decreasing trend of Cropland was recorded at the same time. Grassland areas decreased from 1980 to the 1990 and since this year increasing trend was recorded up to 2005. Since 2005, moderately downward trend has been taking place. Settlements category has continuously increasing trend during the whole period. This situation is mostly caused by the development of transport infrastructure, industrial areas, municipal development and raising the standards and infrastructure in the country and is very often connected with decreasing of the Cropland and Other Land area.




Slovakia provides an explanation for the cause of the abrupt increase in the areas of settlements and decrease in other land occurring around 1995. The abrupt changes in Settlements and Other land occurring around 1995 was likely due to new property owners rushing to get their land recognized as 'settlement' during the country's transition to a market economy. Slovakia assumes that the increase of area in the category of Settlements and reciprocal decline of area in the category of Other Land could be caused by administrative transfer of Other Land to Settlements. The reason for this could be a new territorial administrative division of Slovakia (from 3 to 8 regions) and the effort of the new administrators to obtain property in the form of settlements. This idea results from consultation with the provider of cadastral data (Geodesy, Cartography and Cadastre Authority of the Slovak Republic - GCCA). The abrupt increase in the areas of settlements between 2015 and 2016 in the 4.E.1 LU category was caused by the implementation of the rule about the 20 years long transition period. According to IPCC 2019 GL under the default assumption in every inventory year, the area converted to a land-use category should be added to the category "land converted to" and the same area removed from the land remaining in the land-use category. The area of land that entered that "land converted to" category, 21 years ago (if using the default 20-year period), should be removed and added to the category "land remaining land". Wetland represents 1.9% (94 kha) of the Slovak territory and it is considered constant, not involving any land-use conversions.

**Table 6.4:** The area of LU categories remaining in category in particular years

YEAR	4.A.1	4.B.1	4.C.1	4.E.1	4.F.1
	<i>kha/year</i>				
1990	1 809.15	1 492.15	685.50	94.69	190.37
1995	1 861.77	1 502.19	740.79	102.63	203.45
2000	1 929.76	1 517.42	766.82	109.57	128.14
2005	1 945.13	1 513.92	762.47	116.75	128.01
2010	1 981.89	1 511.70	766.40	116.85	130.80
2011	1 983.77	1 510.36	766.97	117.40	130.65
2012	1 985.11	1 508.36	786.60	117.59	131.46
2013	1 985.74	1 507.23	787.84	117.18	131.36
2014	1 986.15	1 505.97	785.35	117.37	131.13
2015	1 986.73	1 503.58	784.51	117.90	130.04
2016	1 988.25	1 502.40	786.01	184.44	129.49
2017	1 991.52	1 501.95	788.93	206.45	129.33
2018	1 993.56	1 502.51	791.68	206.34	129.57
2019	1 995.57	1 501.94	800.48	206.65	130.00
2020	1 996.76	1 503.21	811.98	207.60	130.41
2021	1 997.86	1 505.04	823.40	210.73	131.12
2022	1 998.59	1 505.25	831.95	211.61	131.18
2023	2 000.17	1 506.27	835.47	212.87	130.89


The land-use matrix shown in [Table A6.1](#) and on [Figure 6.2](#) represents the areas of land-use change among the major categories from 1990 to 2023 for individual years. The annual totals for individual years in the matrix do not correspond to the areas referred to in CRT Tables. These areas account for the progressing for 20 years' transition period beginning in the year 1970. This approach represents tier 1 approach of the IPCC 2006 GL for calculation of soil carbon stocks changes. The areas of biomass carbon pools are not the same as for the soil carbon ones.

**Figure 6.2:** Overall development trends in area of categories from 1970 – 2023 (based on information from the GCCA of the Slovak Republic)



Land-use matrix identifying annual conversions among the categories for the period 1990 – 2023 and describing initial and final areas of particular categories are listed in the [Annex A6.1. \(Table A6.1.1\)](#). The distribution of the LULUCF categories in Slovakia in 2023 is shown on [Figure 6.3](#). Forest Land represents the major category, accounting for 41.4% of the total area, followed by the Cropland with 31.1%, Grassland with 17.3%, Settlements with 4.9%, Other Land with 3.4% and Wetlands with 1.9% of the total country area.

**Figure 6.3:** Distribution of the LULUCF categories in Slovakia in 2023



## 6.2. Category-specific QA/QC and Verification Process

QA/QC procedures in the LULUCF sector are linked with the QA/QC plans for the National Inventory System at the sectoral level and followed basic rules of QA/QC as defined in the IPCC 2006 GL.

The calculation is based on annually submitted or published input data of several institutions:

- 
- the Geodetic and Cartographic Institute Bratislava;
  - the Geodesy, Cartography and Cadastre Authority of the Slovak Republic (GCCA);
  - the Statistical Office of the Slovak Republic (ŠÚ SR);
  - the National Forest Centre - Institute for Forest Resources and Information (NFC-IFRI);
  - the National Forest Centre - Forest Management Planning Institute (NFC-FMPI);
  - the Central Controlling and Testing Institute in Agriculture (ÚKSUP);
  - or information published by the research organizations: Research Institute of Geodesy and Cartography in Bratislava, National Forest Centre - Forest Research Institute (NFC-FRI), National Agriculture and Food Centre - Soil Science and Conservation Research Institute (NAFC - SSCRI) and National Agriculture and Food Centre – Grassland and Mountain Agriculture Research Institute (NAFC-GMARI).

Each of the institutions has internal quality rules depending on the main tasks of the institution. Published data on carbon content in litter, soil and biomass at national level are based on results of laboratories that follow quality management standards in laboratory praxis and successfully participate in the ring tests (international inter-laboratory comparisons).

The primary input data (values, units) are checked for the plausibility and conformity (time series). When possible, data is checked with data from other sources. Data submitted by responsible institution upon request are compared with the relevant published information. The remarkable changes or trend differences in input data are directly discussed and checked with responsible persons and data provider. The input data sets and sources are archived by sectoral expert.

In the process of the emissions calculation and estimation, all procedures are checked (correctness of equations, interim results, units, trend evaluation). Results (output data) are checked according to the QC procedures. Comparison with data in time series and space (results from other countries) are important steps in the data check. Parameters and emission factors used for NID are compared with results and factors in other countries or regions that can be comparable (similar bio-geo-region, site conditions, ways and intensity of land management, etc.).

Methods and emission factors used in the emissions inventory are internally consulted or/and reviewed among experts in the NFC that are not involved in the national emission inventory implementation.

The QC checks (e.g. consistency check between CRT data and national statistics) were done during the CRT and NID compilation by sectoral experts, General QC questionnaire was filled out and archived by the QA/QC manager. The QA is conducted by another LULUCF expert from the NFC and by independent expert from the Ministry of Environment of the Slovak Republic and the Ministry of Agriculture and Rural Development of the Slovak Republic.

### 6.3. Category-specific Recalculations


Recalculations and reallocations made in the LULUCF sector were provided and implemented in line with the Improvement and Prioritization Plan reflecting recommendations made during previous reviews and expert improvement. Short description of recalculations implemented in 2025 submission is in [Table 6.5](#).

**Table 6.5:** Description of recalculations implemented in 2025 submission

NUMBER/ RECOMMENDATION	CATEGORY	DESCRIPTION	REFERENCE
1	4. LULUCF	Changed in activity data in 4.III and HWP categories	Chapter 6
2	4.III Direct & indirect N <sub>2</sub> O emissions from N mineralization/immobilization	Calculation error (incorrect formula used) of indirect N <sub>2</sub> O emissions from N mineralization/immobilization.	Chapters 6.7, 6.8, 6.9, 6.10
3	4.G HWP	Correction of input activity data (Wood base panels Production - years 2021, 2022)	Chapter 6.17


The 4.III Direct & indirect N<sub>2</sub>O emissions from N mineralization/immobilization and Harvested wood products categories within the LULUCF sector were recalculated in 2025 submission. Recalculated values for the whole sector differ from the submission in 2024 by -0.05% to 0.02% in particular years (**Figure 6.4**), the net CO<sub>2</sub> eq. removals decreased by -0.01% in average.

**Figure 6.4:** Comparison of CO<sub>2</sub> eq. (Gg) in the 2024 and 2025 submissions for LULUCF sector



The recalculation was realised also in HWP category in the years 2021 and 2022. The main reason was correction of input activity data – wood base panel and paper and paper board. Recalculated values for the HWP differ from the submission in 2024 of 0.13% in year 2021 and 0.35% in 2022 (**Figure 6.5**), the net CO<sub>2</sub> eq. removals increased by 0.01% in average. These changes improved accuracy of the calculations.

**Figure 6.5:** Comparison of CO<sub>2</sub> eq. (Gg) in the 2024 and 2025 submissions for HWP



## 6.4. Category-specific Improvements and Implementation of Recommendations

All the recommendation from the latest UNFCCC review 2022 were implemented in previous submission, except of recommendation L.1, where the research is still ongoing. ERT recommendation (L.1 - draft ARR 2022) was partially implemented. Continuation of the technical research in order to provide reliable data for estimating CSC in living biomass, dead organic matter and soil organic matter is the long-term process and the results will be implemented in the next submissions. Slovakia clarified that the calculation of CSC in deadwood carbon pools in land converted to forest land, based on partial results from the above-mentioned research, was included in the CRT tables and the NID.

## 6.5. Time-series Consistency and Uncertainties

The time series are consistent in the area of using consistent methodology, consistent way of collection of activity data and use of consistent emission factors and other parameters. Disturbances and fluctuations in time series and in emissions or removals are described in the particular chapters and can be reasonably explained by national circumstances. Three recalculations ([Table 6.6](#)) was performed in this submission.

The uncertainty analysis of the LULUCF sector was performed by the Approach 1 methods using the Equations 3.1 and 3.2 (Volume 1, Chapter 3, IPCC 2006 GL). Used parameters in the Approach 1 uncertainty analyses within the LULUCF sector according to the categories are referred to in [Table 6.6](#). More and detailed information is in the SVK NIR 2018, the [Chapter 6.5 \(Annex A6.2\)](#).

**Table 6.6: Uncertainties of activity data and EFs in individual C pools and LULUCF categories**

LULUCF CATEGORY		ACTIVITY DATA	EMISSION FACTOR	EF REFERENCES
4.A.1	Forest Land remaining Forest Land - living biomass	3%	82.84%	IPCC 2006 GL
4.A.2	Land converted to Forest Land - living biomass	3%	40.61%	IPCC 2006 GL
4.A.2	Land converted to Forest Land – DOM (litter)	3%	75.00%	expert judgement
4.A.2	Land converted to Forest Land - mineral soils	3%	75.00%	expert judgement
4.B.1	Cropland remaining Cropland - living biomass	3%	75.00%	IPCC 2006 GL
4.B.1	Cropland remaining Cropland – mineral soils	3%	76.09%	expert judgement
4.B.2	Land converted to Cropland - living biomass	3%	107.98%	IPCC 2006 GL (tab. 5.1, 6.4), Šmelko et al. 2003
4.B.2	Land converted to Cropland – DOM (DW/litter)	3%	75.24%	SVK NFI, expert judgement
4.B.2	Land converted to Cropland - mineral soils	3%	75.00%	expert judgement
4.C.1	Grassland remaining Grassland - living biomass	3%	75.00%	IPCC 2006 GL
4.C.1	Grassland remaining Grassland – mineral soils	3%	76.09%	expert judgement
4.C.2	Land converted to Grassland - living biomass	3%	107.98%	IPCC 2006 GL (tab. 5.1, 6.4), Šmelko et al. 2003
4.C.2	Land converted to Grassland – DOM (DW/litter)	3%	75.24%	SVK NFI, expert judgement
4.C.2	Land converted to Grassland - mineral soils	3%	75.00%	expert judgement

LULUCF CATEGORY		ACTIVITY DATA	EMISSION FACTOR	EF REFERENCES
4.E.2	Land converted to Settlements - living biomass	3%	107.98%	IPCC 2006 GL (tab. 5.9, 6.4), Šmelko <i>et al.</i> 2003
4.E.2	Land converted to Settlements – DOM (DW/litter)	3%	75.24%	SVK NFI, expert judgement
4.E.2	Land converted to Settlements - mineral soils	3%	75.00%	expert judgement
4.F.2	Land converted to Other Land - living biomass	3%	107.98%	IPCC 2006 GL (tab. 5.9, 6.4), Šmelko <i>et al.</i> 2003
4.F.2	Land converted to Other Land – DOM (DW/litter)	3%	75.24%	SVK NFI, expert judgement
4.F.2	Land converted to Other Land - mineral soils	3%	75.00%	expert judgement
4.G	Harvested Wood Products	5%	50.00%	IPCC 2006 GL

In a reflection to the ERT recommendations made in previous reviews, the NS SR has started preparation work on improvement of uncertainty analyses of the key categories inside the LULUCF sector. In October 2017, the Expert Working Group for LULUCF (EWG LULUCF) was created. The EWG LULUCF consists of the LULUCF sectoral experts, uncertainty expert, expert for emission modelling, QA/QC expert and NS SR coordinator. Independent observers are experts for LULUCF legislation from the Ministry of Environment of the Slovak Republic and Ministry of Agriculture and Rural Development of the Slovak Republic. Main task of the EWG LULUCF is the preparation of higher tier uncertainty analyses and further improvement in this sector. The first meeting of the EWG LULUCF agreed on the Working Plan for the next period of approximately three years.

Working Plan (in shortened version):

- Preparation of detailed key category analysis on level and trend assessment in the LULUCF sector using Approach 1 (IPCC 2006 GL);
- Analysis of key categories by trend and level assessment, incorporating formulas and parameters, including comments on availability of national data on uncertainty, literature;
- Uncertainty expert checks information sent by sectoral experts and set up the range of work and other possibility;
- Cooperation with the Cadastral Office;
- Evaluation of input data;
- Preparation of Monte Carlo model;
- Evaluation of results;
- Further improvements.

During the years 2018 – 2022, work on the improvement of uncertainty analyses for the LULUCF categories was ongoing according to the agreed schedule. Several expert meetings were followed by discussions and email communication. During the first part of work done in 2017, key categories were identified as follow:

- Approach 1 – level assessment (CO<sub>2</sub>): FL remaining FL, L converted to FL, CL remaining CL and HWP;
- Approach 2 – level assessment (CO<sub>2</sub>): FL remaining FL, L converted to FL, CL remaining CL, L converted to GL, L converted to S, L converted to OL and HWP;
- Approach 1 & 2– trend assessment (CO<sub>2</sub>): FL remaining FL, L converted to FL, CL remaining CL, L converted to CL, L converted to GL and HWP;

- 
- From non-CO<sub>2</sub> gases, only N<sub>2</sub>O emissions from L converted to CL is a key category in level and trend assessment.

According to the key category identification, work on the Monte Carlo simulation started in the second half of 2018 and it has continued during 2019, 2020, 2021 and 2022. Preliminary results of the application of Monte Carlo simulations are provided in the [Annex A6.2](#) of this Document. Work will continue following the available capacities and sources.

Analyses of uncertainties using Monte Carlo simulations for the main LULUCF categories (including the HWP), as well as for the whole LULUCF sector, were included in this submission.

## 6.6. Forest Land (CRT 4.A)

Forests currently cover 41.4% of the Slovak Republic. The area of forests in Slovakia is in temperate-zone and is managed. Forests in Slovakia are known for richly diverse species composition mainly with European beech being the dominant forest tree species covering 35.4% of the area, followed by Norway spruce (21.1%), oaks (10.3%) and pine (6.4%). Broadleaved species represent 64.8% of all tree species found in Slovak forests. Percentage of coniferous species (currently at 35.2%) has been steadily decreasing since 1980; since 2000, their presence fell by 7.0%. Due to harmful agents in forests, Norway spruce percentage has fallen from the original 26.8% in 2000 to current 22.1%, a drop by 4.3%. At the same time, the area of European beech has increased by 5.1% whilst the area of noble hardwoods (maples and ash) has grown by 2.4% (Green Report, 2024). In addition to the overall representation of individual tree species, the mixing of tree species in particular forest management units is also an important indicator of species diversity and forest stand stability. At present, the most represented types of forest stands are: beech forests (27.5%), conifer-beech mixtures (25.5%), spruce forests (15.0%) and forests dominated by oak (9.0%). The actual age structure of forest significantly differs from the normal (ideal/optimal) structure. At present, forests 70+ years old are the most represented group of forests. Majority of these forests reached the age when it is desirable to start with their regeneration. Conversely, percentage of young forests (20-70 years old) is below normal. In the last ten years or so, the proportion of the youngest forest stands of the 1<sup>st</sup> and 2<sup>nd</sup> age classes have increased significantly. This is due to the high extent of forest damage caused by harmful agents and subsequent regeneration of damaged forests (Green Report, 2024). At present, forest management is focused more on close-to-nature silvicultural procedures, establishment of forest stands with better structural, species diversity, and higher ecological stability. Split by main species groups reads as follows: coniferous forests 31%, broadleaved forests 50%, and mixed forests 19%.

The growing stock has shown a continual increase in the volume of available timber in forests. The estimated growing stock was 487.1 mil. m<sup>3</sup> (merchantable volume, defined as tree stem and branch volume under bark with minimum diameter threshold of 7 cm) in 2023, and increase of 4.3 mil m<sup>3</sup> compared to 2022. Currently, due to the present age-structure of forests in Slovakia, the growing stock of forests is the highest. However, their volume is already at the culmination point. It is expected that in the coming years and decades these stocks will decrease due to a gradual change in age structure. This trend is also confirmed by the observed decrease in the average annual increase in growing stocks in forests in the SR, which was as follows: 1991 – 1995: +5.9 mil. m<sup>3</sup> annually, 1995 – 2000: +6.4 mil. m<sup>3</sup>, 2000 – 2005: +5.8 mil. m<sup>3</sup>, 2006 – 2010: +4.6 mil. m<sup>3</sup>, 2011 – 2015: +3.2 mil. m<sup>3</sup>; 2016 – 2020: +1.3 mil. m<sup>3</sup>, and the average annual increase in growing stocks was only 0.8 mil. m<sup>3</sup> in 2021 – 2023. A similar trend to the annual change in total growing stocks can be observed also in the development of the annual change in average growing stock per 1 ha. Average hectare growing stock was 250 m<sup>3</sup> in 2023 (Green Report, 2024).

In 2023, the volume of current annual increment (CAI) reached 11.88 mil. m<sup>3</sup>, or 6.16 m<sup>3</sup> per ha of FL. Over the last few decades, CAI gradually grew to 12.126 mil. m<sup>3</sup> (6.25 m<sup>3</sup> per ha) in 2012. However, since 2012 it has decreased by 1%, or 119 000 m<sup>3</sup>, respectively.

---

Healthy and resilient forests are also an important part of the landscape due to their significant contribution to carbon sequestration. They directly contribute to reduction of greenhouse gas emissions, carbon dioxide in particular, as carbon is stored for a long time in forest biomass, soil and wood products. Along with the increase in growing stock in forests and FL, there is also an increase in carbon stock bound in individual balance categories.

According to [Green Report 2024](#), the carbon stock in forests found in living biomass (aboveground and underground), dead organic mass (deadwood, litter) and forest soils reached a volume of 511.0 mil. t in 2023, with the largest amount stored in soils (270.5 mil. t) and aboveground tree biomass (167.0 mil. t). As a result of the current trend in the development of the age structure of forests, a decrease in the amount of carbon stocks in individual balance categories will occur simultaneously with the decrease in the wood stocks in the forests. (Green Report, 2024).

The total volume of harvested timber reached 7.22 mil. m<sup>3</sup> in 2023. Compared to 2022, realized felling decreased by 0.464 mil. m<sup>3</sup>, and it was lower by 2.3 mil. m<sup>3</sup>, as the planned felling calculated using actual harvesting possibilities and forest regeneration on urgency. Of the total volume, 53.3% of harvested timber represents the coniferous wood and 46.7% broadleaved wood. Of the total timber volume, 3.51 mil. m<sup>3</sup> (48.6%) was felled due to natural disturbances and pests, of which 85.0% was coniferous wood. Despite this, the actual felling is still below the level of total current increment (the volume of timber that accrues in forests every year) and has been even lower than planned felling since 2012, except for the year 2014. The realized logging was lower than CAI during the completely reporting period ([Figure 6.8](#)). Planned and actual felling in volume of m<sup>3</sup> are increasing in Slovakia, despite the fact that in 2020 the volume of felling was the lowest in the last 15 years. The main reason behind increased felling volumes is the current age structure of forests with a high proportion of 70+ years old forests. Due to a high percentage of mature forests approaching rotation, the volume of planned felling kept increasing to reach 9.8 million m<sup>3</sup> in 2020, which was 84.9% more than in 2000. Both the growing stock and the area of mature forests have stagnated in recent years, which indicates the onset of a gradual reduction of previously high felling volumes (Green Report, 2024).

All available information about the forests in Slovakia comes from two sources. The first one is the Forest Management Plan (FMP), updated on a regular basis. Investigation is carried out in a 10-years period – i.e. one tenth of the territory is surveyed each year, practically all forest stands are surveyed once in every 10 years. The survey produces detailed maps, as well as description of the forest stands (e.g. species composition, diameter at breast height, mean height, stock volume, number of trees, basal area, crown closure, volume increment etc.). Gathered data are stored in databases and further processed into aggregated files used for reporting and compilation of various documents including the Compendium of Forestry Statistics - the Aggregated Forest Management Plan (AFMP), and the Permanent Forest Inventory (PFI). Professionally and technically competent non-state experts and companies elaborated Forest Management Plans (FMPs). The FMPs are prepared according to the existing legislation, procedures and methodologies. All relations concerning the FMPs can be found in the Act No 326/2005 Coll. on Forests and Public Notice of the Ministry of Agriculture No 453/2006 Coll. on Forest Management Planning and Forest Protection. The FMPs are approved by provincial (governmental) forest authorities and are audited by the National Forest Centre (NFC). The FMPs have been performed for all forests, owners or users within the Slovak territory (Act No 326/2005 Coll.). For the forest management it is mandatory, that activities, including harvest and harvested volume, are recorded and reported yearly to the state authority.


The second source of information are data from the National Forest Inventory and Monitoring (NFIM). The first cycle of the statistical forest inventory (sample based, tree level) was performed during 2005 – 2006 and the second one during 2015 – 2016 by the NFC. The NFIM is a selective statistical method of forest condition inventory. It has two levels – national and regional, and provides data for all forests regardless of land category (forest, non-forest). The NFIM provided a comprehensive set of data on forests relevant to December 31, 2005. Accuracy and reliability of provided outcomes meets the quality




expected at the beginning of investigation (standard error 2.1% for total standing volume). This data source is not usable for emissions reporting of Forest Land, because it does not cover reporting period sufficiently. However, it is usable for estimation of carbon pools for example of dead organic matter – dead wood.

The 4.A category includes emissions and removals of CO<sub>2</sub> (Gg) associated with forest. Category is divided into subcategories: 4.A.1 - FL remaining FL and 4.A.2 Land converted to Forest Land (L converted to FL). **Figure 6.6** shows area changing during years and **Figure 6.7** shows map of Forest Land in Slovakia.

**Figure 6.6:** Development of activity data (kha) for the category 4.A - FL in the period 1990 – 2023



**Figure 6.7:** Distribution of FL calculated as a spatial share within individual cadastral units



### 6.6.1. Forest Land Remaining Forest Land (CRT 4.A.1)

Emissions estimation is based on the methodology from the IPCC 2006 GL and 2019 IPCC Refinement and activity data from the PFI processed continuously on annual basis. Results of estimation were obtained by using the IPCC methodology and national data on area of forested land and land converted to the forest during the inventory year 2023. This category includes carbon stock change in following carbon pools: living biomass (above and below ground), dead organic matter (dead wood and litterfall) and organic soil carbon. Carbon stock change is given by the sum of changes in living biomass, dead organic matter and soil. Total area of Forest Land remaining Forest Land represents 2 000.174 kha.

## Methodological issues – methods, activity data, emission factors and parameters

The carbon stock change in living biomass was estimated using a gain-loss method according to the Equation 2.7 of the IPCC 2006 GL. This method is based on separate estimation of increments, removals and its difference. Calculations of carbon stock changes in living biomass as a result of annual biomass increment and annual biomass loss was carried out following the Equations 2.9 - 2.12 of the IPCC 2006 GL. Current annual increment (CAI) expressed as merchantable volume, defined as tree stem and branch volume under bark with a minimum diameter threshold of 7 cm are the key inputs to calculate the carbon increment. The CAI values are calculated by the NFC-IFRI, which is the FMP database administrator for Slovakia. The calculation is performed at the level of the individual stands and species using the available stand parameters, yield data and models. The CAI is determined based on the average stocks in the different age levels for individual tree species as the sum of the average increment in the different age levels, expressed per unit of actual area of tree species occurrence.

$G_{TOTAL}$  is the expansion of current annual increment of aboveground biomass ( $G_W$ ) to include its belowground part, involving multiplication by the ratio of belowground biomass to aboveground biomass (often called the root-to-shoot ratio that applies to increments). The current annual increment (merchantable volume increment -  $I_V$ ) is converted to the annual biomass increment ( $G_{TOTAL}$ ) using the biomass conversion expansion factor ( $BCEF_I$ ) and root-to-shoot ratio ( $R$ ) (Equation 2.10 (A) and (B) of the IPCC 2006 GL) as follows:

- $G_{TOTAL} = G_W * (1 + R)$
- $G_W = I_V * BCEF_I$

The root-to-shoot ratio was differentiated according to Table 4.4 of the IPCC 2006 GL (0.20 for coniferous and 0.24 for other broadleaved species). The input data and factors used in the calculation of the biomass carbon stock increment for different tree species are presented in [Table 6.7](#).

**Table 6.7: Annual biomass increment for individual forest tree species in the Slovak Republic in 2023**

Tree Species	Current annual increment	Biomass conversion/ expansion factor	Average annual above-ground biomass growth	Ratio of below-ground biomass to above-ground biomass	Average annual biomass growth above- and below-ground
	CAI (m <sup>3</sup> /ha/yr)	$BCEF_I$	$G_W$ (t dm/ha/yr)	$R$	$G_{TOTAL}$ (t dm/ha/yr)
Spruce	8.10	0.45	3.63	0.20	4.36
Fir	7.12	0.45	3.18	0.20	3.82
Pine	6.18	0.67	4.14	0.20	4.97
Larch	6.42	0.80	5.16	0.20	6.20
Other conifer	2.56	0.54	1.37	0.20	1.65
Oak	4.31	0.87	3.74	0.24	4.64
Beech	5.89	0.78	4.57	0.24	5.66
Hornbeam	6.05	0.91	5.52	0.24	6.85
Maple	6.18	0.72	4.44	0.24	5.51
Ash	7.57	0.72	5.44	0.24	6.75
Elm	6.18	0.74	4.58	0.24	5.68
Turkey oak	4.19	0.93	3.91	0.24	4.85
Locust	4.63	0.91	4.22	0.24	5.24
Birch	2.93	0.68	2.01	0.24	2.49
Alder	2.35	0.68	1.61	0.24	1.99
Linden	7.17	0.51	3.68	0.24	4.56
Breeding poplars	9.83	0.48	4.67	0.24	5.80
Poplar	3.31	0.42	1.38	0.24	1.71
Willow	2.54	0.71	1.81	0.24	2.25
Other broadleaves	1.86	0.68	1.27	0.24	1.58

According to present knowledge, about 55-90% (depending on tree species) of the total tree biomass can be assumed stored in the stems (Šebík et al., 1989). The density of wood (at dry weight) varies depending on tree species, from 0.40 to 0.80 t d.m./m<sup>3</sup> in the national conditions (Požgaj et al., 1993). The annual biomass increments per hectare and year (resulting from application of annual wood volume increment data and biomass expansion factor) varies from 1.40 to 6.80 t d.m./ha for different tree species.

The BCEF<sub>1</sub> showed in [Table 6.8](#) were calculated as a ratio of CAI expressed as tree volume over bark and CAI expressed as merchantable volume (defined as tree stem and branch volume under bark with a minimum diameter threshold of 7 cm) for spruce, fir, pine, beech, oaks and poplar tree species. This is multiplied by the basic wood density of individual tree species. The values of CAI for individual tree species were based on national growth and yield tables (Halaj and Petráš, 1998) using values of age and “bonita” degree (yield class) calculated by the NFC-IFRI Zvolen annually.

Estimation of annual increase in carbon stocks due to biomass increment in FL remaining FL requires inputs of actual stand area (A), annual increment of total biomass (G<sub>TOTAL</sub>) and carbon fraction of dry matter and was calculated by the Equation 2.9 of the IPCC 2006 GL as followed:

$$\Delta C_{FFG} = \sum (A * G_{TOTAL}) * CF$$

The middle of the range values for the carbon fraction of above-ground biomass in forest (all, broadleaves and conifers) (Table 4.3 of the IPCC 2006 GL) was implemented. The carbon content of 51% for coniferous and 48% for broadleaved wood was used for calculation of carbon gains in living biomass. The annual increase in carbon stock due to biomass increment in the category FL remaining FL represents 4 949.40 kt C in 2023 and is shown in [Table 6.8](#).


**Table 6.8:** Total carbon uptake increment for individual forest tree species in 2023

Tree Species	Area of tree species for FL remain FL	Average annual biomass growth above- and below-ground	Annual increase in biomass due to biomass growth	Carbon fraction of dry matter	Annual increase in biomass carbon stocks due to biomass growth
	(kha)	(t dm/ha)	(kt/dm/yr)	(tC/tdm)	(kt C yr)
Spruce	422.237	4.36	1841.59	0.51	939.21
Fir	79.807	3.82	304.77	0.51	155.43
Pine	128.011	4.97	636.33	0.51	324.53
Larch	52.405	6.20	324.73	0.51	165.61
Other conifer	20.802	1.65	34.27	0.51	17.48
Oak	205.618	4.64	953.17	0.48	457.52
Beech	708.862	5.66	4015.51	0.48	1927.44
Hornbeam	120.410	6.85	824.26	0.48	395.65
Maple	52.805	5.51	290.77	0.48	139.57
Ash	30.003	6.75	202.37	0.48	97.14
Elm	0.600	5.68	3.41	0.48	1.64
Turkey oak	52.005	4.85	252.39	0.48	121.15
Locust	35.403	5.24	185.47	0.48	89.02
Birch	34.803	2.49	86.53	0.48	41.54
Alder	15.201	1.99	30.31	0.48	14.55
Linden	8.801	4.56	40.16	0.48	19.28
Breeding poplars	8.601	5.80	49.86	0.48	23.93
Poplar	8.001	1.71	13.66	0.48	6.56
Willow	2.000	2.25	4.49	0.48	2.16
Other broadleaves	13.201	1.58	20.84	0.48	10.00
<b>TOTAL</b>	<b>1 999.574</b>		<b>10 114.90</b>		<b>4 949.40</b>


The annual decrease in carbon stocks due to biomass loss in FL remaining FL follows Equations 2.12 of the IPCC 2006 GL. Slovakia reports that main/primary source of information for annual harvesting is the harvest statistics. The annual harvest volume (H) is collected in the mandatory reporting of forest managers and elaborated by the NFC-IFRI. It covers managed forests, as the reporting is an integral mandatory part of forest management and covers any annual harvest data including thinning and final cut. Relevant forest companies, forest owners or users are obligated to provide data on forest management activities (harvest, silviculture) to the central forestry database annually (Regulation No 297/2011 Coll. of the Ministry of Agriculture and Rural Development of the Slovak Republic). Annual data on harvest includes biomass harvested in forest in a reported year. Even the stolen timber is notified by owners and is included in the annual harvest each year. All subjects (users, companies) managing forest, which realized or did not realized harvest have the statutory duty (Act No 326/2005 Coll. on Forests) to inform the NFC - IFRI authorities about the amount and type of harvest throughout districts.

The annual amount of total harvest and fuel wood removals is published annually in the Green Reports. The harvesting volumes of coniferous and broadleaved trees, CAI and total harvest during the reporting period 1990 – 2023 in Slovakia are presented on **Figures 6.7** and **6.8**.

**Figure 6.8:** The harvesting volume in forest (coniferous and broadleaved) (mil. m<sup>3</sup> volume >7 cm under bark) in 1990 – 2023



**Figure 6.9:** Current annual increment (CAI) and total annual harvest (mil. m<sup>3</sup>) in 1990 – 2023



The annual carbon loss due to commercial felling was calculated using the Equation 2.12 of the IPCC 2006 GL:

- $L_{\text{fellings}} = H * BCEF_R * (1+R) * CF$

Biomass conversion and expansion factors (BCEF<sub>R</sub>) were developed based on new NFI data. BCEF<sub>R</sub> were developed for Norway spruce (*Picea abies*), Pine (*Pinus sylvestris*), Oak (*Quercus robur*) and Beech (*Fagus sylvatica*). The methodology follows a common procedure described in literature (Lehtonen et al., 2004) and cited in the IPCC 2006 GL. The BCEF is generally defined as:

$$\text{BCEF}_i = W_i / V$$

Where: i indicates a tree biomass component, W<sub>i</sub> (Mg) is the dry biomass of component, V (m<sup>3</sup>) is the tree merchantable volume.

Tree-level data of new NFI were used to construct age-related BCEFs. Only inventory plots that contained a dominant share (at least 50% of the basal area) of any of the four key tree species (beech, oak, pine and spruce) were used for the analysis. This selected database contained over 22 thousand trees. Tree merchantable volume and tree aboveground biomass were calculated using national methodology (Petras and Pajtik, 1991). The aboveground biomass functions were used from the studies (Wutzler et al., 2008 for beech trees, Cienciala et al., 2008 for oak trees, Cienciala et al., 2006 for pine trees and Wirth et al., 2004 for spruce). More complete description of the BCEF<sub>R</sub> calculation was published in the report “[Different Approaches to Carbon Stock Assessment in Slovakia](#)”, Chapter 13.

The values of BCEF<sub>R</sub> were calculated for each year separately considering actual age structure of forests.

During the review 2022 the ERT suggested recommendation ([L.14 - draft ARR 2022](#)) that BCEF<sub>R</sub> coefficients for coniferous species be divided by 0.92 and BCEF<sub>R</sub> coefficients for broadleaved species be divided by 0.9 for adding bark and harvest losses in accordance with the 2006 IPCC GL. Slovakia revised its calculation on annual carbon losses due to commercial felling for the forest land remaining forest land category of LULUCF and resubmitted the LULUCF CRT tables with the revised calculations, which was accepted by the ERT and the resubmission confirmed. The forest land remaining forest land removals decreased from -7 422.68 Gg CO<sub>2</sub> eq. to -6 290.29 Gg CO<sub>2</sub> eq. (15.3%) for 2020 through this resubmission. This revision affected the whole time series (1990 – 2020).

The CF factors used in calculation are described in [Table 6.9](#). The carbon loss due to fuel wood gathering was not estimated separately as this activity is very rare in Slovakia and fuelwood is included in total harvest. The total annual carbon release from forest harvest was 3 287.39 kt C in 2023.

**Table 6.9: Activity data and BCEFR used in calculation of carbon losses in 2023**

Tree species	Annual wood removal - harvest volume	Biomass conversion/expansion factor	Annual wood removal - biomass	Ratio of below-ground biomass to above-ground biomass	Annual wood removal - biomass	Carbon fraction of dry matter	L wood-removals including fuelwood
	H (m <sup>3</sup> /yr)	BCEF <sub>R</sub>	(t dm/yr)	R	(t dm/yr)	CF (tC/tdm)	(ktC/yr)
Spruce	3 236 582	0.683	2 209 241	0.20	2 651 089	0.51	1 352.06
Fir	260 805	0.683	178 022	0.20	213 626	0.51	108.95
Pine	291 782	0.572	166 678	0.20	200 013	0.51	102.01
Larch	55 958	0.572	31 966	0.20	38 359	0.51	19.56
Other conifer	6 995	0.572	3 996	0.20	4 795	0.51	2.45
Oak	603 886	0.923	556 711	0.24	690 322	0.48	331.35
Beech	2 177 590	0.833	1 813 126	0.24	2 248 276	0.48	1 079.17
Hornbeam	186 965	0.833	155 672	0.24	193 034	0.48	92.66
Locust	61 988	0.833	51 613	0.24	64 001	0.48	30.72
Poplar	76 986	0.833	64 100	0.24	79 485	0.48	38.15
Other broad	262 951	0.833	218 940	0.24	271 486	0.48	130.31
<b>TOTAL</b>	<b>7 222 488</b>		<b>5 450 066</b>		<b>6 654 485</b>		<b>3 287.39</b>

---

According to the ERT recommendation ([L.13 - draft ARR 2022](#)), Slovakia clarified that wooded land which is below the thresholds for forest land (tree species covering less than 0.3 ha or with density lower than 20%, woody vegetation which potentially cannot exceed 5 m height) reported as other conifers under the forest land remaining forest land category. According to the FAO - Global Forest Resources Assessment 2020 report, Slovakia considers as other wooded land the Alpine vegetation zone with *Pinus mugo* plantations, which are reported under forest land. CSC of other wooded land in forest land remaining forest land represents 0.29 to 17.96 kt C/y (0.01 to 2.71% of total removals of FL remaining FL category in individual years). Other wooded land represents a net sink for whole reporting period. The area of other conifers (other wooded land) ranges from 18 to 22 kha (1.00 to 1.12%), of the total forest land area in individual years. All data are reported in NID [Table 6.7](#) and [Table 6.8 \(Chapter 6.6.1\)](#). Current annual increment of biomass varied from 1.39 to 2.60 m<sup>3</sup>/ha/y, BCEF<sub>I</sub> and BCF<sub>R</sub> are similar as pine tree species.

The assessment of the net carbon stock change in DOM includes dead wood and litter pools.

The dead wood carbon pool contains dead trees from standing, stumps, coarse lying dead wood and small-sized lying dead wood not included in litter or soil carbon pools. The information on dead wood stocks was obtained from the first and second National Forest Inventory (NFI) realized in 2005/2006 and 2015/2016. Before realization of the NFIs, no reliable data on dead wood (except for standing dead trees) were available in Slovakia. Quantification of dead wood was performed by the methodology where all components were determined in the same volume units (m<sup>3</sup> over bark) in order to enable their aggregation. The volume of standing dead trees was determined from the volume equations of living trees (HSK). In order to determine the stump volume, new regression equations were derived, where the diameter at the top of the cut area D and the stump height H represent input variables. The volume of the lying dead wood with the top diameter of 7 cm was calculated from the measured diameters d1 and d2 (cm) outside bark at both ends and the length of each piece inside the inventory plot (IP) or a sub-plot using the Smalian equation (Šmelko, 2000). The volume of small-sized lying dead wood (having diameter from 1 to 7 cm) was estimated by the original method, where the volume of small-sized lying dead wood (in m<sup>3</sup>) densely arranged in 1 m<sup>2</sup> is calculated from the biometrical model as a function of the middle diameter of small-sized lying dead wood multiplied by the area of IP, estimated coverage of small-sized lying dead wood, and tree species proportion (Šmelko et al., 2008). The conversion of volume to dry biomass was carried out based on wood density coefficients and using reduction coefficients according to the degrees of decomposition of dead wood (Fresh 1; Hard 0.83; Soft 0.66; Decayed 0.5). The volume was multiplied by the wood dry matter density coefficients according to the NIML List of Forest Tree Abbreviations (Šebeň, 2017) and the above-ground biomass in mass units was calculated. The conversion of biomass to carbon was performed by a coefficient of 0.496 (Šmelko et al., 2011). According the NFIs the average C stock of dead wood was calculated on 6.6 ± 0.5 t C/ha for 2005 as well as 7.4 ± 0.7 t C/ha for 2015 in Forest Land category. Using the estimated trend based on these empirical observations, data for the years between these data points were linearly interpolated and extrapolated accordingly beyond that period. The Equation 2.19 of the IPCC 2006 GL was used for calculation of the net C stocks change of DW.

The litter pool definition used in the inventory includes all non-living biomass with a size less than the minimum diameter defined for dead wood (1 cm). The small-sized lying dead wood (diameter between 1 and 7 cm), in various states of decomposition above the mineral soil are not a part of litter, because they are included in dead wood. The litter includes the surface organic layer (L, F, H horizons) as usually defined in soil profile description and classification. Live fine roots above the mineral or organic soil (of less than the minimum diameter limit chosen for below-ground biomass) are included in litter because they cannot be distinguished empirically. All existing national databases on carbon stocks in forest soil organic layer are based on the same approach and soil data were obtained by standard sampling procedure.


The mean carbon stock in forest litter is 8.3 t C/ha. The value is derived from datasets of the Forest Monitoring System (FMS) and the NFI. The changes of forests management that would dramatically change litter properties and litter carbon changes do not occur, i.e. no significant changes of carbon stocks in litter in the 4.A.1 were assumed (tier 1). Information on soil carbon stocks in forest soils is from soil survey on permanent monitoring plots (16x16 km grid of large-scale forest monitoring), soil survey on the NFI plots and sets of research plots databases. The most detailed information source with respect to soil depth (0-10 cm, 10-20 cm, 20-40 cm, 40-80 cm) and sampling design is the set of 112 plots of large-scale monitoring and 9 intensive monitoring plots. The largest and the most representative information source is the set of plots of the NFI (almost 1 500 plots with sampling depth limited to 20 cm). Carbon stocks per hectare (in both data sources) are calculated using information on carbon concentration in fine soil, bulk density and coarse fragment content. The calculated soil carbon stocks range from 13.7 to 486.8 C t/ha (for the depth 0-20 cm in both the FMS and the NFI datasets). Supplementary information about carbon content and carbon stock in forest soil comes also from other research plots with detailed soil profile description and classification. It is used mainly for derivation of indices for recalculation of carbon stocks for different depths and respective soil types or site units.

For estimation of carbon stock change for mineral soils carbon pool, tier 1 approach was used and assumed that soil carbon stocks change in category 4.A.1 is considered to be zero. Evaluation of results from re-sampling after 13 years (in 16x16 km grid of monitoring plots) has been finished. Though slight increase of soil carbon stocks seems to be possible, tests did not show significant differences (changes). Based on these tests, forest soils (for forests remaining forests) are neither carbon emission source nor sink. Soil data management and evaluation of differences after 10 years from the NFI plots (8x4 km grid of inventory plots) is expected to be done in near future.

In the central European conditions, the mineral soils and the litter are not a source of net emissions (Pavlenda, 2016) in managed Forest Land, based on the principles of sustainable forestry. The same assumption was made in countries with similar soils and climatic conditions (Hungary, the Czech Republic).

Figure 6.10 shows that the net CO<sub>2</sub> removals in the FL remaining FL represent -6 680.76 Gg in 2023. It is necessary to mention that every forest in Slovakia is considered as managed. Uptake of carbon into the biomass of forest trees has slightly increased since 1990 and then decreased after 2004, however fluctuations can be observed in time series of harvested volume, especially in the last ten years, which can be attributed to fluctuations of salvage logging after disturbances.

Figure 6.10: Summary results of CO<sub>2</sub> removals (Gg) from FL-FL subcategory in 1990 – 2023



## 6.6.2. Biomass Burning (CRT 4.A.1 - 4(IV))

The biomass burning activity in 4.A.1 - 4(V) includes emissions of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O associated with forest fires and biomass burning on forest areas. The National Forest Centre – Forest Protection Service summarized activity data from controlled burning and forest fires since 1999.

Slovak harvesting system partly includes burning of harvesting residues if decided by forest managers and the risk of fire is limited (at cleared plots after processing of trees infested by bark beetle or after clear cuts). The harvesting residues are burned on about 50% of the forest clearing area. The differences are in the quantity of burning biomass. For coniferous 10% and for broadleaved about 25% of above ground biomass is burned. Because there is no official estimation of amount of post logging slash, the expert judgment was used for calculation. The biomass fraction burned on clearing areas was quantified on the basis of annually reported amount of main felling, separately for coniferous and broadleaved species as well as the BCEF<sub>R</sub> were applied in calculation of harvest losses in FL remaining FL. The emissions from biomass residues burning were calculated according to the Equation 2.27 and the default emission factors provided in Table 2.5 (IPCC 2006 GL). Default combustion factor value for post logging slash burn in other temperate forests is 0.62 according to Table 2.6 (IPCC 2006 GL).

The main information sources on wildfires or forest fires are the internal fires statistics of the Ministry of Interior and the “Reports of the occurrence of harmful agents in Slovakia”. Reported forest fires in Slovakia were at the area of 29.46 ha in 2023. This number decreased compared to the previous year 2022, when the total burnt area was 1210.55 ha. The average burnt forest area per one fire was 0.5 ha. The largest forest area damaged by fire was 14 ha. The forest fires occurred mostly in spring and in the summer. The GHG<sub>s</sub> emissions from wildfires were calculated based on Equation 2.27 (IPCC 2006 GL) and the mass of fuel available for combustion derived using known areas burnt annually. The average stock per hectare (250 m<sup>3</sup>/ha in 2023) and biomass expansion factor was used for estimation. The GHG emissions from wildfires were calculated based on known annual burnt area and the average stock per hectare. **Table 6.10** shows biomass burned in forests with emissions in the same units.

**Table 6.10: Biomass burned in Forest land remaining Forest land category, CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from wildfires and controlled burning in particular years**

YEAR	BIOMASS BURNED (t d.m.)	AREA BURNED (ha)	CO <sub>2</sub> EMISSIONS (kt)*		CH <sub>4</sub> EMISSIONS (t)		N <sub>2</sub> O EMISSIONS (t)	
	Controlled Burning	Wildfires	Controlled Burning	Wildfires	Controlled Burning	Wildfires	Controlled Burning	Wildfires
1990	104 472.85	208.94	IE	45.83	304.43	137.28	16.84	7.59
1995	89 822.48	65.48	IE	15.57	261.74	46.63	14.48	2.58
2000	132 254.40	892.90	IE	231.79	385.39	694.34	21.32	38.41
2005	214 689.72	511.65	IE	141.62	625.61	424.22	34.61	23.47
2010	218 608.12	189.12	IE	54.72	637.02	163.92	35.24	9.07
2011	210 905.06	396.75	IE	115.83	614.58	346.97	34.00	19.19
2012	126 556.09	1 658.91	IE	490.48	368.78	1469.26	20.40	81.28
2013	127 582.17	266.23	IE	79.35	371.77	237.71	20.57	13.15
2014	252 540.53	188.74	IE	56.32	735.90	168.72	40.71	9.33
2015	241 696.53	346.65	IE	103.94	704.30	311.37	38.96	17.22
2016	234 857.16	171.87	IE	51.79	684.37	155.15	37.86	8.58
2017	229 233.90	292.80	IE	88.30	667.99	264.50	36.95	14.63
2018	239 997.14	244.33	IE	73.77	699.35	220.99	38.69	12.22
2019	228 119.75	454.87	IE	138.07	664.74	413.59	36.77	22.88
2020	189 340.40	465.32	IE	141.36	551.74	423.45	30.52	23.42
2021	193 403.60	156.50	IE	47.77	563.58	143.10	31.18	7.92



YEAR	BIOMASS BURNED (t d.m.)	AREA BURNED (ha)	CO <sub>2</sub> EMISSIONS (kt)*		CH <sub>4</sub> EMISSIONS (t)		N <sub>2</sub> O EMISSIONS (t)	
	Controlled Burning	Wildfires	Controlled Burning	Wildfires	Controlled Burning	Wildfires	Controlled Burning	Wildfires
2022	190 943.68	1192.11	IE	360.17	556.41	1078.91	30.78	59.68
2023	176 438.98	29.01	IE	8.87	514.14	26.56	28.44	1.47

\*tier 1 approach, CO<sub>2</sub> emissions from controlled burning are included in the total biomass loss associated with harvesting (CRT Table 4.A).

### Controlled burning

Total methane emissions from controlled burning were 514.14 t and total emissions of N<sub>2</sub>O were 28.44 t in 2023. CO<sub>2</sub> emissions from controlled burning are included in the total biomass loss associated with harvesting in CRT Table 4.A.

### Wildfires

Total methane emissions from wildfires were 26.56 t and total emissions of N<sub>2</sub>O were 1.47 t in 2023. CO<sub>2</sub> emissions were 8.87 Gg in 2023.

### 6.6.3. Land converted to Forest Land (CRT 4.A.2)

This category includes all processes connected with conversion of lands into Forest Land. This activity is closely connected with afforestation or reforestation. The changes in the FL were following: CL converted to FL 1.869 kha, GL converted to FL 19.701 kha, and OL converted to FL 8.679 kha in 2023. Total FL area was 2 030.423 kha in 2023.

#### Methodological issues – methods, activity data, emission factors and parameters

This category includes the calculation of net carbon stock changes in living biomass, DOM and in the mineral soil. Tier 1 and tier 2 approaches (IPCC 2006 GL) were used for calculation of carbon stocks change in living biomass and DOM. Carbon stocks changes in living biomass in the 4.A.2 through the forest regeneration were estimated using the Equation 2.7 (IPCC 2006 GL). The carbon increment is proportional to the extent of afforested areas and the yearly growing biomass. The new afforested areas were determined from the cadastral database. The annual increment of the total tree biomass for four main species (Norway spruce, Scotch pine, European beech and Sessile oak) were selected from experimental database of the NFC-IFRI. These data were published (Priwitzer et al., 2008, Priwitzer et al., 2009 and Pajčík et al., 2011). The annual increment of the above-ground tree biomass (dry mass) for the four main tree species included in the inventory are following: spruce 2.74 t C/ha/y, pine 3.17 t C/ha/y, beech 2.32 t C/ha/y and oak 1.23 t C/ha/y. The activity data comes from representative experimental plots. Then, whole-tree samples including foliage, branches, stem and coarse roots were taken, oven-dried and weighed. Allometric relationships for all tree compartments using tree height and/or diameter on stem base as independent variables were constructed. The tree biomass was measured at the sites and calculated by different compartment (stem, branches, roots and foliage) using allometric functions. Moreover, soil cores for fine roots (diameter up to 2 mm) estimation were taken. Biomass for all tree compartments was calculated on a hectare base.

The annual increments of the below-ground biomass (dry mass) for the four main tree species included in the inventory are following: spruce 0.56 t C/ha/y, pine 0.40 t C/ha/y, beech 0.57 t C/ha/y and oak 0.90 t C/ha/y. The ratio of main tree species from reforestation for different years was taken from the Statistical Office of the Slovak Republic and represented 31% for spruce, 12% for pine, 51% for beech and 6% for oak in 2023.

---

The carbon loss connected with living biomass due to silvicultural cuttings in the subcategory L-FL was assumed to be insignificant (zero). The reason is that the first significant thinning occurs in older age forest stands.

The net changes of carbon stock in dead organic matter (DOM) were estimated in accordance with the guidance of the Tier 1 method (IPCC 2006 GL), using available country specific information. The changes in DOM were estimated separately for deadwood and litter C pools.

According to the NFIs the average C stock of dead wood was calculated on  $6.6 \pm 0.5$  t C/ha for 2005 as well as  $7.4 \pm 0.7$  t C/ha for 2015 in Forest Land category. Using the estimated trend based on these empirical observations, data for the years between these data points were linearly interpolated and extrapolated accordingly beyond that period. The mean net annual accumulation of dead wood over 10-years period is 0.08 t C/ha/y. The net C stocks change of DW was calculated by the Equation 2.23 of the IPCC 2006 GL.

Methodology for carbon estimation in dead wood pool follows conversion of land to forest land just prior to and just following conversion. Most of the categories (CL, GL, OL) does not produce dead wood, so the corresponding carbon pools prior to conversion are zero.

The changes in living biomass and deadwood are assumed to be zero at conversion due to common afforestation practices, if any vegetation exists in Cropland or Grassland it is not removed before conversion to FL and remains in afforested areas. Due to economic reasons, Land converted to FL is located exclusively in mountainous regions of the Carpathians on the steeper slopes with less productive soil, while rich soil in the lowlands remain under managed Cropland or Grassland. Therefore, when converted to Forest Land, existing grass vegetation is not removed to prevent intensive soil erosion on mountain slopes. There is no tree biomass considered present on Grassland. On Cropland, tree biomass is neglected as the Perennial Croplands with tree biomass (orchards, gardens) composes less than 5% of the managed Cropland area. Moreover, orchards and gardens are mostly situated close to built-up area and therefore usually are not subject of conversion to Forest Land.

The net carbon stock change in litter was estimated using the country specific tier 2 approach. It was based on existing data sets from soil inventories and published information (Šály, 1998, Kobza et al., 2002, 2009, 2014, Pavlenda, 2008) with the default assumption of 20 years period for carbon stock equilibrium in "new land-use" conditions. The mean value 8.3 t C/ha for carbon stocks in litter (representing surface organic layer) was used for calculation of net carbon stock change in litter. The mean net annual accumulation of litter over length of transition period is 0.415 t C/ha/y. The Equation 2.23 (IPCC 2006 GL) was used for calculation of annual changes in carbon stocks in litter for this subcategory. The change in litter carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with this subcategory.

The net carbon stock change in mineral soil was estimated using the country specific tier 2 approach. It was based on existing data sets from soil inventories and published information (Šály, 1998, Kobza et al., 2002, 2009, 2014, Pavlenda, 2008) with the default assumption of 20 years period for carbon stock equilibrium in „new land-use“ conditions. The approach for calculation of the organic carbon stocks in soil is consistent with the previous submission. Mean values of soil organic carbon stocks in each category were calculated from datasets of FMS (112 representative monitoring plots in forests) and Soil Monitoring System (318 monitoring plots). Data was recalculated to 30 cm soil layer (topsoil) and compared for three altitudinal zones in each category. The significant changes in soil carbon were caused by land-use change during decades and are only in topsoil (soil layers near the soil surface). Partial results were published in several articles (Barančíková et al. 2013, Barančíková et al. 2016, Pavlenda et al., 2016). The case study using different approach (transections at local level for GL, FL, GL converted to FL) proved very similar results (Pavlenda et al. 2015).

For respective categories, following values (calculated as weighted average) were used in calculations of carbon stock changes in mineral soils (0-30 cm, without any surface organic layer):

- Forest Land 89.02 t C/ha
- Cropland 60.11 t C/ha
- Grassland 74.95 t C/ha
- Settlements 53.85 t C/ha
- Other Land 53.85 t C/ha

The average annual carbon stock change in mineral soil for different conversion of Land to FL was calculated as:

- Annual changes in mineral soil carbon stocks for Land converted to FL = average annual change of SOC over length of transition period (t C/ha/y) \* converted area (kha). Average annual change of SOC over length of transition period = (mean SOC stock of FL - mean SOC stock of Land converted to FL)/20.

The following factors (mean annual change of soil carbon stock) were calculated for different types of conversion:

- CL converted to FL 1.446 t C/ha/y
- GL converted to FL 0.704 t C/ha/y
- S converted to FL 1.758 t C/ha/y
- OL converted to FL 1.758 t C/ha/y

The change in soil carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to Forest Land.

As mentioned in the category FL-FL, the same values as in previous documents were used. For FL, the carbon stock in surface organic layer is separated from carbon stock in mineral soils.

The land-use matrix from 2003 to 2023 is provided in [Table 6.12](#).


The results from the category Land converted to FL are summarized in [Table 6.11](#) and on [Figure 6.11](#).

**Table 6.11: Results for the subcategory Land converted to Forest Land in 2023**

LAND USE CATEGORY	CARBON STOCK CHANGE IN LIVING BIOMASS (kt C)			NET CARBON STOCK CHANGE IN DOM	NET CARBON STOCK CHANGE IN SOIL	NET CO <sub>2</sub> EMISSIONS/REMOVALS
	gains	losses	net change	(kt C)	(kt C)	(kt CO <sub>2</sub> )
Land - FL	45.27	NO	45.27	14.97	31.83	-337.62
GL - FL	29.49	NO	29.49	9.75	13.87	-194.73
CL - FL	2.80	NO	2.80	0.93	2.70	-23.56
WL - FL	NO	NO	NO	NO	NO	NO
S - FL	NO	NO	NO	NO	NO	NO
OL - FL	12.99	NO	12.99	4.30	15.26	-119.33

The estimated removals for Land converted to Forest Land were -337.62 Gg CO<sub>2</sub> in 2023. In 2023, the net carbon stock change in living biomass, DOM and soil from Land converted to Forest Land represented gains of 45.57, 15.07 and 32.02 kt of C respectively.

Figure 6.11: Summary results of CO<sub>2</sub> removals (Gg) in L-FL subcategory in 1990 – 2023



**Table 6.12: The land-use matrix from 2003 – 2023**

Land use	Forest Land managed	Forest Land unmanaged	Cropland annual	Cropland perennial	Grassland managed	Grassland unmanaged	Wetland managed	Wetland unmanaged	Settlements	Other Land	Total unmanaged	Initial area (2003)
Category	(kha)											
Forest Land (managed)	2 000.174	0.000	0.159	0.000	1.078	0.000	0.000	0.000	0.939	1.750	0.000	2 004.100
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	1.869	0.000	1 382.897	0.198	11.120	0.000	0.000	0.000	18.896	15.217	0.000	1 430.197
Cropland perennial	0.000	0.000	3.872	119.301	0.000	0.000	0.000	0.000	0.000	0.000	0.000	123.173
Grassland (managed)	19.701	0.000	13.599	0.000	835.469	0.000	0.000	0.000	7.324	7.413	0.000	883.506
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	212.869	11.802	0.000	224.671
Other Land	8.679	0.000	2.477	0.000	0.031	0.000	0.000	0.000	1.795	130.891	0.000	143.873
<b>Total unmanaged land</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
<b>Final area (2023)</b>	<b>2 030.423</b>	<b>0.000</b>	<b>1 403.004</b>	<b>119.499</b>	<b>847.698</b>	<b>0.000</b>	<b>94.000</b>	<b>0.000</b>	<b>241.823</b>	<b>167.073</b>	<b>0.000</b>	<b>4 903.520</b>
<b>Net change</b>	<b>26.323</b>	<b>0.000</b>	<b>-27.193</b>	<b>-3.674</b>	<b>-35.808</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>	<b>17.152</b>	<b>23.200</b>	<b>0.000</b>	

#### 6.6.4. Biomass Burning (CRT 4.A.2 - 4(IV))

The biomass burning activity in 4.A.2 - 4(V) includes emissions of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O associated with the forest fires and biomass burning on forest areas. The National Forest Centre – Forest Protection Service Activity summarized data from the forest fires (wildfires) since 1999. The emissions from wildfires ([Table 6.13](#)) were calculated according to the Equation 2.27 and Table 2.4 (IPCC 2006 GL) using the default emission factors - available mass of fuel for combustion was used according to Table 2.4 (IPCC 2006 GL).

**Table 6.13: Burned forest area, CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from wildfires in particular years**

Year	AREA BURNED (ha)	CO <sub>2</sub> EMISSIONS (t)	CH <sub>4</sub> EMISSIONS (t)	N <sub>2</sub> O EMISSIONS (t)
1990	23.06	911.86	2.73	0.15
1995	4.94	195.15	0.58	0.03
2000	34.35	1358.27	4.07	0.23
2005	16.31	645.00	1.93	0.11
2010	2.84	112.43	0.34	0.02
2011	5.80	229.22	0.69	0.04
2012	24.55	970.61	2.91	0.16
2013	4.03	159.46	0.48	0.03
2014	2.99	118.17	0.35	0.02
2015	5.92	234.24	0.70	0.04
2016	3.01	119.19	0.36	0.02
2017	4.90	194.17	0.58	0.03
2018	4.05	159.95	0.48	0.03
2019	7.30	288.72	0.86	0.05
2020	7.36	291.03	0.87	0.05
2021	2.44	96.40	0.29	0.02
2022	18.44	729.12	2.18	0.12
2023	0.45	17.62	0.05	0.003


#### Wildfires

Total methane emissions from wildfires in category 4.A.2 were 0.05 t and total emissions of N<sub>2</sub>O were 0.003 t in 2023. Total CO<sub>2</sub> emissions were 17.61 t in 2023.


### 6.7. Cropland (CRT 4.B)

The GHGs emissions and removals in this category were estimated using the 2019 IPCC Refinements methodology the IPCC 2006 GL for AFOLU and national data on area of Cropland and Land converted to Cropland in 2023. The total area of Cropland represented 1 522.503 kha in 2023, i. e. 31.1% of the total country area. This category has been constantly decreasing during reporting period, even since 1970. The total area of Cropland remaining Cropland (CL-CL) represents 1 506.268 kha, of which Annual Cropland remaining Annual Cropland (CLA-CLA) is 1 382.897 kha, Perennial Cropland remaining Perennial Cropland (CLP-CLP) is 119.301 kha, changes from Annual Cropland converted to Perennial Cropland (CLA-CLP) is 0.198 kha and the changes from Perennial Cropland converted to Annual Cropland (CLP-CLA) is 3.872 kha. The changes in the Cropland were following: FL converted to CL 0.159 kha, GL converted to CL 13.599 kha and OL converted to the CL 2.477 kha in 2023 as shown on [Figures 6.12](#) and [6.13](#).

**Figure 6.12:** Development of activity data (kha) for 4.B Cropland in the period 1990 – 2023



**Figure 6.13:** Distribution of Cropland in Slovakia – calculated as a spatial share within individual cadastral units



### 6.7.1. Cropland Remaining Cropland (CRT 4.B.1)

The emissions inventory in this category included net carbon stock change in living biomass of Perennial Cropland remaining Perennial Cropland (CLP-CLP) and carbon stock changes in biomass due to land-use change between Annual Cropland (CLA) and Perennial Cropland (CLP) and net carbon stock change in soil of Annual Cropland remaining Annual Cropland (CLA-CLA) and Perennial Cropland remaining Perennial Cropland (CLP-CLP) and due to land-use change between CLA and CLP. The CLA represented arable land planted with annual crops (cereals, oilseeds, crop roots, technical crops, fodder and other) and its area was 1 382.897 kha in 2023. The CLP including vineyards, orchards, hop-gardens and gardens represented 119.301 kha in 2023.

#### Methodological issues – methods, activity data, emission factors and parameters

Change in biomass carbon stocks of Cropland remaining Cropland were estimated by tier 1 approach.

---

## Changes of carbon stocks in biomass of Annual Cropland remaining Annual Cropland and Perennial Cropland remaining Perennial Cropland

In general, Cropland has no dead wood and only little crop residues or litter, with the exception of agroforestry systems which can be accounted under either Cropland or Forest Land, depending upon definitions adopted by country. Tier 1 approach assumes that dead wood and litter stocks are not present in Cropland or are at equilibrium like in agroforestry systems and orchards.

The carbon stock changes of living biomass in the CLA remaining CLA are estimated to be zero. For annual crops increase in biomass stocks in the CLA remaining CLA in a single year is assumed to be equal to biomass losses from harvest and mortality in the same year – thus there are no net emissions/removals from biomass in the CLA remaining CLA (Chapter 5.2.1.1 of the IPCC 2006 GL).

The emissions/removals were estimated for the changes in woody perennial biomass stocks of the CLP remaining CLP (above-ground and below-ground biomass). So, these emissions/removals were estimated for CLA converted to CLP and vice versa (Chapter 5.3 of the IPCC 2006 GL and in 2019 IPCC Refinements remain unchanged; CLA biomass is not resolved/equilibrium. Also, CLP biomass will not change, as tier 2 is used here for vineyards and orchards). For that purpose, the carbon stock of annual and perennial crops has been estimated and applied in the LUC calculation subsequently. The annual change of carbon stocks in biomass was calculated using the Equation 2.7 of the IPCC 2006 GL.

The immature CLP area accumulates carbon at a rate of approximately 2.35 t for orchards and 4.43 t for vineyards of average carbon stock in living biomass per hectare per year. The emission factors taken from Hungarian inventory was used due to consideration, that carbon accumulation is similar as in Slovakia. The value of above ground biomass carbon stock at harvest is 70.5 t C/ha for orchards and 132.90 t C/ha for vineyards. For gardens and hop-gardens default value was used for CLP (Table 5.1 of the IPCC 2006 GL; Table 5.1 a 5.3 in 2019 IPCC Refinement).

The periodic cuttings, pruning and thinning are not included in the estimation of annual losses in perennial croplands due to low acreage of this areas, lack of historical data and use of this cut material in the production of mulch.

## Changes of carbon stocks in biomass of Annual Cropland converted to Perennial Cropland

Total area of CLA converted to CLP was 0.198 kha in 2023. This type of conversion occurred previous year after several years (to 2017 was zero area of CLA-CLP). The applied method follows entirely the IPCC 2006 GL (Chapter 5.3, Chapter 5.3.1.1). The 2019 IPCC Refinements and the IPCC 2006 GL do not foresee any method for land-use change within the Cropland. CLA and CLP have completely different C stocks and C accumulation rates in biomass and soil. For the calculation of the annual change in carbon stock in living biomass of Land converted to Cropland, the equations 2.15 and 2.16 (IPCC 2006 GL) were applied. For CLP, an annual growth 2.1 t C/ha according to the IPCC 2006 GL (Chapter 5.2.1.2, Table 5.1) was assumed for each year of the whole transition period of 20 years.

Annual change in biomass = conversion area for a transition period of 20 years \*  $\Delta C_{\text{growth}}$  + annual area of currently converted land \*  $L_{\text{conversion}}$

Where:  $L_{\text{conversion}} = C_{\text{after}} - C_{\text{before}}$ ;

$C_{\text{after}}$  = carbon stock immediately after conversion is 0;

$\Delta C_{\text{growth}}$  = default value for perennial crops carbon accumulation rate is 2.1 t C/ha/y (annual growth rate in each year of the whole LUC transition period of 20 years);

$C_{\text{before}}$  = country specific value of carbon stock of annual crops before conversion is 3.25 t C/ha/y (biomass loss accounted only for the year of LUC).

Biomass losses in the year of LUC from CLA to CLP used the country specific average biomass stock in CLA. The average carbon stock of living biomass in CLA was calculated by using country specific



---

data from the ŠÚ SR (Statistical Yearbook of the Slovak Republic, 2016). For all annual crops mentioned in the Statistical Yearbook, the harvested yield biomass (1990 – 2016) has been taken and calculated with use of national coefficient of carbon stocks for crops in total living biomass (Bielek, Jurčová, 2010, Torma and Vilček, 2017). This country specific value (3.25 t C/ha/y) is used for estimates of LUCs to and from CLA and is 35% lower than default value (5.0 t C/ha/y, IPCC 2006 GL).

### Changes of carbon stocks in biomass of Perennial Cropland converted to Annual Cropland

Total land-use change area from CLP converted to CLA was 3.872 kha in 2023. The rationale for these estimates and used methods are described in the [Chapter 6.7.1](#). For the calculation of the annual change in carbon stocks of living biomass of CLP converted to CLA the Equations 2.15 and 2.16 were used (IPCC 2006 GL). According to the 2006 IPCC GL, the gains of the CLA biomass during LUCs to CLA are accounted only once, in the initial year of LUC to CLA ([Chapter 6.7.1](#) in more details):

$$\text{Annual change in biomass} = \text{annual area of currently converted land} * (L_{\text{conversion}} + \Delta C_{\text{growth}})$$

Where:

$$L_{\text{conversion}} = C_{\text{after}} - C_{\text{before}};$$

$C_{\text{after}}$  = carbon stock immediately after conversion is 0;

$C_{\text{before}}$  = country specific value of carbon stock of annual crops before conversion is 3.25 t C/ha/y;

$\Delta C_{\text{growth}}$  = annual growth rate of perennial woody biomass is 2.1 t C/ha/y.

The calculation according to the Austrian methodology was applied.  $C_{\text{before}}$  = country specific value of carbon stock of annual crops before conversion is 3.25 t C/ha/y. Annual change of perennial woody biomass (biomass loss accounted only for the year when the Land use change occurred- CLA-CLP Biomass Loss). This happened for years 2019 and 2020. The difference is that with CLA changed to CLP and vice versa, the values 2.1 (annual growth rate of perennial woody biomass) and 3.25 stand (with CLA changed to CLP only in the year of change) on opposite sides of the equation.

### Changes of carbon stocks in mineral soils of Annual Cropland remaining Annual Cropland and Perennial Cropland remaining Perennial Cropland

The Cropland category was recalculated this year due to a change in the soil management (FMG) and soil land use (FLU), as the new values of these factors were applied according to the 2019 IPCC Refinements of methodological manuals document. CLA set aside instead of Land use (FLU) Long-term cultivated instead of 0.80 we will use 0.77; Tillage (FMG) instead of 1.10 we will use 1.00.

CLP Land use (FLU) 1.00, we will use 0.72; Tillage (FMG) instead of 1.02 we will use 0.98. These changes mainly concerned GHG removal and emissions from the soil. The decrease in GHG removals was caused especially by the FMG factor for CLP. According to the IPCC 2006 GL, value of FMG was 1 ton C/ha/year, and according to the IPCC 2019 Refinement, value is 0.75 tons C/ha/year. Depends on the available capacities and resources, we will improve estimate. Equations for calculating the balance of GHG removals and emissions from biomass and soil from Cropland category and subcategories remain unchanged - no Refinement.

The emissions and removals of the soil carbon stock change in CLA-CLA were calculated using a country specific tier 2 approach. Mean values of soil organic carbon stocks in CLA by the Soil Monitoring System (318 monitoring plots) is 60.11 t C/ha (Barančíková et al. 2013, Barančíková et al. 2016). Mean values of soil organic carbon stocks in CLP was calculated from LUCAS Topsoil Survey (LUCAS data) (Tóth, Jones and Montanarella, 2013). Soil Monitoring System does not contain soil organic carbon stock in CLP, so LUCAS data were used for estimation of the soil carbon stocks of CLP. Mean values of soil organic carbon stocks in CLP (two samples, more samples will be added in the near future) is 66.54 t C/ha (0-30 cm).

The SVK NID text reports only the summary results of GHG removals for the entire CLP category. But this result is the sum of all four subcategories: orchards, vineyards, gardens and hop gardens (these are calculated separately in the calculation file). The chosen factors are the immature CLP area accumulates carbon at a rate of approximately 2.35 t for orchards and 4.43 t for vineyards of average carbon stock in living biomass per hectare per year. The emission factors taken from the Hungarian inventory were used due to consideration, that carbon accumulation for this specific category is similar as in Slovakia. Only for gardens and hop-gardens default value was used for CLP (Table 5.1 of the IPCC 2006 GL; Table 5.1 and 5.3 in 2019 IPCC Refinement).

### Changes of carbon stocks in mineral soils of Annual Cropland converted to Perennial Cropland

The area of CLA converted to CLP changed from 17.266 kha to 0.125 kha from 1990 to 2005. In the year 2018, the area of CLA converted to CLP increased after several years up to 0.150 kha. Total area of CLA converted to CLP was 0.198 kha in 2023. C before = country specific value of carbon stock of annual crops before conversion is 3.25 t C/ha/y. Annual change of perennial woody biomass (biomass loss accounted only for the year when the Land use change occurred- CLA-CLP Biomass Loss). This happened for years 2019 and 2020. The difference is that with CLA changed to CLP and vice versa, the values 2.1 (Annual growth rate of perennial woody biomass) and 3.25 stand (with CLA changed to CLP only in the year of change) on opposite sides of the equation.

According to the Equation 2.25 of the IPCC 2006 GL, annual rates of carbon stock change are estimated as the difference in stocks at two points in time divided by the time dependence of the stock change factors. Annual change in carbon stock of mineral soils in CLA converted to CLP =  $\Delta\text{SOC}_{20}$  \* conversion area for a transition period of 20 years

$$\Delta\text{SOC} = (\text{SOC}_0 - \text{SOC}_{0-T})/20 = 0.322 \text{ t C/ha/y}$$

Where:  $\Delta\text{SOC}_{20}$  = average annual carbon stock change in soils of annual cropland converted to perennial cropland (t C/ha/y) over land-use change transition period of 20 years;  $\text{SOC}_0$  = average c stock in 0-30 cm of CLP soils in Slovakia – 66.54 t C/ha;  $\text{SOC}_{0-T}$  = average c stock in 0-30 cm of CLA soils in Slovakia – 60.11 t C/ha.

For a total area of CLA-CLP (0.198 kha in 2020 and also in 2023), the  $\Delta\text{SOC}_{20}$  is in both years 0.06 kt C.

### Changes of carbon stocks in mineral soils of Perennial Cropland converted to Annual Cropland


The area of CLP converted to CLA changed from 1.435 kha to 3.872 kha from 1990 to 2023. According to the Equation 2.25 of the IPCC 2006 GL, annual rates of carbon stock change are estimated as the difference in stock at two points in time divided by the time dependence of the stock change factors. Annual change in carbon stock of mineral soils in CLP converted to CLA =  $\Delta\text{SOC}_{20}$  \* conversion area for a transition period of 20 years.

$$\Delta\text{SOC} = (\text{SOC}_0 - \text{SOC}_{0-T})/20 = -0.3215 \text{ t C/ha/y}$$

Where:  $\Delta\text{SOC}_{20}$  = average annual carbon stock change in soils of perennial cropland converted to annual cropland (t C/ha/y) over land-use change transition period of 20 years.

For a total area of CLP – CLA (3.872 kha), the  $\Delta\text{SOC}_{20}$  represented -1.25 kt C. [Figure 6.14](#) shows the net CO<sub>2</sub> removals in the category 4.B.1 Cropland remaining Cropland.

**Figure 6.14:** Summary results of CO<sub>2</sub> removals (Gg) in CL-CL subcategory in 1990 – 2023



### 6.7.2. Land Converted to Cropland (CRT 4.B.2)

This category includes all processes connected with the conversion of Land converted to Cropland. Land conversion to Cropland from Forest Land and Grassland usually results in a net loss carbon from biomass and soils to the atmosphere. With regard to changes in carbon stocks in living biomass, only losses for conversion from FL and Grassland were calculated.

#### Methodological issues – methods, activity data, emission factors and parameters

According to the ERT recommendation ([L.15 - draft ARR 2022](#)), Slovakia changed the structure of the AD of forest land converted to cropland by species with weighted tree species proportion, which it used for the calculation of BCEF coefficients and revised BCEF<sub>R</sub> values. Slovakia revised its estimation of biomass CSC of forest land converted to cropland by changing the AD structure by tree species and revised BCEF<sub>R</sub> coefficients for the forest land converted to cropland category of LULUCF with revised BCEF<sub>R</sub> coefficients. Slovakia resubmitted the LULUCF CRT tables with the revised calculation, which was accepted by the ERT. The forest land converted to cropland emissions increased from 2.86 Gg CO<sub>2</sub> eq. to 3.02 Gg CO<sub>2</sub> eq. (5.5%) for 2020 through this resubmission. These revised estimates were also reflected in the entire time series (1990 – 2020) in the resubmission.

Carbon stock changes in biomass were calculated using tier 1 and tier 2 approaches (IPCC 2006 GL). Tier 1 follows the approach used in Land converted to FL where the amount of biomass cleared for Cropland is estimated by multiplying the area converted in one year by the average carbon stock in biomass in FL or GL prior to conversion. For calculation of biomass carbon stocks of FL prior conversion, the annually updated average growing stock volumes, BCEF<sub>R</sub> (0.657 for conifers and 0.853 for broadleaves) and default carbon content (0.51 for coniferous and 0.48 for broadleaves) were used. For biomass carbon stock of GL prior the conversion, default values of 13.6 t/ha for above ground and below ground biomass were used (Table 6.4, IPCC 2006 GL). Amount of biomass after land conversion to Cropland was assumed zero (0 t/ha).

Estimated emissions/removals of carbon in dead organic matter pools following conversion of Forest Land to another type of land-use categories (CL, GL, S, OL) require estimates of the carbon stock just prior to and just after conversion. The data obtained from the two National Forest Inventories (NFIs) realized in 2005/2006 and 2015/2016 was used in estimation of dead wood prior the conversion in FL. The NFIs provide data on the mean dead wood biomass stocks (m<sup>3</sup>/ha) separately for coniferous and broadleaved trees in the following categories: standing dead trees, stumps, coarse lying dead wood and small-sized lying dead wood. Each of the mentioned categories was classified according to decomposition degree as a fresh, hard, soft and decomposed dead wood. The conversion of volume to dry biomass was carried out based on wood density coefficients and using reduction coefficients

---

according to the degrees of decomposition of dead wood (Fresh 1; Hard 0.83; Soft 0.66; Decayed 0.5). The volume was multiplied by the dry wood density coefficients according to the NIML List of Forest Tree Abbreviations (Šebeň 2017) and the above-ground biomass in mass units was calculated. The conversion of biomass to carbon was performed by a coefficient of 0.496 (Šmelko et al., 2011).

To construct the data series for entire reporting period, data of NFI1 to represent year 2005, and NFI2 to represent year 2015. The average C stocks of dead wood represents  $6.6 \pm 0.5$  t C/ha in 2005 as well as  $7.4 \pm 0.7$  t C/ha in 2015 in national conditions. Using the estimated trend based on these empirical observations, data for the years between these data points were linearly interpolated and extrapolated accordingly beyond that period.

Because the Cropland does not produce dead wood, these carbon pools after conversion can be considered as zero (default assumption).

The net carbon stock change in litter was estimated by using the country specific tier 2 approach. It was based on existing data sets from soil inventories and published information (Šály, 1998, Kobza et al., 2002, 2009, 2014, Pavlenda, 2008). The mean value of 8.3 t C/ha/y for carbon stocks in litter (representing surface organic layer) were used for calculation of net carbon stock change in litter. The Equation 2.23 (IPCC 2006 GL) was used for calculation of annual changes in carbon stocks in litter for Land converted to CL. To apply instant oxidation of carbon in litter, litter stock under the “new category” was set to zero and transition period to be one. The change in litter carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to CL.

The calculation of carbon stock change in mineral soils was based on the data from the soil inventory with the default assumption of 20 years' period for carbon stock equilibrium in „new category” conditions. Calculations of carbon stock change in mineral soil as a result of FL and GL conversions to CL were carried out following the IPCC 2006 GL. The net carbon stock change in mineral soils was estimated by using country specific tier 2 approach described in detail in the [Chapter 6.6.3](#) of this Document. For estimation of net carbon stock change in mineral soil, the average carbon stocks per hectare were used. The soil carbon stock was calculated for the depth 30 cm for each category ([Chapter 6.6.3](#)). Current results of monitoring of agricultural soil and updated databases were used for calculation.

The average annual carbon stock change in mineral soil for different conversion of Land to CL was calculated as follows:

- Annual changes in mineral soil C stocks for Land converted to CL = average annual change of SOC over length of transition period (t C/ha/y) x converted area (kha);
- Average annual change of SOC over length of transition period = (mean SOC stock of CL – mean SOC stock of land converted to CL)/20.

The following factors (mean annual change of soil carbon stock) were calculated for different types of conversion:

- FL converted to CL -1.446 t C/ha/y
- GL converted to CL -0.742 t C/ha/y
- S converted to CL 0.313 t C/ha/y
- OL converted to CL 0.313 t C/ha/y


The change in soil carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to Cropland. The land-use matrix from 2003 to 2023 is provided in [Table 6.12](#). The results for the subcategory Land converted to Cropland are summarized in [Table 6.14](#), summary of CO<sub>2</sub> emissions is showed in [Figure 6.15](#).

**Table 6.14: Result for the Land converted to Cropland subcategory in 2023**

LAND USE CATEGORY	CARBON STOCK CHANGE IN LIVING BIOMASS (kt C)			NET CARBON STOCK CHANGE IN DOM	NET CARBON STOCK CHANGE IN SOIL	NET CO <sub>2</sub> EMISSIONS/REMOVALS
	gains	losses	net change	(kt C)	(kt C)	(Gg CO <sub>2</sub> )
Land - CL	NO	-0.01	-0.01	NO	-9.55	35.05
FL – CL	NO	NO	NO	NO	-0.23	0.84
GL – CL	NO	-0.01	-0.01	NO	-10.09	37.05
WL – CL	NO	NO	NO	NO	NO	NO
S – CL	NO	NO	NO	NO	NO	NO
OL – CL	NO	NO	NO	NO	0.78	-2.84

The Land converted to Cropland represents net emissions 35.05 Gg of CO<sub>2</sub> in 2023. In 2023, the net carbon stock change in living biomass, DOM and soil from Land converted to Cropland represented losses of -0.01, and -9.55 kt of C respectively.

**Figure 6.15: Summary of CO<sub>2</sub> emissions (Gg) in L-CL subcategory in 1990 – 2023**



### 6.7.1. Grassland (CRT 4.C)

The GHG emissions and removals in this category were obtained by using the 2019 IPCC Refinements methodology and the IPCC 2006 GL for LULUCF and national data on Grassland and Land converted to Grassland area in 2023. The total area of Grassland represented 847.698 kha in 2023; this is approximately 17.3% of the total country area. Grassland area decreased from 1980 to beginning of 1990 and since this year increased up to 2005. Since 2005, area of Grassland shows moderately decreasing trend. [Figures 6.16](#) and [6.17](#) show activity data and map of Grassland area in Slovakia.

Figure 6.16: Development of activity data (kha) for 4.C Grassland in the period 1990 – 2023





Figure 6.17: Distribution of Grassland in Slovakia – calculated as a spatial share within individual cadastral units



The total area of Grassland remaining Grassland was 835.469 kha in 2023, the changes in Grassland were following: Forest Land converted to Grassland 1.078 kha, Cropland converted to Grassland 11.120 kha, Other Land converted to Grassland 0.031 kha in 2023.

6.8.1. Grassland Remaining Grassland (CRT 4.C.1)

According to the tier 1, no change in living biomass in Grassland remaining Grassland occurred. This approach was used in the emissions/removals estimation in this category. This is a conservative approach for the national conditions, where any application of higher tiers would be justified with respect to data requirements and the expected insignificant stock changes. There were no changes in either type or intensity of management and biomass will be in an approximate steady state (carbon accumulation through plant growth is roughly balanced by losses through grazing, decomposition and fire) in Grassland. The CO<sub>2</sub> emissions are considered insignificant as no change in DOM (dead wood and litter). This is a conservative assumption, if the country did not expect significant changes in categories, disturbance or management regimes within the reporting year (tier 1, IPCC 2006 GL). There are no changes in soil carbon for mineral soils for grassland remaining grassland in case there have been no change to the stock change factors for grassland management (see table 6.2 in the 2006 IPCC Guidance). In CRT Table 4.C.1 notation key “NA” is reported. The limestone application is not a practice

---

in Grassland remaining Grassland category in Slovakia and biomass burning activities are strictly prohibited by the Act No 314/2001 Coll. on Fire Protection.

### 6.8.2. Land Converted to Grassland (CRT 4.C.2)

This category includes all processes connected with conversion of Land into Grassland. For calculation of carbon stock changes in biomass, Tier 1 and tier 2 were used. Tier 1 requires estimate of the biomass of the category before conversion and after conversion. It is assumed, that all biomass is cleared when preparing a site for Grassland, therefore the default value for biomass immediately after conversion is 0 t/ha. Tier 1 follows the approach described in the [Chapter 6.6](#) of this Document where the amount of biomass that is cleared for Grassland is estimated by multiplying the area converted in one year by the average carbon stock in biomass in the Forest Land or Cropland prior to conversion. The default carbon stock values before conversion for the perennial woody crops in accordance with the IPCC 2006 GL, for carbon stocks in CL converted to GL have been implemented. The conversion of perennial CL to GL does not exist in the national conditions. Slovakia estimates and reports the carbon stock change only for CLA converted to CLP and CLP converted to CLA since 2018 submission. This estimation includes the carbon stock changes in living biomass, DOM and mineral soil carbon pools. More information about the AD and EF used is in the [Chapter 6.7.1](#).

#### Methodological issues – methods, activity data, emission factors and parameters

According to the ERT recommendation ([L.16 - draft ARR 2022](#)), Slovakia changed the structure of the AD of forest land converted to grassland by species with weighted tree species proportion, which it used for the calculation of BCEF coefficients and revised BCEF<sub>R</sub> values. Slovakia revised its estimation of biomass CSC of forest land converted to grassland by changing the AD structure by tree species and revised BCEF<sub>R</sub> coefficients for the forest land converted to grassland category of LULUCF with revised BCEF<sub>R</sub> coefficients. Slovakia resubmitted the LULUCF CRT tables with the revised calculation, which was accepted by the ERT. The forest land converted to grassland emissions increased from 7.52 Gg CO<sub>2</sub> eq. to 7.87 Gg CO<sub>2</sub> eq. (4.7%) for 2020 through this resubmission. These revised estimates were also reflected in the entire time series (1990–2020) in the resubmission.

The annually updated average growing stock volumes, BCEF<sub>R</sub> (0.657 for conifers and 0.853 for broadleaves) and default carbon content (0.51 for coniferous and 0.48 for broadleaves) were used for calculation of biomass carbon stocks in FL prior conversion. The default values of 4.7 t C/ha for herbaceous above ground and below ground biomass were used for biomass carbon stock on Grassland prior conversion. Carbon stock from one-year growth Grassland vegetation following the conversion was 13.6 t C/ha (Table 6.4, IPCC 2006 GL).

Estimation of DOM emissions include the emissions from changes in dead wood related to conversion of Forest Land. The calculation procedure is identical with the estimation described in the Land converted to Cropland category.

The net carbon stock change in litter was estimated by using the country specific tier 2. It was based on existing data sets from soil inventories and published information (Šály, 1998, Kobza et al., 2002, 2009, 2014, Pavlenda, 2008). The mean value of 8.3 t C/ha/y for carbon stocks in litter (representing surface organic layer) was used for calculation of net carbon stock change in litter. The Equation 2.23 (IPCC 2006 GL) was used for calculation of annual changes in carbon stocks in litter for Land converted to CL. To apply instant oxidation of carbon in litter, litter stock under the “new category” was set to zero and transition period to be one. The change in litter carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to GL.

The calculation of carbon stock change in mineral soils was based on the data from the soil inventory with the default assumption of 20 years' period for carbon stock equilibrium in „new category” conditions. Calculations of carbon stock change in mineral soil as a result of FL and GL conversions to CL were carried out following the IPCC 2006 GL. For estimation of net carbon stock change in mineral soil, the

average carbon stocks per hectare were used. The soil carbon stock was calculated for the depth 30 cm for each category ([Chapter 6.6.3](#)). Current results of monitoring of agricultural soil and updated databases were used for calculation.

The calculation of carbon stock change in litter was separated from calculations of changes in soil. The information on carbon stocks in surface organic layer of forest soils (based on the data from the soil inventory) was used for calculation of carbon stock change in dead organic matter (FL converted to GL) with the default 20 years' period for carbon stock equilibrium in „new category” conditions. The average annual C stock change in mineral soil for different conversion of the Land converted to GL was calculated as follows:

- Annual changes in mineral soil C stocks for Land converted to GL = average annual change of SOC over length of transition period (t C/ha/y) \* converted area (kha).
- Average annual change of SOC over length of transition period = (mean SOC stock of GL - mean SOC stock of land converted to GL)/20.

The following factors (mean annual change of soil carbon stock) were calculated for different types of conversion:

- FL converted to GL     -0.704 t C/ha/y
- CL converted to GL     +0.742 t C/ha/y
- OL converted to GL     +1.055 t C/ha/y

The change in soil carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to Grassland. The land-use matrix from 2003 to 2023 is provided in [Table 6.12](#). The results of balance in the Land converted to Grassland subcategory are summarized in [Table 6.15](#).


**Table 6.15: Results for Land converted to Grassland subcategory in 2023**

LAND USE CATEGORY	CARBON STOCK CHANGE IN LIVING BIOMASS (kt C)			NET CARBON STOCK CHANGE IN DOM	NET CARBON STOCK CHANGE IN SOIL	NET CO <sub>2</sub> EMISSIONS/REMOVALS
	gains	losses	net change	(kt C)	(kt C)	(Gg CO <sub>2</sub> )
<b>Land - GL</b>	<b>0.07</b>	<b>NO</b>	<b>0.07</b>	<b>NO</b>	<b>7.52</b>	<b>-27.85</b>
FL - GL	NO	NO	NO	NO	-0.76	2.78
CL - GL	0.07	NO	0.07	NO	8.25	-30.51
WL - GL	NO	NO	NO	NO	NO	NO
S - GL	NO	NO	NO	NO	NO	NO
OL - GL	NO	NO	NO	NO	0.03	-0.12

Total removals estimated in this category were -27.85 Gg CO<sub>2</sub> in 2023. The net carbon stock change in mineral soils for this category represented gains of 7.52 kt C, but the net carbon stock change in living biomass from Land converted to Grassland represented the gains of 0.07 kt C in the reporting year 2023. Summary of CO<sub>2</sub> removals are shown on [Figure 6.18](#).



**Figure 6.18:** Summary of CO<sub>2</sub> removals (Gg) in the L-GL subcategory in 1990 – 2023



## 6.8. Wetlands (CRT 4.D)

The responsible body for Wetlands conservation and management in Slovakia is the Ministry of Environment of the Slovak Republic (MŽP SR). The MŽP SR represents the national Administrative Authority for the Convention on Wetlands (Ramsar Convention). The MŽP SR administers the protection of Wetlands, the Integrated River Basin Management and planning, monitoring, national and international cooperation. Practical measures concerning Wetlands conservation, management and restoration are carried out by organisations established by the MŽP SR, especially the State Nature Conservancy of the Slovak Republic, the Slovak Water Management Enterprise (state-owned) and Water Management Research Institute.

The Ministry of Agriculture and Rural Development of the Slovak Republic and its organisations are responsible for the inventory of GHGs within the LULUCF sector. There is ongoing update of the cross-sectoral and the inter-institutional coordination for ensuring necessary collection and processing of wetlands relevant data. Administrative steps were already taken in the area of future cooperation in the Wetlands inventory between the Ministry of Environment, the Ministry of the Agriculture and Rural Development of the Slovak republic and corresponding research institutions (the State Environmental Protection agency and the NPPC-VÚPOP).


Based on the cadastral data the area of this category is 94 kha, corresponding to 1.9% of the whole country area. Wetlands consist of surface waters (water courses and water bodies). The share of this category is unchanged since 1990. Permanent surface waters have no carbon stock by definition.

## 6.9. Settlements (CRT 4.E)


Settlements category was reported separately for the first time in the reporting year 2009. This category represents 4.9% of the total country area. Total area of settlements was 241.823 kha in 2023. The increasing trend of settlements area is visible in the time series. This situation is mostly caused by the development of transport infrastructure, industrial areas, municipal development and raising the standards and infrastructure. It is very often connected with decreased area of Cropland and other categories.

Total area of Settlements remaining Settlements is 212.869 kha, the changes in the Settlements were as follows: FL converted to S 0.939 kha, CL converted to S 18.896 kha, GL converted to S 7.324 kha and OL converted to S 1.795 kha in 2023, as described on [Figures 6.19](#) and [6.20](#).

**Figure 6.19:** Development of activity data (kha) in the 4.E Settlements in the period 1990 – 2023



**Figure 6.20:** Distribution of Settlements in Slovakia – calculated as a spatial share within individual cadastral units



### 6.10.1. Settlements Remaining Settlements (CRT 4.E.1)

For this category, CO<sub>2</sub> emissions are considered insignificant as no change in living biomass, DOM (dead wood and litter) and soil carbon pools is assumed (tier 1, IPCC 2006 GL). This is a conservative assumption, if the country did not expect significant changes in land-use types, disturbance or management regimes within the reporting year.

### 6.10.2. Land Converted to Settlements (CRT 4.E.2)

This category includes all processes connected with conversion of Land into Settlements.

#### Methodological issues – methods, activity data, emission factors and parameters

Tier 1 and tier 2 approaches from the IPCC 2006 GL, Vol. 4 were used for carbon stock changes in biomass calculation. Tier 1 requires estimation of the biomass before and after conversion. It is assumed that all biomass is cleared when preparing a site for Settlements, therefore the default value for biomass immediately after conversion is 0 t/ha. Tier 1 follows the approach where the amount of biomass that is cleared for Settlements is estimated by multiplying the area converted in one year by the average carbon stock in biomass in the FL, CL or GL prior to conversion. The calculation procedure is identical as described in detail in the chapters above.

Estimation of DOM includes the emission changes in dead wood related to conversion of Forest Land. The calculation procedure is identical as described in detail in the Chapter Land converted to Cropland.

The net carbon stock change in litter was estimated by using the country specific tier 2 approach. It was based on existing data sets from soil inventories and published information (Šály, 1998, Kobza et al., 2002, 2009, 2014, Pavlenda, 2008). The mean value of 8.3 t C/ha/y for carbon stocks in litter (representing surface organic layer) was used for calculation of net carbon stock change in litter. The Equation 2.23 (IPCC 2006 GL) was used for calculation of annual changes in carbon stocks in litter for Land converted to CL. To apply instant oxidation of carbon in litter, litter stock under the “new category” was set to zero and transition period to be one.

The calculation of carbon stock changes in mineral soils was based on the data from the soil inventory. The default 20 years period for carbon stock equilibrium in „new category“ conditions was applied. The net carbon stock change in mineral soils was estimated by using country specific tier 2 applying factors for mean annual change of soil carbon stock described below. Net carbon stock change in mineral soil was used for estimation of the average carbon stock per hectare mentioned also in Land converted to FL subcategory. The soil carbon stocks were calculated for the depth to 30 cm for each category. More information is in the [Chapter 6.6.3](#) of this Document.

The average annual C stock change in mineral soil for different conversion of Land to Settlement was calculated as follows:

- Annual changes in mineral soil C (kt) stocks for Land converted to S = average annual change of SOC over length of transition period (t C/ha/y) x converted area (kha);
- Average annual change of SOC = (mean SOC stock of S – mean SOC stock of land converted to S).

The following factors (mean annual change of soil carbon stock) were calculated for different types of conversion:

- FL converted to S        -1.758 t C/ha/y
- CL converted to S        -0.313 t C/ha/y
- GL converted to S        -1.055 t C/ha/y


The change in soil carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to Settlements. The land-use matrix from 2003 to 2023 is provided in [Table 6.12](#). The results for Land converted to Settlements subcategory are summarized in [Table 6.16](#). Summary of CO<sub>2</sub> removals are shown on [Figure 6.21](#).

**Table 6.16: Results for the subcategory Land converted to Settlements in 2023**

LAND USE CATEGORY	CARBON STOCK CHANGE IN LIVING BIOMASS (kt C)			NET CARBON STOCK CHANGE IN DOM	NET CARBON STOCK CHANGE IN SOIL	NET CO <sub>2</sub> EMISSIONS/REMOVALS
	gains	losses	net change	(kt C)	(kt C)	(Gg CO <sub>2</sub> )
<b>Land – S</b>	<b>NO</b>	<b>-5.58</b>	<b>-5.58</b>	<b>-0.11</b>	<b>-15.29</b>	<b>76.94</b>
FL – S	NO	-0.80	-0.80	-0.11	-1.65	9.41
CL – S	NO	-3.21	-3.21	NO	-5.91	33.44
GL – S	NO	-1.57	-1.57	NO	-7.73	34.10
WL – S	NO	NO	NO	NO	NO	NO
OL – S	NO	NO	NO	NO	NO	NO

In the reporting year 2023, the total emissions estimated in this category were 76.94 Gg CO<sub>2</sub>, the net CSC in living biomass, DOM and soil for this category represented losses of -5.58 kt C, -0.11 kt C and -15.29 kt C respectively.

**Figure 6.21:** Summary of CO<sub>2</sub> emissions (Gg) in the subcategory Land-S in 1990 – 2023




## 6.10. Other Land (CRT 4.F)


The emissions and removals of GHGs in this category were estimated using the IPCC 2006 GL and 2019 IPCC Refinement as well as national data on area of Other Land and Land converted to Other Land during the inventory year 2023. Total area of Other Land represented 167.073 kha in 2023, which is 3.4% of the total country area. Other Land area decreased between 1995 and 1997, since that year the trend was balanced and slightly increasing, especially after 2007.

Total area of Other Land remaining Other Land was 130.891 kha, the changes in Other Land were following: FL converted to OL 1.750 kha, CL converted to OL 15.217 kha, GL converted to OL 7.413 kha, S converted to OL 11.802 kha in 2023, as is described on [Figures 6.22](#) and [6.23](#).

**Figure 6.22:** Development of activity data (kha) for 4.F Other Land in the period 1990 – 2023



**Figure 6.23:** Distribution of Other Land in Slovakia – calculated as a spatial share within individual cadastral units



#### 6.11.1. Other Land Remaining Other Land (CRT 4.F.1)

The CO<sub>2</sub> emissions are insignificant as no change in living biomass, DOM (dead wood and litter) and soil carbon pools occurred (tier 1, IPCC 2006 GL) in this category. This is a conservative assumption, if the country did not experience significant changes in land-use types, disturbance or management regimes within the reporting year.

#### 6.11.2. Land Converted to Other Land (CRT4.F.2)

This category includes all processes connected with conversion of Land into Other Land. Tier 1 and tier 2 approaches (IPCC 2006 GL) for carbon stock changes in biomass calculation were used. Tier 1 requires estimates of the biomass before and after conversion. It is assumed that all biomass is cleared when preparing a site for other land, thus the default value for biomass immediately after conversion is 0 t/ha.

#### Methodological issues – methods, activity data, emission factors and parameters

Tier 1 and tier 2 approaches follow the approach described in section Forest Land, where the amount of biomass that is cleared for Other Land is estimated by multiplying the area converted in one year by the average carbon stock in biomass in the Forest Land, Cropland or Grassland prior to conversion. The calculation procedure is identical as described in detail in the chapters above.

Estimation of DOM includes the emissions changes in dead wood in Forest Land. The calculation procedure is identical as described in detail in the chapter Land Converted to Settlements.

The net carbon stock change in litter was estimated using the country specific tier 2. It was based on existing data sets from soil inventories and published information (Šály, 1998, Kobza et al., 2002, 2009, 2014, Pavlenda, 2008) and total loss of litter in the year of conversion. The mean value 8.3 t C/ha/y for carbon stocks in litter was used for calculation of net carbon stock change in litter as follows:

- Annual changes in litter C (kt) stocks for Forest Land converted to OL = mean value of carbon in litter in forests (t C/ha/y) \* converted area (kha).

The change in litter carbon stock in each year was calculated as the sum of annual changes in C stocks for each category associated with FL converted to OL. To apply instant oxidation of carbon in litter, litter stock under the “new category” was set to zero and transition period to one year.

The calculation of carbon stock changes in mineral soils was based on the data from the soil inventory. The default 20 years period for carbon stock equilibrium in „new category“ conditions was applied. The net carbon stock change in mineral soils was estimated by using country specific tier 2 approach applying factors for mean annual change of soil carbon stock described below. Net carbon stock change in mineral soil was used for estimation of the average carbon stock per hectare mentioned also in Land converted to FL subcategory. The soil carbon stocks were calculated for the depth to 30 cm for each category. More information is in the [Chapter 6.6.3](#) of this Document.

The average annual C stock change in mineral soil for different conversion of Land to OL was calculated as follows:

- Annual changes in mineral soil C (kt) stocks for Land converted to OL = average annual change of SOC over length of transition period (t C ha/y) \* converted area (kha).
- Average annual change of SOC (kt) over length of transition period = (mean SOC stock of OL - mean SOC stock of land converted to OL)/20.

The following factors (mean annual change of soil carbon stock) were calculated for different types of conversion:

- FL converted to OL -1.758 t C ha/y
- CL converted to OL -0.313 t C ha/y
- GL converted to OL -0.704 t C ha/y


The change in soil carbon stock in each year was calculated as the sum of annual changes in carbon stocks for each category associated with Land converted to Other Land. The land-use matrix from 2003 to 2023 is provided in [Table 6.12](#). The results from the subcategory Land converted to Other Land are summarized in [Table 6.17](#) and summary of CO<sub>2</sub> emissions during the years on [Figure 6.24](#).

**Table 6.17: Results for the subcategory Land converted to Other Land in 2023**

LAND USE CATEGORY	CARBON STOCK CHANGE IN LIVING BIOMASS (kt C)			NET CARBON STOCK CHANGE IN DOM	NET CARBON STOCK CHANGE IN SOIL	NET CO <sub>2</sub> EMISSIONS/REMOVALS
	gains	losses	net change	(kt C)	(kt C)	(Gg CO <sub>2</sub> )
<b>Land - OL</b>	<b>NO</b>	<b>-7.62</b>	<b>-7.62</b>	<b>-0.93</b>	<b>-15.66</b>	<b>88.78</b>
FL – OL	NO	-6.52	-6.52	-0.93	-3.08	38.59
CL – OL	NO	-1.10	-1.10	NO	-4.76	21.51
GL – OL	NO	NO	NO	NO	-7.82	28.68
WL – OL	NO	NO	NO	NO	NO	NO
S - OL	NO	NO	NO	NO	NO	NO

Total emissions estimated in this category were 88.78 Gg CO<sub>2</sub> in 2023. The net carbon stock change in living biomass, DOM and soil for this category represented losses of -7.62 kt C, -0.93 kt C and -15.66 kt C, respectively.

**Figure 6.24:** Summary of CO<sub>2</sub> emissions (Gg) in L-OL subcategory in 1990 – 2023



### 6.11. Direct and indirect nitrous oxide (N<sub>2</sub>O) emissions from nitrogen (N) inputs to managed soils (CRT 4(I))

Direct & indirect nitrous oxide (N<sub>2</sub>O) emissions from nitrogen (N) inputs to managed soils (CRT 4 I):

There are no direct and indirect N<sub>2</sub>O emissions from N fertilization on Forest Land, Wetlands or Settlements as there is no practice of nitrogen fertilization of forest stands in Slovakia.

### 6.12. Emissions and removals from drainage and rewetting and other management of organic and mineral soils (CRT 4(II))

Emissions and removals from drainage and rewetting and other management of organic and mineral soils (CRT 4 II):

There are no reported CO<sub>2</sub> and non-CO<sub>2</sub> emissions related to drainage and rewetting and other management of organic and mineral soils. The reason is very simple, because the drainage and rewetting and other management of organic and mineral soils are no practice in Slovakia. Only few spots of wet forest soils classified as peat land exist in Slovakia, they are very rare and therefore this land belongs to protected areas without active management. According to (Stanová et al., 2000) the area of peat lands in Slovakia covered only 2 773 ha in 2000.

### 6.13. Direct and indirect nitrous oxide (N<sub>2</sub>O) emissions from nitrogen (N) mineralization/immobilization associated with loss/gain of soil organic matter resulting from change of land use or management of mineral soils (CRT 4(III))

The direct N<sub>2</sub>O emissions (the annual release of N<sub>2</sub>O from soils due to mineralisation of soil organic matter after disturbance) were calculated by default tier 1 (Equations 11.8, IPCC 2006 GL). N<sub>2</sub>O emissions were estimated based on the detected changes in mineral soils on respective areas of FL

and GL converted to CL, S, OL using default emission factor 0.0125 kg N<sub>2</sub>O-N/kg N, and C:N ratio = 12. Direct N<sub>2</sub>O emissions from N mineralization/immobilization are summarized in [Table 6.18](#) and on [Figure 6.25](#).


**Table 6.18: Results for 4(III) – Direct & indirect N<sub>2</sub>O emissions from N mineralization/immobilization in LULUCF sector in 2023**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	ACTIVITY DATA AND OTHER RELATED INFORMATION		IMPLIED EMISSION FACTORS		N <sub>2</sub> O EMISSIONS		
	Land area converted	N mineralised in mineral soils associated with loss of soil C from soil organic matter	N <sub>2</sub> O–N emissions per area	N <sub>2</sub> O–N emissions per unit of N lost through leaching and run-off	Direct Emissions	Indirect Emissions	Total Emissions
CATEGORY	<i>kha</i>	<i>t N/year</i>	<i>kg N<sub>2</sub>O–N/ha</i>	<i>(kg N<sub>2</sub>O–N/kg N)</i>	<i>kt</i>		
<b>Total all land-use categories</b>	160.38	3 502.62	0.24	0.00	0.06	0.01	0.07
<b>A. Forest land</b>	NO	NO	NO	NO	NO	NO	NO
1. Forest land remaining forest land	NO	NO	NO	NO	NO	NO	NO
2. Lands converted to forest land	NO	NO	NO	NO	NO	NO	NO
<b>B. Cropland</b>	13.76	860.03	1.23	0.00	0.03	0.00	0.03
2. Lands converted to cropland	13.76	860.03	1.23	0.00	0.03	0.00	0.03
<b>C. Grasslands</b>	1.08	63.24	0.47	0.00	0.00	0.00	0.00
1. Grasslands remaining grasslands	NO	NO	NO	NO	NO	NO	NO
2. Lands converted to grasslands	1.08	63.24	0.47	0.00	0.00	0.00	0.00
<b>D. Wetlands</b>	94.00	NO	NO	NO	NO	NO	NO
1. Wetlands remaining wetlands	94.00	NO	NO	NO	NO	NO	NO
2. Lands converted to wetlands	NO	NO	NO	NO	NO	NO	NO
<b>E. Settlements</b>	27.16	1 274.34	0.38	0.00	0.02	0.00	0.02
1. Settlements remaining settlements	NO	NO	NO	NO	NO	NO	NO
2. Lands converted to settlements	27.16	1 274.34	0.38	0.00	0.02	0.00	0.02
<b>F. Other land</b>	24.38	1 305.01	0.43	0.00	0.02	0.00	0.02
2. Lands converted to other land	24.38	1 305.01	0.43	0.00	0.02	0.00	0.02

The indirect nitrous oxide (N<sub>2</sub>O) emissions from managed soil were calculated using Equation 11.10 with *FSOM* based on Equation 11.8, *FracLEACH-(H)* (0.30 - default Table 11.3) and *EF5* (0.0075 - default Table 11.3) of the IPCC 2006 GL. Time series was calculated and included firstly in 2018 submission. The resulting values are reported in CRT Table 4(III) and on [Figure 6.25](#) and [in Table 6.18](#). Indirect N<sub>2</sub>O emissions from Nitrogen Leaching and Run-off represented 0.01 Gg in 2023.



**Figure 6.25:** The direct and indirect N<sub>2</sub>O emissions (Gg) from N mineralization/immobilization in LULUCF sector in 1990 – 2023



## 6.14. Biomass Burning (CRT 4(IV))

Calculation of GHG emissions from biomass burning is included in the categories Forest Land remaining Forest Land as well as Land converted to Forest Land. Biomass burning is not common practice in Cropland, Grassland or another land use category in Slovakia, these activities are strictly prohibited by the Act No 314/2001 Coll. on Fire Protection.

## 6.15. Harvested Wood Products (HWP) (CRT4.Gs1-2)

Slovakia started to report on the carbon stock changes and associated emissions and removals of CO<sub>2</sub> from the Harvested Wood Products (HWP) pool in 2015. The wood products in the country define HWP activities as a carbon pool. This carbon pool includes products generated from the wood production in the categories FL remaining FL and Land converted to FL. Harvested timber is converted into a wide variety of wood products. Their carbon content moves through different levels during their life cycle. After their use, products are recycled in some cases and ultimately burned or deposited in landfills where they slowly decay (reported in Waste sector). The carbon stored in wood, which was initially captured from the atmosphere, is finally released back into the atmosphere.

For the carbon balance estimation, the round wood is split into industrial round wood and fuelwood. Contrary to the energetic use of wood (fuelwood) for which an instantaneous oxidation is applied, the long-term used HWP as sawn wood, wood-based panels and paper represent a carbon pool with specific half-lives.

For the assessment, the half-lives were applied according to Table 2.8.2 in the IPCC 2006 GL: 35 years for sawn wood, 25 years for wood-based panels and 2 years for paper products were used. According to the ERT recommendation (L.19 - draft ARR 2022), Slovakia provides further information on parameters for estimating CSC for HWP, following default conversion factors (from the Kyoto Protocol Supplement, table 2.8.1) for estimating CSC of HWP were used: sawnwood (aggregate) 0.229, wood-based panels (aggregate) 0.269 and paper and paperboard (aggregate) 0.386.

The approach applied for HWP accounting calculates delayed emissions based on the annual stock change of semi-finished wood products using the first order decay function following Equation 12.1 (Chapter 12, IPCC 2006 GL). The carbon stock changes in forests are estimated in the 4.A (FL).

### 6.17.1. Methodological Issues – Methods, Activity Data, Emission Factors

The activity data (production and trade of sawn wood, wood-based panels and paper and paperboard) are taken from the [FAO database](#) on wood production and trade. The data are available since 1961, however, data for Slovakia (SR) and the Czech Republic (ČR) are aggregated before the split of Czechoslovakia (ČS) in 1993. To calculate the share of the SR and the ČR on individual HWP in the period 1961 – 1992, ČS figures were multiplied by the country specific share on the sum of figures for both countries in the period of five years 1993 – 1997 (Raši et al. 2015), i.e., correspondingly as applied earlier in the Czech Republic (Cienciala & Palán 2014).

The share of the ČR and SR production, import and export quantities of main HWP categories, calculated as an average of country specific shares according to the FAO data in the period 1993 – 1997, is provided in [Table 6.19](#).

**Table 6.19: The share of the ČR and SR on the HWP in the period 1993 – 1997 and default half-lives**

WOOD PRODUCT	FAO CODE	PRODUCTION		IMPORT		EXPORT		DEFAULT HALF-LIFE (y)
		ČR	SR	ČR	SR	ČR	SR	
Sawn wood	1 872	0.834	0.166	0.868	0.132	0.723	0.277	35
Wood based boards	1 873	0.716	0.284	0.719	0.281	0.851	0.149	25
Paper and paperboards	1 876	0.655	0.345	0.772	0.228	0.598	0.402	2

The change in carbon stocks was estimated separately for each product category by applying Equation 2.8.4 (IPCC 2013 GL). Instantaneous oxidation was applied to HWPs originating from deforestation, which results in a conservative estimate of carbon stock changes in the HWP-pool. The results of CO<sub>2</sub> gains and losses from domestically produced and used HWP are provided in [Table 6.20](#) and on [Figure 6.26](#).


**Table 6.20: Greenhouse gas emissions (positive values) and removals (negative values) from HWP from Forest Land in particular years**

CO <sub>2</sub> emissions and removals from HWP	1990	1995	2000	2005	2010	2015
	<i>Net Emissions/Removals in Gg of CO<sub>2</sub> eq.</i>					
4.G (UNFCCC)	-470.4	-58.8	-920.1	-1 996.5	-1 334.5	-940.7
gains sawn wood	644.3	528.7	1 027.0	2 144.2	1 972.9	1 235.6
gains wood panels	381.9	327.9	330.0	582.4	619.5	866.0
gains paper	606.8	382.8	1 107.8	993.8	710.3	770.2
losses sawn wood	-482.8	-498.8	-526.2	-593.8	-726.0	-785.4
losses wood panels	-268.6	-277.6	-282.0	-299.0	-357.5	-392.7
losses paper	-411.2	-404.2	-736.5	-831.1	-884.6	-752.9

CO <sub>2</sub> emissions and removals from HWP	2018	2019	2020	2021	2022	2023
	<i>Net Emissions/Removals in Gg of CO<sub>2</sub> eq.</i>					
4.G (UNFCCC)	-889.2	-644.9	-247.3	-382.2	-144.4	-291.9
gains sawn wood	1 231.2	1 125.0	904.1	950.8	977.6	908.4
gains wood panels	921.0	882.9	708.4	692.9	524.9	656.6
gains paper	770.8	671.4	652.0	756.4	660.2	746.5
losses sawn wood	-812.6	-819.7	-823.6	-825.6	-828.3	-830.6
losses wood panels	-436.9	-449.6	-459.1	-465.7	-469.6	-472.9
losses paper	-784.4	-765.0	-734.6	-726.6	-720.4	-716.2

According to the ERT recommendation (L.18 - draft ARR 2022), Slovakia provides an explanation of the trend of CSC of HWP. The CSC of HWP follows the production approach, and the real use of wood products in Slovakia differs owing to trade with wood products. The HWP production structure in countries differs according to the wood industry structure. HWP production culminated in 2006 – 2007, just before the 2008 global financial crisis; in Slovakia, the production of sawnwood also accelerated owing to greater availability of wood processed after the destruction of spruce stands by a windstorm in November 2004. While the production of wood-based panels, paper and paperboard is more stable, sawnwood shows higher fluctuations. The wood production and processing sectors in Slovakia as a relatively small country are sensitive to disturbances, for example, the availability of wood due to disturbances in forests, technological processes in wood-processing factories and the situation in the wood products market.

**Figure 6.26:** CO<sub>2</sub> emissions (positive values) and removals (negative values) from HWP in Slovakia in 1990 – 2023 originating from Forest Land



The course of carbon stored in the HWP pool (Figure 6.26) shows that the 1990-2000 following 1990 was characterized by balanced losses and gains of carbon in the pool and a trend of increasing carbon gains in sawnwood and paper is evident. The second decennium was characterized by the growth of the production of sawnwood and wood-based panels and increasing carbon gains in these HWP. Later years are characterized by a drop of production in all HWP categories, which is reflected in the annual CSC in HWP (Figure 6.26) and 2008 (the start of the economic crisis) can be identified as a break point when the trend of increasing gains in the HWP carbon pool turned into a decrease. It is noticeable that in the years since 2008 felling in Slovakia has been higher than in the previous period, indicating an increase in an alternative use of wood, such as for energy purposes. The inventory results indicate that the HWP pool is a carbon sink; however, if the market does not recover and the production stagnates or drops down, the HWP pool may become a source of carbon emissions owing to the decline in the higher gains accumulated in the past. In addition, since 2018 there has been a decrease in timber harvesting in Slovakia, which has caused a decrease in the supply of wood to the domestic market. The decrease in the wood supply since 2019 was due to a decrease in timber harvesting, mainly owing to the coronavirus disease 2019 pandemic, restrictions by nature conservation authorities and the unfavourable situation in the softwood market.

## Annex A6.1. Land-Use Matrix

**Table A6.1:** Land-use matrixes identifying annual conversions among the LUC for the period 1990 – 2022, describing initial and final areas of LUC (kha)

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1989)
Forest Land (managed)	1 985.219	0.000	0.010	0.000	0.353	0.000	0.000	0.000	0.028	0.418	0.000	1 986.028
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.088	0.000	1 507.845	0.000	0.754	0.000	0.000	0.000	0.352	0.000	0.000	1 509.039
Cropland perennial	0.000	0.000	0.203	130.675	0.000	0.000	0.000	0.000	0.000	0.000	0.000	131.081
Grassland (managed)	1.421	0.000	1.407	0.000	807.184	0.000	0.000	0.000	1.293	1.391	0.000	812.696
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	124.361	0.747	0.000	125.108
Other Land	2.261	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	243.307	0.000	245.568
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1990)	1 988.989	0.000	1 509.465	130.878	808.291	0.000	94.000	0.000	126.034	245.863	0.000	4 903.520
Net change	2.961	0.000	0.426	-0.203	-4.405	0.000	0.000	0.000	0.926	0.295	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1990)
Forest Land (managed)	1 988.001	0.000	0.045	0.000	0.678	0.000	0.000	0.000	0.075	0.190	0.000	1 988.989
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.012	0.000	1 507.130	0.000	2.323	0.000	0.000	0.000	0.000	0.000	0.000	1 509.465
Cropland perennial	0.000	0.000	0.486	129.906	0.000	0.000	0.000	0.000	0.000	0.000	0.000	130.878
Grassland (managed)	0.325	0.000	0.941	0.000	806.475	0.000	0.000	0.000	0.356	0.194	0.000	808.291
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	126.034	0.000	0.000	126.034

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1990)
Other Land	1.626	0.000	0.144	0.000	0.000	0.000	0.000	0.000	0.126	243.967	0.000	245.863
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1991)	1 989.964	0.000	1 508.746	130.392	809.476	0.000	94.000	0.000	126.591	244.351	0.000	4 903.520
Net change	0.975	0.000	-0.719	-0.486	1.185	0.000	0.000	0.000	0.557	-1.512	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1991)
Forest Land (managed)	1 989.640	0.000	0.002	0.000	0.146	0.000	0.000	0.000	0.063	0.113	0.000	1 989.964
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.202	0.000	1 484.552	0.000	22.173	0.000	0.000	0.000	0.492	1.327	0.000	1 508.746
Cropland perennial	0.000	0.000	0.692	129.008	0.000	0.000	0.000	0.000	0.000	0.000	0.000	130.392
Grassland (managed)	0.196	0.000	0.793	0.000	808.322	0.000	0.000	0.000	0.165	0.000	0.000	809.476
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	126.591	0.000	0.000	126.591
Other Land	1.069	0.000	0.000	0.000	0.770	0.000	0.000	0.000	0.174	242.338	0.000	244.351
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1992)	1 991.107	0.000	1 486.039	129.700	831.411	0.000	94.000	0.000	127.485	243.778	0.000	4 903.520
Net change	1.143	0.000	-22.707	-0.692	21.935	0.000	0.000	0.000	0.894	-0.573	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1992)
Forest Land (managed)	1 990.741	0.000	0.002	0.000	0.175	0.000	0.000	0.000	0.071	0.118	0.000	1 991.107
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.008	0.000	1 480.682	0.000	4.595	0.000	0.000	0.000	0.285	0.469	0.000	1 486.039
Cropland perennial	0.000	0.000	0.953	127.794	0.000	0.000	0.000	0.000	0.000	0.000	0.000	129.700
Grassland (managed)	0.227	0.000	0.975	0.000	829.862	0.000	0.000	0.000	0.268	0.079	0.000	831.411

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1992)
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	127.485	0.000	0.000	127.485
Other Land	0.487	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.158	243.133	0.000	243.778
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1993)	1 991.463	0.000	1 482.612	128.747	834.632	0.000	94.000	0.000	128.267	243.799	0.000	4 903.520
Net change	0.356	0.000	-3.427	-0.953	3.221	0.000	0.000	0.000	0.782	0.021	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1993)
Forest Land (managed)	1 991.112	0.000	0.014	0.000	0.186	0.000	0.000	0.000	0.025	0.126	0.000	1 991.463
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.019	0.000	1 481.597	0.000	0.869	0.000	0.000	0.000	0.127	0.000	0.000	1 482.612
Cropland perennial	0.000	0.000	0.767	127.213	0.000	0.000	0.000	0.000	0.000	0.000	0.000	128.747
Grassland (managed)	0.308	0.000	0.553	0.000	833.771	0.000	0.000	0.000	0.000	0.000	0.000	834.632
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	128.267	0.000	0.000	128.267
Other Land	0.232	0.000	0.292	0.000	0.000	0.000	0.000	0.000	0.044	243.231	0.000	243.799
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1994)	1 991.671	0.000	1 483.223	127.980	834.826	0.000	94.000	0.000	128.463	243.357	0.000	4 903.520
Net change	0.208	0.000	0.611	-0.767	0.194	0.000	0.000	0.000	0.196	-0.442	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1994)
Forest Land (managed)	1 991.536	0.000	0.002	0.000	0.063	0.000	0.000	0.000	0.023	0.047	0.000	1 991.671

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1994)
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.028	0.000	1 477.809	0.000	5.386	0.000	0.000	0.000	0.000	0.000	0.000	1 483.223
Cropland perennial	0.000	0.000	0.465	127.050	0.000	0.000	0.000	0.000	0.000	0.000	0.000	127.980
Grassland (managed)	0.556	0.000	0.725	0.000	833.333	0.000	0.000	0.000	0.212	0.000	0.000	834.826
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	128.463	0.000	0.000	128.463
Other Land	0.137	0.000	0.103	0.000	0.243	0.000	0.000	0.000	0.291	242.583	0.000	243.357
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1995)	1 992.257	0.000	1 479.104	127.515	839.025	0.000	94.000	0.000	128.989	242.630	0.000	4 903.520
Net change	0.586	0.000	-4.119	-0.465	4.199	0.000	0.000	0.000	0.526	-0.727	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1995)
Forest Land (managed)	1 991.789	0.000	0.098	0.000	0.280	0.000	0.000	0.000	0.032	0.058	0.000	1 992.257
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.107	0.000	1 470.639	0.000	4.015	0.000	0.000	0.000	0.474	0.000	0.000	1 479.104
Cropland perennial	0.000	0.000	0.245	126.674	0.000	0.000	0.000	0.000	0.000	0.000	0.000	127.515
Grassland (managed)	1.113	0.000	0.610	0.000	837.302	0.000	0.000	0.000	0.000	0.000	0.000	839.025
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	128.989	0.000	0.000	128.989
Other Land	0.357	0.000	0.000	0.000	0.117	0.000	0.000	0.000	66.648	175.508	0.000	242.630
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1996)	1 993.366	0.000	1 472.124	126.919	841.714	0.000	94.000	0.000	196.143	175.566	0.000	4 903.520
Net change	1.109	0.000	-3.443	-0.245	2.689	0.000	0.000	0.000	67.154	-67.064	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1996)
Forest Land (managed)	1 992.978	0.000	0.026	0.000	0.203	0.000	0.000	0.000	0.065	0.094	0.000	1 993.366
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.130	0.000	1 470.639	0.000	4.634	0.000	0.000	0.000	0.164	0.000	0.000	1 472.124
Cropland perennial	0.000	0.000	0.245	126.674	0.000	0.000	0.000	0.000	0.000	0.000	0.000	126.919
Grassland (managed)	0.311	0.000	1.214	0.000	840.189	0.000	0.000	0.000	0.000	0.000	0.000	841.714
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	196.143	0.000	0.000	196.143
Other Land	2.954	0.000	0.000	0.000	0.565	0.000	0.000	0.000	22.212	149.835	0.000	175.566
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1997)	1 996.373	0.000	1 472.124	126.919	845.591	0.000	94.000	0.000	218.584	149.929	0.000	4 903.520
Net change	3.007	0.000	-3.443	-0.245	3.877	0.000	0.000	0.000	22.441	-25.637	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1997)
Forest Land (managed)	1 995.995	0.000	0.004	0.000	0.294	0.000	0.000	0.000	0.000	0.080	0.000	1 996.373
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.067	0.000	1 466.916	0.000	4.724	0.000	0.000	0.000	0.000	0.417	0.000	1 472.124
Cropland perennial	0.000	0.000	0.675	125.569	0.000	0.000	0.000	0.000	0.000	0.000	0.000	126.919
Grassland (managed)	0.845	0.000	1.575	0.000	843.171	0.000	0.000	0.000	0.000	0.000	0.000	845.591
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	218.084	0.500	0.000	218.584
Other Land	1.376	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	148.553	0.000	149.929
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000



CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1997)
Final area (1998)	1 998.283	0.000	1 469.170	126.244	848.189	0.000	94.000	0.000	218.084	149.550	0.000	4 903.520
Net change	1.910	0.000	-2.954	-0.675	2.598	0.000	0.000	0.000	-0.500	-0.379	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1998)
Forest Land (managed)	1 997.986	0.000	0.009	0.000	0.086	0.000	0.000	0.000	0.029	0.173	0.000	1 998.283
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.067	0.000	1 458.684	0.000	10.057	0.000	0.000	0.000	0.287	0.075	0.000	1 469.170
Cropland perennial	0.000	0.000	1.042	124.160	0.000	0.000	0.000	0.000	0.000	0.000	0.000	126.244
Grassland (managed)	0.831	0.000	0.868	0.000	846.284	0.000	0.000	0.000	0.000	0.206	0.000	848.189
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	218.084	0.000	0.000	218.084
Other Land	1.204	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.027	148.319	0.000	149.550
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (1999)	2 000.088	0.000	1 460.603	125.202	856.427	0.000	94.000	0.000	218.427	148.773	0.000	4 903.520
Net change	1.805	0.000	-8.567	-1.042	8.238	0.000	0.000	0.000	0.343	-0.777	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1999)
Forest Land (managed)	1 999.961	0.000	0.005	0.000	0.023	0.000	0.000	0.000	0.008	0.091	0.000	2 000.088
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.096	0.000	1 447.768	0.000	12.214	0.000	0.000	0.000	0.244	0.281	0.000	1 460.603
Cropland perennial	0.000	0.000	0.247	124.708	0.000	0.000	0.000	0.000	0.000	0.000	0.000	125.202
Grassland (managed)	0.693	0.000	2.471	0.000	852.983	0.000	0.000	0.000	0.192	0.088	0.000	856.427
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (1999)
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	218.250	0.177	0.000	218.427
Other Land	0.503	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.643	147.627	0.000	148.773
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2000)	2 001.253	0.000	1 450.491	124.955	865.220	0.000	94.000	0.000	219.337	148.264	0.000	4 903.520
Net change	1.165	0.000	-10.112	-0.247	8.793	0.000	0.000	0.000	0.910	-0.509	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2000)
Forest Land (managed)	2 000.951	0.000	0.039	0.000	0.101	0.000	0.000	0.000	0.040	0.122	0.000	2 001.253
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.013	0.000	1 437.399	0.000	12.113	0.000	0.000	0.000	0.212	0.754	0.000	1 450.491
Cropland perennial	0.000	0.000	1.129	122.697	0.000	0.000	0.000	0.000	0.000	0.000	0.000	124.955
Grassland (managed)	0.422	0.000	2.596	0.000	862.202	0.000	0.000	0.000	0.000	0.000	0.000	865.220
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	219.337	0.000	0.000	219.337
Other Land	0.743	0.000	0.000	0.000	0.000	0.000	0.000	0.000	2.886	144.635	0.000	148.264
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2001)	2 002.129	0.000	1 441.163	123.826	874.416	0.000	94.000	0.000	222.475	145.511	0.000	4 903.520
Net change	0.876	0.000	-9.328	-1.129	9.196	0.000	0.000	0.000	3.138	-2.753	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2001)
Forest Land (managed)	2 001.980	0.000	0.006	0.000	0.064	0.000	0.000	0.000	0.021	0.058	0.000	2 002.129
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.008	0.000	1 431.567	0.000	8.980	0.000	0.000	0.000	0.263	0.345	0.000	1 441.163

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2001)
Cropland perennial	0.000	0.000	0.535	122.756	0.000	0.000	0.000	0.000	0.000	0.000	0.000	123.826
Grassland (managed)	0.509	0.000	1.094	0.000	872.813	0.000	0.000	0.000	0.000	0.000	0.000	874.416
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	222.475	0.000	0.000	222.475
Other Land	0.276	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.596	144.639	0.000	145.511
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2002)	2 002.773	0.000	1 433.202	123.291	881.857	0.000	94.000	0.000	223.355	145.042	0.000	4 903.520
Net change	0.644	0.000	-7.961	-0.535	7.441	0.000	0.000	0.000	0.880	-0.469	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2002)
Forest Land (managed)	2 002.452	0.000	0.009	0.000	0.185	0.000	0.000	0.000	0.065	0.062	0.000	2 002.773
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.050	0.000	1 428.082	0.000	4.562	0.000	0.000	0.000	0.379	0.129	0.000	1 433.202
Cropland perennial	0.000	0.000	0.118	123.055	0.000	0.000	0.000	0.000	0.000	0.000	0.000	123.291
Grassland (managed)	1.110	0.000	1.988	0.000	878.759	0.000	0.000	0.000	0.000	0.000	0.000	881.857
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	223.355	0.000	0.000	223.355
Other Land	0.488	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.872	143.682	0.000	145.042
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2003)	2 004.100	0.000	1 430.197	123.173	883.506	0.000	94.000	0.000	224.671	143.873	0.000	4 903.520
Net change	1.327	0.000	-3.005	-0.118	1.649	0.000	0.000	0.000	1.316	-1.169	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2003)
Forest Land (managed)	2 003.934	0.000	0.005	0.000	0.020	0.000	0.000	0.000	0.050	0.091	0.000	2 004.100
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.086	0.000	1 427.075	0.000	2.156	0.000	0.000	0.000	0.517	0.363	0.000	1 430.197
Cropland perennial	0.000	0.000	0.073	123.027	0.000	0.000	0.000	0.000	0.000	0.000	0.000	123.173
Grassland (managed)	0.815	0.000	3.443	0.000	878.878	0.000	0.000	0.000	0.370	0.000	0.000	883.506
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	224.427	0.244	0.000	224.671
Other Land	0.091	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.192	143.590	0.000	143.873
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
<b>Final area (2004)</b>	<b>2 004.926</b>	<b>0.000</b>	<b>1 430.596</b>	<b>123.100</b>	<b>881.054</b>	<b>0.000</b>	<b>94.000</b>	<b>0.000</b>	<b>225.556</b>	<b>144.288</b>	<b>0.000</b>	<b>4 903.520</b>
Net change	0.826	0.000	0.399	-0.073	-2.452	0.000	0.000	0.000	0.885	0.415	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2004)
Forest Land (managed)	2 004.392	0.000	0.015	0.000	0.219	0.000	0.000	0.000	0.038	0.262	0.000	2 004.926
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.023	0.000	1 428.075	0.000	1.146	0.000	0.000	0.000	0.601	0.751	0.000	1 430.596
Cropland perennial	0.000	0.000	0.443	122.214	0.000	0.000	0.000	0.000	0.000	0.000	0.000	123.100
Grassland (managed)	0.455	0.000	0.506	0.000	879.918	0.000	0.000	0.000	0.175	0.000	0.000	881.054
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	225.405	0.151	0.000	225.556
Other Land	0.364	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.038	143.886	0.000	144.288
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
<b>Final area (2005)</b>	<b>2 005.234</b>	<b>0.000</b>	<b>1 429.039</b>	<b>122.657</b>	<b>881.283</b>	<b>0.000</b>	<b>94.000</b>	<b>0.000</b>	<b>226.257</b>	<b>145.050</b>	<b>0.000</b>	<b>4 903.520</b>

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2004)
Net change	0.308	0.000	-1.557	-0.443	0.229	0.000	0.000	0.000	0.701	0.762	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2005)
Forest Land (managed)	2 004.995	0.000	0.000	0.000	0.109	0.000	0.000	0.000	0.024	0.106	0.000	2 005.234
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.044	0.000	1 426.698	0.000	0.984	0.000	0.000	0.000	0.801	0.512	0.000	1 429.039
Cropland perennial	0.000	0.000	0.207	122.243	0.000	0.000	0.000	0.000	0.000	0.000	0.000	122.657
Grassland (managed)	0.504	0.000	0.452	0.000	879.779	0.000	0.000	0.000	0.366	0.182	0.000	881.283
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	225.901	0.356	0.000	226.257
Other Land	1.397	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	143.653	0.000	145.050
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2006)	2 006.940	0.000	1 427.357	122.450	880.872	0.000	94.000	0.000	227.092	144.809	0.000	4 903.520
Net change	1.706	0.000	-1.682	-0.207	-0.411	0.000	0.000	0.000	0.835	-0.241	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2006)
Forest Land (managed)	2 006.486	0.000	0.068	0.000	0.144	0.000	0.000	0.000	0.047	0.195	0.000	2 006.940
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.065	0.000	1 424.648	0.000	1.085	0.000	0.000	0.000	0.742	0.817	0.000	1 427.357
Cropland perennial	0.000	0.000	0.368	121.714	0.000	0.000	0.000	0.000	0.000	0.000	0.000	122.450
Grassland (managed)	0.365	0.000	0.811	0.000	879.692	0.000	0.000	0.000	0.004	0.000	0.000	880.872
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2006)
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	227.092	0.000	0.000	227.092
Other Land	0.226	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.045	144.538	0.000	144.809
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2007)	2 007.142	0.000	1 425.895	122.082	880.921	0.000	94.000	0.000	227.930	145.550	0.000	4 903.520
Net change	0.202	0.000	-1.462	-0.368	0.049	0.000	0.000	0.000	0.838	0.741	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2007)
Forest Land (managed)	2 006.819	0.000	0.010	0.000	0.119	0.000	0.000	0.000	0.058	0.136	0.000	2 007.142
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.084	0.000	1 420.579	0.000	1.248	0.000	0.000	0.000	2.479	1.505	0.000	1 425.895
Cropland perennial	0.000	0.000	0.310	121.462	0.000	0.000	0.000	0.000	0.000	0.000	0.000	122.082
Grassland (managed)	0.847	0.000	0.772	0.000	878.485	0.000	0.000	0.000	0.711	0.106	0.000	880.921
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	225.811	2.119	0.000	227.930
Other Land	0.507	0.000	0.182	0.000	0.000	0.000	0.000	0.000	0.000	144.861	0.000	145.550
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2008)	2 008.257	0.000	1 421.853	121.772	879.852	0.000	94.000	0.000	229.059	148.727	0.000	4 903.520
Net change	1.115	0.000	-4.042	-0.310	-1.069	0.000	0.000	0.000	1.129	3.177	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2008)
Forest Land (managed)	2 007.795	0.000	0.014	0.000	0.050	0.000	0.000	0.000	0.262	0.136	0.000	2 008.257
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.044	0.000	1 416.273	0.000	1.264	0.000	0.000	0.000	3.371	0.901	0.000	1 421.853
Cropland perennial	0.000	0.000	0.291	121.190	0.000	0.000	0.000	0.000	0.000	0.000	0.000	121.772

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2008)
Grassland (managed)	0.472	0.000	1.244	0.000	877.156	0.000	0.000	0.000	0.550	0.430	0.000	879.852
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	225.206	3.853	0.000	229.059
Other Land	0.532	0.000	0.162	0.000	0.000	0.000	0.000	0.000	0.550	147.483	0.000	148.727
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
<b>Final area (2009)</b>	<b>2 008.843</b>	<b>0.000</b>	<b>1 417.984</b>	<b>121.481</b>	<b>878.470</b>	<b>0.000</b>	<b>94.000</b>	<b>0.000</b>	<b>229.939</b>	<b>152.803</b>	<b>0.000</b>	<b>4 903.520</b>
Net change	0.586	0.000	-3.869	-0.291	-1.382	0.000	0.000	0.000	0.882	4.022	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2009)
Forest Land (managed)	2 008.517	0.000	0.022	0.000	0.156	0.000	0.000	0.000	0.066	0.082	0.000	2 008.843
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.035	0.000	1 415.108	0.000	0.562	0.000	0.000	0.000	1.324	0.955	0.000	1 417.984
Cropland perennial	0.000	0.000	0.308	120.865	0.000	0.000	0.000	0.000	0.000	0.000	0.000	121.481
Grassland (managed)	1.218	0.000	0.778	0.000	875.766	0.000	0.000	0.000	0.524	0.184	0.000	878.470
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	228.150	1.789	0.000	229.939
Other Land	1.479	0.000	0.416	0.000	0.000	0.000	0.000	0.000	0.524	150.384	0.000	152.803
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
<b>Final area (2010)</b>	<b>2 011.249</b>	<b>0.000</b>	<b>1 416.632</b>	<b>121.173</b>	<b>876.484</b>	<b>0.000</b>	<b>94.000</b>	<b>0.000</b>	<b>230.588</b>	<b>153.394</b>	<b>0.000</b>	<b>4 903.520</b>
Net change	2.406	0.000	-1.352	-0.308	-1.986	0.000	0.000	0.000	0.649	0.591	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2010)
Forest Land (managed)	2 011.162	0.000	0.000	0.000	0.013	0.000	0.000	0.000	0.023	0.051	0.000	2 011.249
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.115	0.000	1 414.162	0.000	0.157	0.000	0.000	0.000	0.713	1.485	0.000	1 416.632
Cropland perennial	0.000	0.000	0.238	120.697	0.000	0.000	0.000	0.000	0.000	0.000	0.000	121.173
Grassland (managed)	0.933	0.000	1.073	0.000	874.054	0.000	0.000	0.000	0.424	0.000	0.000	876.484
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	230.588	0.000	0.000	230.588
Other Land	0.126	0.000	0.180	0.000	0.000	0.000	0.000	0.000	0.219	152.869	0.000	153.394
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2011)	2 012.336	0.000	1 415.653	120.935	874.224	0.000	94.000	0.000	231.967	154.405	0.000	4 903.520
Net change	1.087	0.000	-0.979	-0.238	-2.26	0.000	0.000	0.000	1.379	1.011	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2011)
Forest Land (managed)	2 012.214	0.000	0.002	0.000	0.011	0.000	0.000	0.000	0.037	0.072	0.000	2 012.336
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.274	0.000	1 412.856	0.000	0.546	0.000	0.000	0.000	0.725	1.252	0.000	1 415.653
Cropland perennial	0.000	0.000	0.027	120.881	0.000	0.000	0.000	0.000	0.000	0.000	0.000	120.935
Grassland (managed)	1.044	0.000	0.746	0.000	870.767	0.000	0.000	0.000	0.574	1.093	0.000	874.224
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	231.263	0.704	0.000	231.967
Other Land	0.527	0.000	0.108	0.000	0.000	0.000	0.000	0.000	0.000	153.770	0.000	154.405
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2012)	2 014.059	0.000	1 413.739	120.908	871.324	0.000	94.000	0.000	232.599	156.891	0.000	4 903.520



CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2011)
Net change	1.723	0.000	-1.914	-0.027	-2.900	0.000	0.000	0.000	0.632	2.486	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2012)
Forest Land (managed)	2 013.955	0.000	0.006	0.000	0.016	0.000	0.000	0.000	0.036	0.046	0.000	2 014.059
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.057	0.000	1 411.632	0.000	0.258	0.000	0.000	0.000	0.915	0.877	0.000	1 413.739
Cropland perennial	0.000	0.000	0.405	120.098	0.000	0.000	0.000	0.000	0.000	0.000	0.000	120.908
Grassland (managed)	0.800	0.000	0.872	0.000	867.787	0.000	0.000	0.000	0.952	0.913	0.000	871.324
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	231.402	1.197	0.000	232.599
Other Land	0.556	0.000	0.214	0.000	0.000	0.000	0.000	0.000	0.000	156.121	0.000	156.891
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2013)	2 015.368	0.000	1 413.129	120.503	868.061	0.000	94.000	0.000	233.305	159.154	0.000	4 903.520
Net change	1.309	0.000	-0.610	-0.405	-3.263	0.000	0.000	0.000	0.706	2.263	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2013)
Forest Land (managed)	2 015.219	0.000	0.004	0.000	0.052	0.000	0.000	0.000	0.037	0.056	0.000	2 015.368
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.168	0.000	1 411.008	0.000	0.113	0.000	0.000	0.000	0.604	1.236	0.000	1 413.129
Cropland perennial	0.000	0.000	0.372	119.759	0.000	0.000	0.000	0.000	0.000	0.000	0.000	120.503
Grassland (managed)	1.582	0.000	0.675	0.000	864.516	0.000	0.000	0.000	0.420	0.868	0.000	868.061
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2013)
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	233.305	0.000	0.000	233.305
Other Land	0.136	0.000	0.169	0.000	0.000	0.000	0.000	0.000	0.05	158.799	0.000	159.154
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2014)	2 017.105	0.000	1 412.228	120.131	864.681	0.000	94.000	0.000	234.416	160.959	0.000	4 903.520
Net change	1.737	0.000	-0.901	-0.372	-3.380	0.000	0.000	0.000	1.111	1.805	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2014)
Forest Land (managed)	2 016.971	0.000	0.008	0.000	0.006	0.000	0.000	0.000	0.039	0.081	0.000	2 017.105
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.273	0.000	1 409.012	0.000	0.448	0.000	0.000	0.000	0.651	1.844	0.000	1 412.228
Cropland perennial	0.000	0.000	0.409	119.313	0.000	0.000	0.000	0.000	0.000	0.000	0.000	120.131
Grassland (managed)	2.302	0.000	1.299	0.000	858.147	0.000	0.000	0.000	0.407	2.526	0.000	864.681
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	233.414	0.002	0.000	234.416
Other Land	0.57	0.000	0.566	0.000	0.000	0.000	0.000	0.000	0.000	159.823	0.000	160.959
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2015)	2 020.116	0.000	1 411.294	119.722	858.601	0.000	94.000	0.000	235.511	164.276	0.000	4 903.520
Net change	3.011	0.000	-0.934	-0.409	-6.080	0.000	0.000	0.000	1.095	3.317	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2015)
Forest Land (managed)	2 020.055	0.000	0.000	0.000	0.007	0.000	0.000	0.000	0.014	0.040	0.000	2 020.116
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.090	0.000	1 409.400	0.000	0.187	0.000	0.000	0.000	1.045	0.572	0.000	1 411.294
Cropland perennial	0.000	0.000	0.054	119.614	0.000	0.000	0.000	0.000	0.000	0.000	0.000	119.722

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2015)
Grassland (managed)	1.908	0.000	0.179	0.000	855.688	0.000	0.000	0.000	0.327	0.499	0.000	858.601
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	234.895	0.616	0.000	235.511
Other Land	0.469	0.000	0.145	0.000	0.000	0.000	0.000	0.000	0.000	163.662	0.000	164.276
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2016)	2 022.522	0.000	1 409.778	119.668	855.882	0.000	94.000	0.000	236.281	165.389	0.000	4 903.520
Net change	2.406	0.000	-1.516	-0.054	-2.719	0.000	0.000	0.000	0.770	1.113	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2016)
Forest Land (managed)	2 022.396	0.000	0.000	0.000	0.010	0.000	0.000	0.000	0.060	0.056	0.000	2 022.522
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.271	0.000	1 408.090	0.000	0.344	0.000	0.000	0.000	0.497	0.576	0.000	1 409.778
Cropland perennial	0.000	0.000	0.131	119.537	0.000	0.000	0.000	0.000	0.000	0.000	0.000	119.668
Grassland (managed)	1.506	0.000	0.389	0.000	853.403	0.000	0.000	0.000	0.569	0.015	0.000	855.882
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	235.853	0.428	0.000	236.281
Other Land	0.201	0.000	0.050	0.000	0.000	0.000	0.000	0.000	0.000	165.138	0.000	165.389
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
Final area (2017)	2 024.374	0.000	1 408.660	119.537	853.757	0.000	94.000	0.000	236.979	166.213	0.000	4 903.520
Net change	1.852	0.000	-1.118	-0.131	-2.125	0.000	0.000	0.000	0.698	0.824	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2017)
Forest Land (managed)	2 024.125	0.000	0.000	0.000	0.094	0.000	0.000	0.000	0.018	0.137	0.000	2 024.374
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.136	0.000	1 407.487	0.150	0.106	0.000	0.000	0.000	0.557	0.224	0.000	1 408.660
Cropland perennial	0.000	0.000	0.000	119.537	0.000	0.000	0.000	0.000	0.000	0.000	0.000	119.537
Grassland (managed)	1.118	0.000	0.132	0.000	851.485	0.000	0.000	0.000	0.447	0.575	0.000	853.757
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	236.867	0.112	0.000	236.979
Other Land	0.648	0.000	0.110	0.000	0.000	0.000	0.000	0.000	0.000	165.455	0.000	166.213
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2018)	2 026.027	0.000	1 407.729	119.687	851.685	0.000	94.000	0.000	237.889	166.503	0.000	4 903.520
Net change	1.653	0.000	-0.931	0.000	0.094	0.000	0.000	0.000	0.018	0.137	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2018)
Forest Land (managed)	2 025.937	0.000	0.001	0.000	0.026	0.000	0.000	0.000	0.034	0.029	0.000	2 025.937
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.000	0.000	1 406.257	0.026	0.225	0.000	0.000	0.000	0.778	0.443	0.000	0.000
Cropland perennial	0.000	0.000	0.000	119.687	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Grassland (managed)	1.162	0.000	0.121	0.000	850.349	0.000	0.000	0.000	0.053	0.000	0.000	1.162
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	0.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	237.855	0.034	0.000	0.000
Other Land	0.000	0.000	0.020	0.000	0.000	0.000	0.000	0.000	0.000	166.483	0.000	0.000
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2019)	2 027.099	0.000	1 406.399	119.713	850.600	0.000	94.000	0.000	238.720	166.989	0.000	4 903.520

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2018)
Net change	1.072	0.000	-1.330	0.026	-1.085	0.000	0.000	0.000	0.831	0.486	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2019)
Forest Land (managed)	2 026.996	0.000	0.004	0.000	0.009	0.000	0.000	0.000	0.023	0.067	0.000	2 027.099
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.046	0.000	1 405.177	0.022	0.160	0.000	0.000	0.000	0.782	0.212	0.000	1 406.399
Cropland perennial	0.000	0.000	0.000	119.713	0.000	0.000	0.000	0.000	0.000	0.000	0.000	119.713
Grassland (managed)	0.639	0.000	0.024	0.000	849.858	0.000	0.000	0.000	0.051	0.028	0.000	850.600
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	238.591	0.129	0.000	238.720
Other Land	0.171	0.000	0.058	0.000	0.000	0.000	0.000	0.000	0.000	166.760	0.000	166.989
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2020)	2 027.852	0.000	1 405.263	119.735	850.027	0.000	94.000	0.000	239.447	167.196	0.000	4 903.520
Net change	0.753	0.000	-1.136	0.022	-0.573	0.000	0.000	0.000	0.727	0.207	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2020)
Forest Land (managed)	2 027.779	0.000	0.000	0.000	0.016	0.000	0.000	0.000	0.036	0.021	0.000	2 027.852
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.020	0.000	1 404.459	0.000	0.019	0.000	0.000	0.000	0.545	0.220	0.000	1 405.263
Cropland perennial	0.000	0.000	0.024	119.711	0.000	0.000	0.000	0.000	0.000	0.000	0.000	119.735
Grassland (managed)	0.598	0.000	0.037	0.000	849.238	0.000	0.000	0.000	0.154	0.000	0.000	850.027
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2020)
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	239.435	0.012	0.000	239.447
Other Land	0.112	0.000	0.059	0.000	0.000	0.000	0.000	0.000	0.000	167.025	0.000	167.196
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2021)	2 028.509	0.000	1 404.579	119.711	849.273	0.000	94.000	0.000	240.170	167.278	0.000	4 903.520
Net change	0.657	0.000	-0.684	-0.024	-0.754	0.000	0.000	0.000	0.723	0.082	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2021)
Forest Land (managed)	1 998.590	0.000	0.168	0.000	1.263	0.000	0.000	0.000	0.997	1.755	0.000	2 002.773
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	1.899	0.000	1 381.760	0.198	15.641	0.000	0.000	0.000	18.593	15.111	0.000	1 433.202
Cropland perennial	0.000	0.000	3.900	119.391	0.000	0.000	0.000	0.000	0.000	0.000	0.000	123.291
Grassland (managed)	19.834	0.000	15.579	0.000	831.953	0.000	0.000	0.000	7.078	7.413	0.000	881.857
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	211.609	11.746	0.000	223.355
Other Land	8.712	0.000	2.457	0.000	0.031	0.000	0.000	0.000	2.667	131.175	0.000	145.042
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Final area (2022)	2 029.035	0.000	1 403.864	119.589	848.888	0.000	94.000	0.000	240.944	167.200	0.000	4 903.520
Net change	26.262	0.000	-29.338	-3.702	-32.969	0.000	0.000	0.000	17.589	22.158	0.000	

CATEGORY	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2022)
Forest Land (managed)	2 028.971	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.007	0.057	0.000	2 029.035
Forest Land (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Cropland annual	0.020	0.000	1 402.886	0.000	0.041	0.000	0.000	0.000	0.682	0.235	0.000	1 403.864
Cropland perennial	0.000	0.000	0.090	119.499	0.000	0.000	0.000	0.000	0.000	0.000	0.000	119.589

<b>CATEGORY</b>	Forest Land (managed)	Forest Land (unmanaged)	Cropland annual	Cropland perennial	Grassland (managed)	Grassland (unmanaged)	Wetland (managed)	Wetland (unmanaged)	Settlements	Other Land	Total unmanaged land	Initial area (2022)
Grassland (managed)	0.977	0.000	0.008	0.000	847.657	0.000	0.000	0.000	0.246	0.000	0.000	848.888
Grassland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Wetland (managed)	0.000	0.000	0.000	0.000	0.000	0.000	94.000	0.000	0.000	0.000	0.000	94.000
Wetland (unmanaged)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Settlements	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	240.888	0.056	0.000	240.944
Other Land	0.455	0.000	0.020	0.000	0.000	0.000	0.000	0.000	0.000	166.725	0.000	167.200
Total unmanaged land	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
<b>Final area (2023)</b>	<b>2 030.423</b>	<b>0.000</b>	<b>1 403.004</b>	<b>119.499</b>	<b>847.698</b>	<b>0.000</b>	<b>94.000</b>	<b>0.000</b>	<b>241.823</b>	<b>167.073</b>	<b>0.000</b>	<b>4 903.520</b>
Net change	1.388	0.000	-0.860	-0.090	-1.190	0.000	0.000	0.000	0.879	-0.127	4 903.520	

## Annex A6.2. Uncertainty Analyses in the LULUCF Sector

This Annex provides results of the application of Monte Carlo simulations uncertainty analyses in the LULUCF sector. The methodology of calculations of GHG emissions and removals follows the methods described in this Document. If compared to previous submission, analyses of uncertainties of the GHG emissions and removals in the whole LULUCF sector are provided, including deadwood carbon pool in CRT 4A category Forest Land and CRT 4(IV) category Indirect N<sub>2</sub>O emission from managed soils (Nitrogen Leaching and Run-off). These were missing in submission 2022.

In order to apply the Monte Carlo iterated simulations, calculations were automated using the Python programming language. Input data and factors (constant values) were modified for each iteration using the level of uncertainty (if known) according to normal or triangle distribution. [Table A6.2.1](#) shows the levels of uncertainties. The number of iterations was set to 100.000.

Results of the Monte Carlo simulations for the main LULUCF categories and HWP, as well as for the whole LULUCF sector, are shown in [Tables A6.2.2 – A6.2.8](#) and on [Figures A6.2.1 – A6.2.7](#).

**Table A6.2.1: The levels of uncertainty for input data and factors**

LULUCF CATEGORY	DATA / FACTOR	DATA TYPE (D-DEFAULT, N-NATIONAL)	UNCERTAINTY IF KNOWN (%)
4.A.1 Forest Land remaining Forest land – Carbon stock change emissions (Gain-Loss method according to the equation 2.7 of the IPCC 2006 GL. Calculations of carbon stock changes in living biomass following the equations 2.9 - 2.12 of the IPCC 2006 GL)	Area of LULUCF category (and transitions, all categories)	N	3
	Share of tree species	N	15
	Mean yield class of tree species	N	
	Mean age of tree species	N	
	Current annual increment	N	30
	Wood density	N	
	Root-to-shoot	D	30
	Carbon fraction	D	2
	Yield tables	N	25
	Harvested wood volume	N	20
	Growing stock	N	20
	Carbon stock in dead wood and its annual change	N	8.5
	NFI data	N	
4.A.2 Land converted to Forest land – Carbon stock change emissions	Share of tree species on afforested land	N	
	Mean annual increment of living biomass	N	
	Mean annual accumulation of litter	N	
	Mean annual carbon stock change in dead wood	N	8.5
	Mean annual carbon stock change in mineral soil	N	75
4.A Forest Land – Biomass burning	Share of area with burned harvesting residues (from total harvested area)	N	
	Biomass fraction burned on clearing areas	N	
	Combustion factor	D	
	BCEF	N	25
	Emission factors	D	
	Area of forest fires	N	
	Available mass of fuel for combustion (4.A.2)	D	
4.B.1 Cropland remaining cropland	Share of used arable land	N	
	Annual growth rate of perennial woody biomass	N, D	0, 75
	Average biomass stock of perennial crops	N, D	0, 75
	Annual growth rate of perennial woody biomass	D	75




LULUCF CATEGORY	DATA / FACTOR	DATA TYPE (D-DEFAULT, N-NATIONAL)	UNCERTAINTY IF KNOWN (%)
	Annual change of perennial woody biomass	D	0
	Mean values of soil organic carbon stocks	D	
	Relative stock change factor (FLU)	D	9, 50
	Relative stock change factor (FMG)	D	5, 6
	Relative stock change factor (FI)	D	0
Land converted to category (4.B.2, 4.C.2, 4.E.2, 4.F.2)	Mean growing stock	N	20
	Mean dead wood biomass stocks	N	75.24
	Mean carbon stock in litter	N	75.24
	Mean carbon stock in mineral soil	N	75
4.G Harvested Wood products	FAO data (roundwood, other)	D	5, 10
	Carbon content	D	10
	Conversion factors	D	25
	Half-lives	D	50

## LULUCF Categories

**Table A6.2.2: Results of Monte Carlo simulation for category 4.A Forest Land (Gg CO<sub>2</sub> eq.)**

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	-8 243.35	-8 265.46	-8 191.53	2 086.36	-12 591.60	-4 407.27	-52.34	46.68
1995	-8 585.14	-8 612.05	-8 526.09	2 118.50	-13 026.16	-4 700.32	-51.25	45.42
2000	-7 455.80	-7 528.37	-7 445.75	2 237.14	-12 166.05	-3 397.14	-61.60	54.88
2005	-1 831.42	-1 899.23	-1 835.88	2 590.22	-7 178.81	3 006.97	-277.99	258.33
2010	-2 807.93	-2 861.14	-2 787.36	2 622.42	-8 233.59	2 067.48	-187.77	172.26
2015	-3 897.01	-3 959.41	-3 884.20	2 722.23	-9 520.20	1 188.01	-140.45	130.00
2016	-3 685.05	-3 750.43	-3 684.97	2 745.73	-9 345.03	1 441.15	-149.17	138.43
2017	-3 561.81	-3 616.54	-3 549.28	2 755.30	-9 225.28	1 578.52	-155.09	143.65
2018	-2 853.97	-2 903.97	-2 838.61	2 810.87	-8 583.56	2 450.36	-195.58	184.38
2019	-3 787.62	-3 839.84	-3 768.90	2 741.95	-9 436.37	1 340.49	-145.75	134.91
2020	-5 925.67	-5 979.27	-5 893.60	2 545.79	-11 235.17	-1 235.64	-87.90	79.33
2021	-5 768.52	-5 818.83	-5 730.40	2 547.48	-11 069.68	-1 068.08	-90.24	81.64
2022	-6 573.79	-6 639.48	-6 530.39	2 771.12	-12 380.84	-1 489.34	-86.47	77.57


**Figure A6.2.1: Probability distribution function for the category 4.A Forest Land, 2022**



**Table A6.2.3: Results of Monte Carlo simulation for category 4.B Cropland (Gg CO<sub>2</sub> eq.)**

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	-399.36	-399.88	-399.99	348.98	-1 083.54	279.86	-170.97	169.98
1995	-311.62	-311.99	-310.01	331.33	-962.45	333.63	-208.48	206.93
2000	-410.39	-410.55	-410.96	320.67	-1 037.34	214.65	-152.67	152.28
2005	-495.65	-496.78	-496.52	317.90	-1 119.19	120.66	-125.29	124.29
2010	-546.98	-547.38	-546.70	313.09	-1 159.63	59.99	-111.85	110.96
2015	-489.40	-489.65	-488.70	312.10	-1 101.93	113.53	-125.04	123.19
2016	-578.01	-578.17	-577.77	311.42	-1 185.83	26.95	-105.10	104.66
2017	-609.93	-610.40	-610.26	310.77	-1 218.82	-7.13	-99.68	98.83
2018	-616.29	-616.95	-616.51	311.62	-1 224.90	-12.10	-98.54	98.04
2019	-622.07	-624.33	-623.23	309.67	-1 230.76	-21.55	-97.17	96.55
2020	-474.26	-474.51	-474.53	342.30	-1 142.40	188.26	-140.75	139.67
2021	-645.34	-645.47	-644.25	309.37	-1 250.88	-47.21	-93.79	92.69
2022	-641.69	-642.09	-641.41	307.79	-1 245.83	-45.13	-94.03	92.97

**Figure A6.2.2: Probability distribution function for the category 4.B Cropland, 2022**



**Table A6.2.4: Results of Monte Carlo simulation for category 4.C Grassland (Gg CO<sub>2</sub> eq.)**

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	-194.92	-192.69	-192.36	119.10	-427.03	41.51	-121.62	121.54
1995	-256.86	-256.10	-255.78	93.07	-439.37	-255.78	-71.56	71.47
2000	-308.98	-309.46	-309.45	94.86	-495.98	-122.83	-60.28	60.31
2005	-199.80	-200.04	-199.61	116.70	-429.40	29.36	-114.66	114.67
2010	-214.87	-214.92	-214.69	109.08	-429.01	-0.80	-99.62	99.63
2015	-190.43	-190.36	-190.36	74.12	-336.48	-44.89	-76.57	76.44
2016	-178.13	-178.26	-178.09	70.09	-316.22	-40.58	-77.39	77.23
2017	-164.48	-164.58	-164.17	65.67	-293.44	-35.01	-78.40	78.71
2018	-110.54	-110.53	-110.24	61.00	-230.29	9.67	-108.36	108.75
2019	-117.74	-117.74	-117.52	50.66	-217.34	-17.91	-84.59	84.79
2020	-92.54	-92.43	-92.25	38.12	-167.25	-17.23	-80.95	81.36
2021	-55.53	-55.68	-55.87	25.47	-105.89	-5.35	-90.17	90.39
2022	-36.32	36.33	36.30	16.33	-68.54	-4.11	-88.64	88.70

Figure A6.2.3: Probability distribution function for the category 4.C Grassland, 2022





Table A6.2.5. Results of Monte Carlo simulation for category 4.E Settlements (Gg CO<sub>2</sub> eq.)

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	100.40	100.63	100.61	13.47	74.29	127.09	-26.17	26.30
1995	64.73	65.13	65.08	11.57	42.47	87.99	-34.79	34.94
2000	57.22	57.24	57.22	10.16	37.39	77.26	-34.68	34.98
2005	64.66	64.71	64.69	7.49	50.13	79.40	-22.53	22.70
2010	104.85	104.84	104.83	9.54	86.25	123.60	-17.73	17.89
2015	89.18	89.15	89.13	12.02	65.58	112.65	-26.44	26.36
2016	84.06	84.03	84.01	12.48	59.64	108.49	-29.03	29.10
2017	104.80	104.77	104.77	13.47	78.49	131.23	-25.09	25.25
2018	85.56	85.52	85.50	13.97	58.15	112.97	-32.00	32.09
2019	88.29	88.25	88.23	14.21	60.40	116.22	-31.56	31.68
2020	83.37	83.63	83.61	14.15	55.93	111.44	-33.12	33.26
2021	89.25	89.26	89.25	14.47	61.02	117.80	-31.63	31.98
2022	83.18	83.20	83.20	14.57	54.80	111.78	-34.13	34.35


Figure A6.2.4: Probability distribution function for the category 4.E Settlements, 2022



**Table A6.2.6: Results of Monte Carlo simulation for category 4.F Other Land (Gg CO<sub>2</sub> eq.)**

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	301.65	304.83	304.66	31.39	243.74	366.48	-20.04	20.23
1995	110.67	111.45	111.36	22.15	68.30	155.05	-38.71	39.12
2000	110.86	110.88	110.83	17.67	76.17	145.75	-31.31	31.44
2005	190.18	190.13	189.97	17.88	155.60	225.59	-18.16	18.65
2010	92.77	92.78	92.77	8.66	75.93	109.85	-18.16	18.40
2015	189.92	189.90	189.86	14.66	161.33	218.69	-15.05	15.16
2016	103.73	103.73	103.77	14.46	75.47	132.11	-27.24	27.36
2017	99.84	99.84	99.85	14.62	71.26	128.63	-28.62	28.83
2018	148.56	148.57	148.52	16.22	116.80	180.44	-21.38	21.45
2019	85.11	85.11	85.14	14.90	56.00	114.28	-34.20	34.28
2020	99.26	100.10	100.07	15.02	70.72	129.51	-29.35	29.39
2021	75.93	75.93	75.87	14.63	47.46	104.64	-37.50	37.82
2022	79.37	79.45	79.43	14.57	50.87	108.06	-35.97	36.00

**Figure A6.2.5: Probability distribution function for the category 4.F Other Land, 2022**



**Table A6.2.7: Results of Monte Carlo simulation for category 4.G HWP (Gg CO<sub>2</sub> eq.)**

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	-470.41	-354.88	-394.37	275.81	-708.00	226.16	-99.50	163.73
1995	-58.77	28.54	-13.56	282.06	-311.70	595.40	-1 192.09	1 986.06
2000	-920.07	-811.04	-856.62	340.35	-1 279.98	-80.34	-57.82	90.09
2005	-1 996.46	-1 874.87	-1 923.86	407.15	-2 440.37	-1 033.88	-30.16	44.86
2010	-1 334.60	-1 197.06	-1 255.53	445.43	-1 802.39	-260.02	-50.57	78.28
2015	-940.70	-798.31	-861.87	445.43	-1 379.40	139.95	-72.79	117.53
2016	-1 063.66	-920.35	-983.03	467.04	-1 518.13	29.29	-64.95	103.18
2017	-1 077.04	-1 002.96	-1 044.34	302.68	-1 458.70	-298.10	-45.44	70.28

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
2018	-889.19	-818.78	-859.45	292.39	-1 264.15	-132.62	-54.39	83.80
2019	-644.91	-578.34	-617.60	282.21	-1 009.64	86.18	-74.57	114.90
2020	-247.28	-182.93	-219.02	272.48	-602.74	457.66	-229.48	350.18
2021	-382.71	-248.80	-283.15	264.92	-660.92	371.56	-165.65	249.34
2022	-144.36	4.763	-62.44	473.89	-529.20	969.90	-12 532	20 262

Figure A6.2.6: Probability distribution function for the category 4.G HWP, 2022





Table A6.2.8: Results of Monte Carlo simulation for LULUCF sector (Gg CO<sub>2</sub> eq.)

YEAR	NIR 2023	RESULTS OF MONTE CARLO SIMULATIONS						
		Average	Median	Standard deviation	2.5%	97.5%	Percentile 2.5	Percentile 97.5
		Gg CO <sub>2</sub> eq.					%	
1990	-8 892.53	-8 802.10	-8 733.80	2 139.78	-13 207.45	-4 810.37	-50.05	45.35
1995	-9 030.43	-8 973.09	-8 893.42	2 230.88	-13 592.81	-4 830.51	-51.48	46.17
2000	-8 922.53	-8 879.53	-8 802.73	2 353.24	-13 702.57	-4 494.53	-54.32	49.38
2005	-4 264.77	-4 199.74	-4 167.69	2 868.42	-9 912.75	1 343.60	-136.03	131.99
2010	-4 704.36	-4 620.48	-4 546.54	2 685.70	-10 112.91	441.94	-118.87	109.56
2015	-5 234.27	-5 154.70	-5 084.65	2 780.10	-10 808.11	114.55	-109.67	102.22
2016	-5 312.70	-5 234.99	-5 173.91	2 804.15	-10 895.04	68.43	-108.12	101.31
2017	-5 204.24	-5 112.08	-5 046.63	2 820.26	-10 824.08	220.20	-111.74	104.31
2018	-4 231.31	-4 135.03	-4 076.84	2 867.62	-9 909.92	1 330.26	-139.66	132.17
2019	-4 994.47	-4 900.12	-4 830.12	2 800.23	-10 585.86	399.58	-116.03	108.15
2020	-6 552.69	-6 458.00	-6 388.46	2 609.11	-11 805.60	-1 571.39	-82.81	75.67
2021	-6 682.58	-6 582.07	-6 502.14	2 611.91	-11 917.01	-1 694.48	-81.05	74.26
2022	-7 229.29	-7 146.17	-7 051.54	2 827.83	-12 958.24	-1 874.43	-81.33	73.77

Figure A6.2.7: Probability distribution function for LULUCF sector, 2022



---

<b>CHAPTER 7. WASTE (CRT 5)</b> .....	<b>384</b>
7.1. OVERVIEW OF THE WASTE SECTOR.....	384
7.2. CATEGORY-SPECIFIC QA/QC AND VERIFICATION PROCESS.....	388
7.3. CATEGORY-SPECIFIC RECALCULATIONS.....	390
7.4. CATEGORY-SPECIFIC IMPROVEMENTS AND IMPLEMENTATION OF RECOMMENDATIONS.....	394
7.5. SOLID WASTE DISPOSAL (CRF 5.A).....	395
7.5.1. <i>Municipal Waste Disposal Sites (Managed)</i> .....	396
7.5.2. <i>Non-municipal Disposal Sites (Industrial)</i> .....	401
7.6. BIOLOGICAL TREATMENT OF SOLID WASTE (CRT 5.B) .....	404
7.6.1. <i>Composting (CRT 5.B.1)</i> .....	404
7.6.2. <i>Methodological Issues</i> .....	405
7.6.3. <i>Anaerobic Digestion at Biogas Facilities (CRT 5.B.2)</i> .....	405
7.6.4. <i>Uncertainties and Time-series Consistency</i> .....	406
7.6.5. <i>Category-specific Recalculations</i> .....	407
7.7. WASTE INCINERATION AND OPEN BURNING OF WASTE (CRT 5.C) .....	408
7.7.1. <i>Waste Incineration (CRT 5.C.1)</i> .....	408
7.7.2. <i>Open Burning of Waste (CRT 5.C.2)</i> .....	412
7.8. WASTEWATER TREATMENT AND DISCHARGE (CRT 5.D).....	412
7.8.1. <i>Domestic Wastewater (CRF 5.D.1)</i> .....	415
7.8.2. <i>Industrial Wastewater (CRF 5.D.2)</i> .....	420
7.9. MEMO ITEMS (CRT 5.F) .....	424

## CHAPTER 7. WASTE (CRT 5)

This Chapter was prepared using GWP<sub>100</sub> taken from the 5<sup>th</sup> Assessment Report of the IPCC by the sectoral experts and institutions involved in the National Inventory System of the Slovak Republic:

INSTITUTE	CHAPTER	SECTORAL EXPERT
External expert of the SHMÚ	Category 5.A, 5.F	Marek Hrabčák
Slovak Hydrometeorological Institute (SHMÚ)	Category 5.B	Kristína Tonhauzer
Slovak Hydrometeorological Institute (SHMÚ)	Category 5.C	Zuzana Jonáček
FCHPT STU Bratislava	Category 5.D	Igor Bođík
Slovak Hydrometeorological Institute (SHMÚ)	Chapter 7.7.2 (partly)	Lea Mrafková

### 7.1. OVERVIEW OF THE WASTE SECTOR

Inventory of the Waste sector includes direct (CH<sub>4</sub>, CO<sub>2</sub>, N<sub>2</sub>O) and indirect GHG emissions (NMVOCs). Methane is generated from solid waste disposal sites, biological treatment of waste, waste incineration and wastewater treatment. The main source of CO<sub>2</sub> emissions is waste incineration. N<sub>2</sub>O emissions are generated from the biological treatment of waste and from wastewater treatment. Estimation of the following emission categories in 2025 submission is presented in this chapter:

- 5.A Solid waste disposal;
- 5.B Biological treatment of solid waste;
- 5.C Incineration and open burning of waste;
- 5.D Wastewater treatment and discharge;
- 5.F Memo Items (HWP).

In 2023, total aggregated GHG net emissions from the Waste sector are relatively stable over the entire period 1990 – 2023 as is shown on [Figure 7.1](#). Total aggregated emissions from the Waste sector were 1 671.86 Gg of CO<sub>2</sub> eq. in 2023 and they decreased by 4% compared to the previous year, due to a main decrease of the amount of SWDS category responsible for a decrease in methane emissions. Compared to the reference year 1990, total GHG emissions increased by 41%. The increase of emissions in SWDS and biological treatment was compensated by the decrease of emissions from incineration of waste without energy use and Wastewater. Emissions from waste incineration with energy use were allocated into the Energy sector (1.A.1.a – Other Fuels for municipal waste and 1.A.2.c&1.A.2.f for industrial waste incineration).

Emissions from landfilled waste (5.A) have changed their current trend and decreased on the level of 2010, following the decreasing trend in last years. The emissions growth from waste disposal slowed down after 2011 and peaked in 2018, since then was already a decrease in time series (albeit minimal) recognised. New methane emissions from landfilled waste in 2023 are slightly lower than in 2022 by -3%. Emissions from industrial landfilled waste (ISW) have been steadily declining since 2008 due to recycling strategy.

Emissions from biological treatment (5.B) do not vary significantly, but there is a decrease in the last year 2023 due to decreasing amounts of waste sent for composting by more than 11%.



Emissions from waste incineration without energy recovery (5.C) were recalculated due to reconsideration of the methodological approach and new activity data. The significant decrease in emissions was due to the new operational hours of heat exchange facilities in one facility that use waste to generate energy (closed in 2022). The waste was therefore incinerated with energy recovery, again.

Total emissions from the category 5.D were gradually decreasing since 1990, but in the last 5-7 years a slight increase in total emissions from N<sub>2</sub>O emissions from nitrogen removal processes on WWTPs there has been occurred.

**Figure 7.1:** Trend of Waste sector emissions by categories in 1990 – 2023





Figure 7.2 below shows that the most important source of GHG emissions is solid waste disposal (69.5%), followed by biological treatment (18.7%) and wastewater treatment (11.2%) and incineration of waste without energy recovery (0.6%). The Waste sector contributed 4.6% to total GHG emissions in 2023.

**Figure 7.2:** The share of categories in waste sector in 2023



The majority of GHG emissions from the Waste sector are in form of CH<sub>4</sub> with 88% share followed by 12% of N<sub>2</sub>O and 0.14% of CO<sub>2</sub> as shows in Table 7.1 and on Figure 7.3.


Summary of the GHG emissions inventory expressed in GWP taken from the AR5 is visible in the Table 7.1.

**Table 7.1: GHG emissions in the Waste sector according to the gases and categories in particular years**

YEAR	TOTAL CO <sub>2</sub> *	TOTAL CH <sub>4</sub>	TOTAL N <sub>2</sub> O	GHG	TOTAL 5.A	TOTAL 5.B	TOTAL 5.C	TOTAL 5.D
	Gg of CO <sub>2</sub> eq.			Gg of CO <sub>2</sub> eq.				
1990	6.68	1 061.18	116.80	<b>1 184.66</b>	781.78	113.98	31.95	256.96
1995	6.69	1 016.81	100.97	<b>1 124.46</b>	751.87	116.69	31.73	224.17
2000	7.21	1 125.48	96.67	<b>1 229.36</b>	865.28	116.85	34.41	212.83
2005	17.47	1 283.32	105.33	<b>1 406.11</b>	1 021.21	126.26	63.41	195.23
2010	10.43	1 419.50	115.42	<b>1 545.35</b>	1 172.08	143.15	42.59	187.54
2011	8.46	1 466.90	120.19	<b>1 595.55</b>	1 214.42	163.05	37.31	180.78
2012	6.80	1 498.64	114.21	<b>1 619.65</b>	1 239.76	186.33	29.42	164.14
2013	8.23	1 516.83	113.08	<b>1 638.14</b>	1 256.53	182.39	31.20	168.02
2014	6.81	1 513.91	108.20	<b>1 628.92</b>	1 244.94	209.60	27.83	146.56
2015	9.68	1 578.83	147.10	<b>1 735.61</b>	1 263.61	254.84	39.22	177.93
2016	2.58	1 534.65	135.90	<b>1 673.14</b>	1 263.67	218.78	13.34	177.35
2017	2.20	1 553.56	153.88	<b>1 709.64</b>	1 264.20	248.20	12.12	185.11
2018	5.40	1 567.85	155.19	<b>1 728.43</b>	1 264.98	258.29	23.88	181.28
2019	5.28	1 546.20	160.04	<b>1 711.52</b>	1 255.69	243.26	23.11	189.47
2020	4.22	1 568.65	186.89	<b>1 759.76</b>	1 243.93	305.41	19.22	191.19
2021	2.26	1 548.77	178.29	<b>1 729.32</b>	1 238.16	297.00	12.15	182.01
2022	3.31	1 537.32	202.05	<b>1 742.68</b>	1 199.72	346.01	16.04	180.92
2023	2.38	1 472.24	197.24	<b>1 671.86</b>	1 162.62	312.49	10.03	186.72

\*Only non-bio CO<sub>2</sub> included in category 5.C

**Figure 7.3: Trend in aggregated emissions by gases within the waste in 1990 – 2023**



The general approach to estimate emissions in the Waste sector is to use the default parameters taken from the IPCC 2006 GL and country-specific data. Overview of used tiers by category is summarised in **Table 7.2**.

**Table 7.2: Overview of tiers used in the Waste sector in 2023**

EMISSION CATEGORY	GAS/TIER USED		NOTE (RESPONSES TO DECISION TREE)
5.A Solid Waste Disposal	CH <sub>4</sub>	T2/CS	Good quality CS AD are available, except of composition of waste landfilled.
			CS models and parameters partly available.
5.B Biological Treatment	CH <sub>4</sub> , N <sub>2</sub> O	T1/D	CS data on waste available.
			CS emission factors not available.
	CO <sub>2</sub>	T2/CS, D	Plant specific data not available.

EMISSION CATEGORY	GAS/TIER USED		NOTE (RESPONSES TO DECISION TREE)
5.C Incineration and Open Burning			CS data on waste available.
			CS emission factors not available.
5.C Incineration and Open Burning	CH <sub>4</sub> , N <sub>2</sub> O	T2/CS, D	Plant specific data not available.
			CS data on waste available.
5.D Wastewater	CH <sub>4</sub> , N <sub>2</sub> O	T1, T2/D	Wastewater treatment pathways characterised.
			Measurements are available (BOD, COD, N <sub>tot</sub> ), but CS method not available.
			CS emission factors not available, but CS model developed.
			Wastewater is a key category.

European Waste Catalogue (EWC) – the division of waste to the Waste Groups List defined in the European System of Waste Classification (Commission Decision 2000/532/EC) was used for estimating of the emissions. The “municipal solid waste” (MSW) means all waste reported in the Waste Group 20. All the other waste types from Waste Groups 1 – 19 are called “industrial solid waste” (ISW). Statistical data on waste generation, disposal, incineration and recovery by waste groups are published by the ŠÚ SR annually in publication “Odpady v Slovenskej republike” (*Waste in the Slovak Republic*). This is primary source of activity data for estimation of emissions in the Waste sector. **Table 7.3** presents overview of the mass flows in percent for the different waste types in 2023, from generation to the different treatment options, including recycling and landfilling.

**Table 7.3: Overview of generated waste and mass flows for the different waste types according to the national statistics in 2023**

CATEGORY	WASTE TOTAL	RECOVERY, REUSE				DISPOSAL			STORAGE
		A	B	C	D	E	F	G	
	tons	share				share			
<b>SR Total</b>	<b>13 566 348</b>	<b>38.68%</b>	<b>2.96%</b>	<b>11.45%</b>	<b>0.69%</b>	<b>18.52%</b>	<b>0.07%</b>	<b>2.37%</b>	<b>25.26%</b>
01 Wastes resulting from exploration, mining, quarrying, physical and chemical treatment of minerals	114 432	63%	0%	1%	0%	18%	0%	0%	18%
02 Wastes from agriculture, horticulture, aquaculture, forestry, hunting and fishing, food preparation and processing	588 008	44%	3%	25%	1%	1%	0%	6%	21%
03 Wastes from wood processing and the production of panels and furniture, pulp, paper and cardboard	558 487	2%	3%	34%	4%	8%	0%	0%	47%
04 Wastes from the leather, fur and textile industries	16 948	12%	2%	13%	0%	12%	0%	0%	60%
05 Wastes from petroleum refining, natural gas purification and pyrolytic treatment of coal	2 741	9%	0%	0%	0%	1%	9%	24%	58%
06 Wastes from inorganic chemical processes	3 758	18%	0%	40%	0%	2%	0%	14%	26%
07 Wastes from organic chemical processes	65 682	15%	2%	23%	2%	8%	0%	1%	50%
08 Wastes from the manufacture, formulation, supply and use (MFSU) of coatings (paints, varnishes and vitreous enamels),	15 885	4%	1%	0%	0%	8%	0%	17%	70%

CATEGORY	WASTE TOTAL	RECOVERY, REUSE				DISPOSAL			STORAGE
		A	B	C	D	E	F	G	H
	<i>tons</i>	<i>share</i>				<i>share</i>			
adhesives, sealants and printing inks									
09 Wastes from the photographic industry	143	8%	0%	0%	0%	0%	0%	23%	68%
10 Wastes from thermal processes	1 234 642	7%	0%	0%	0%	69%	0%	6%	16%
11 Wastes from chemical surface treatment and coating of metals and other materials; non-ferrous hydro-metallurgy	38 894	20%	0%	4%	0%	2%	0%	40%	33%
12 Wastes from shaping and physical and mechanical surface treatment of metals and plastics	765 360	95%	0%	0%	0%	0%	0%	1%	3%
13 Oil wastes and wastes of liquid fuels (except edible oils, 05 and 12)	38 328	34%	0%	0%	0%	0%	0%	25%	39%
14 Waste organic solvents, refrigerants and propellants (except 07 and 08)	2 630	40%	1%	0%	0%	0%	0%	13%	45%
15 Waste packaging; absorbents, wiping cloths, filter materials and protective clothing not otherwise specified	229 940	10%	1%	57%	0%	7%	0%	1%	23%
16 Wastes not otherwise specified in the list	329 655	55%	0%	4%	0%	7%	0%	23%	11%
17 Construction and demolition wastes (including excavated soil from contaminated sites)	5 153 080	50%	0%	0%	1%	5%	0%	0%	45%
18 Wastes from human or animal health care and/or related research (except kitchen and restaurant wastes not arising from immediate health care)	13 868	7%	12%	1%	0%	3%	27%	3%	46%
19 Wastes from waste management facilities, off-site wastewater treatment plants and the preparation of water intended for human consumption and water for industrial use	1 698 231	42%	9%	13%	0%	17%	0%	4%	15%
20 Municipal waste (household waste and similar commercial, industrial and institutional wastes) including separately collected fractions	2 560 971	19%	8%	31%	1%	39%	0%	0%	2%

A=material, B=energy, C=compost, D=other, E=landfilling, F=incineration, G=other

## 7.2. CATEGORY-SPECIFIC QA/QC AND VERIFICATION PROCESS

QA/QC procedures in the Waste sector are linked with the QA/QC plans for the National Inventory System at the sectoral level and follow basic rules and activities of QA/QC as defined in IPCC 2006 GL. The QC checks (e.g. consistency check between CRT data and national statistics) were done during the CRT and NID compilation, General QC questionnaire was filled and archived by QA/QC manager.

---

Due to larger revisions and recalculations provided currently in the categories 5.A – Solid Waste Disposal Sites and 5.D – Wastewater Treatment, implementation process was finalised on national level followed EU voluntary review of the inventory submitted in 2024. Presentation of new methodology and resulting emissions from the municipal and industrial solid waste disposal sites followed by discussion introduced several interesting areas for further improvements, however the principles and results of the recalculation were accepted on national level.

Verification of activity data used for estimation of emissions from municipal solid waste disposed to SWDS was performed by comparing reported year data to previous years' data. Verification on MSW data was strengthened by correlation with the direct data from disposal sites operators. The data provided by the MŽP SR was verified by the official statistics and implemented into inventory.

The period 1950 – 1990 was estimated based on economic growth according to the procedure given in the previous submissions. For the period 1990 – 2004, statistical input data on waste production was available, however the EWC was not adopted until 2001, thus the groups and types of waste from this period are not entirely consistent with the EWC. Therefore, the data on the composition of waste for the period 1990 – 2004 are extrapolated. For the period from 2005 – 2023, summary statistical data on waste production and composition were used according to real data from disposals' operators collected by the MŽP SR. Data were further analysed up to the level of individual types of waste according to the EWC as maintained in the Information System Waste (IS Waste). These data are sufficiently reliable and valid. In addition, new data source for several parameters (composition of waste) was introduced into inventory for the first time on national level.

In the retrospective review, inventory is relied on the period since sufficiently reliable statistical data on the waste production and management (2005 – 2023) is available. Details on the recalculations and revisions of landfill data since the previous submission are given in [Chapter 7.5.1](#).

Verification of data on recovered methane from landfill gas is ensured by the use of national database of electricity produced from renewable sources, annually published by the Regulatory Office for Network Industries ([Chapter 7.5.1](#)). Verification of activity data used for estimation of emissions from agricultural and industrial solid waste disposed to SWDS was performed by comparing reported year data to previous years' data.

Verification of data on biological treatment was done by comparing data from the ŠÚ SR with the National Strategy of Biodegradable Waste Management provided by the Ministry of Environment of the Slovak Republic (MŽP SR).

Verification of activity data and estimated emissions from MSW incinerators is ensured by comparing results of modelling with the Reports on Operation and Monitoring of Waste incinerators and data reported to the NEIS database and the Annual Reports from companies OLO Bratislava and KOSIT Košice. Verification of activity data and estimated emissions from the non-MSW incinerators is ensured by a modelling results comparison with the information provided in the Reports on Operation and Monitoring of Waste incinerators and the NEIS database and the Annual Reports from companies incinerating and co-incinerating waste. Activity data are available from the Statistical Yearbook and the NEIS database for the waste incineration. Default emission factors were used, and these were verified to fully comply with the IPCC 2019 Refinements to the IPCC 2006 Guidelines. Because the Slovak incinerators do not monitor dry matter content, parameters for wet weight were used consistently for all calculations.

Data on population were obtained from the demographic information updated by the ŠÚ SR, from the Report on Water Management prepared by the Water Research Institute (VÚVH) and from the national censuses. Data on protein consumption are published annually by the ŠÚ SR, however by December 2023, actual data for 2022 is missing. Therefore, the protein consumption for the year 2022 was provided

based on extrapolated real data from the ŠÚ SR for the last 5 years. Sewage sludge data were obtained from the Report on Water Management prepared by the VÚVH.

Data on use of retention tanks (cesspools and septic tanks) are based on population censuses done in years 1991, 2001 and 2011. These censuses are also used for verification of population distribution to individual wastewater pathways. Additional information used in wastewater estimation was collected by the SHMÚ. Data published in statistical reports are verified by a comparison in category and time series. Data on population connected to cesspools and septic tanks, domestic WWTPs as well as others are estimated according discussion with wastewater experts on Slovak University of Technology Bratislava, Association of wastewater treatment experts of Slovak Republic, VÚVH Bratislava a Ministry of Environment SR.

Data on BOD<sub>5</sub>, COD and N<sub>tot</sub> in influents as well as effluents from all Slovak WWTPs was obtained based on information provided by the ŠÚ SR and from the SHMÚ. Additional information used in wastewater estimation was collected by the SHMÚ and the wastewater treatment experts. Data published in statistical reports are verified by a comparison in category and time series.

Information about industrial wastewater is also registered in the Database of Wastewaters at the SHMÚ (Department of Water Quality) and is published by the ŠÚ SR.

### 7.3. CATEGORY-SPECIFIC RECALCULATIONS

Sectoral experts made some smaller revisions of the methodological approaches and used activity data also in 2024 submission. After analysis, several improvements introduced in this submission led to recalculation or reallocation of data from several categories. This recalculation work is reaction on the implementation of the new ETF system and connected with the implementation of the 2019 IPCC Refinements.

In addition, waste composting was prepared by the sectoral expert for agriculture with the cross-checked of data provided between the Agriculture and Waste sectors. The air pollution expert with the cooperation of the energy sectoral expert prepared inventory in the waste incineration category (without energy use). The crosscheck was done between the Energy and Waste sectors in this submission.

In line with the Improvement and Prioritization Plan for 2024, minor correction of data (waste incineration) took place in this submission. These reflecting recommendations made during previous reviews and suggested experts' improvements.

**Table 7.4:** Description of recalculations implemented in 2025 submission

RECOMMENDATION NO.	CATEGORY	DESCRIPTION	REFERENCE
1.	5.A	<p><b>2011-2022:</b> Recalculation based on revision MSW composition (% share of paper + garden + food) based on consideration of recycling share.</p> <p><b>2022:</b> Recovery disposal gas was wrongly reported by the external organisation for 2022 (ÚRSO), data was corrected and lowered.</p> <p><b>2010-2019:</b> Activity data for waste disposal was updated by the Statistical Office of the Slovak Republic, minor changes.</p> <p><b>2020-2022:</b> Activity data for waste disposal was updated according real data from the disposal companies, approved by MŽP SR, minor changes 5-10%.</p>	Chapter 7.5
2.	5.B	This recalculation is connected with the correction of activity data of digestion in <b>2001 – 2022</b> . The revision of new data is connected with data refinement provided by the Statistical Office of the Slovak Republic.	Chapter 7.6
3.	5.C.1.a Biogenic and	Emissions of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O from ISW incineration were recalculated for all-time series <b>1990 – 2022</b> due	Chapter 7.7

RECOMMENDATION NO.	CATEGORY	DESCRIPTION	REFERENCE
	5.C.1.b Non-Biogenic	inclusion of the waste incinerated in the clinical waste incinerators These recalculations increased biogenic as well as non-biogenic GHG emissions in equivalents.	
4.	5.D.1	Recalculations of methane emissions based on the implementation of different MCFs used for methane emission in individual retention tanks - cesspools. Explained different methane production in septic tanks and cesspools.	Chapter 7.8
5.	5.F	Recalculations are connected with recalculations in 5.A category for SWDS and parameters.	Chapter 7.9

**Ad. 1:** This recalculation is connected with the correction of activity data of annual waste disposal on the SWDS and correction of the composition of the waste in 2011 – 2022. The revision of new data is connected with data refinement provided by the Statistical Office of the Slovak Republic and new evidence system of real data on waste amount and composition directly from the disposal sites operators. **Table 7.5** is showing changes led to increase or decrease of emissions in this category.

**Table 7.5: Recalculations of the category 5.A for 2010 – 2022 and comparison of the submissions**

YEAR	5.A.1.a – ANNUAL WASTE AT THE SWDS			5.A.1.a – CH <sub>4</sub> EMISSIONS		
	kt			Gg		
	2025	2024	%	2025	2024	%
2010	1 620.73	1 620.73	100.00%	41.86	41.865	99.99%
2011	1 577.26	1 577.26	100.00%	43.372	43.508	99.69%
2012	1 488.80	1 488.80	100.00%	44.277	44.593	99.29%
2013	1 419.77	1 419.77	100.00%	44.876	45.382	98.89%
2014	1 358.48	1 358.48	100.00%	44.462	45.080	98.63%
2015	1 457.46	1 461.23	99.74%	45.129	45.785	98.57%
2016	1 433.30	1 437.94	99.68%	45.131	45.819	98.50%
2017	1 426.78	1 431.32	99.68%	45.150	46.053	98.04%
2018	1 350.73	1 356.96	99.54%	45.178	46.230	97.72%
2019	1 292.73	1 297.06	99.67%	44.846	45.998	97.50%
2020	1 279.21	1 318.50	97.02%	44.426	45.590	97.45%
2021	1 136.31	1 171.69	96.98%	44.220	45.626	96.92%
2022	1 116.89	1 091.50	102.33%	42.847	43.110	99.39%

**Ad 2:** This recalculation is connected with the correction of activity data on digestion of other waste in years 2001 – 2022. The revision was made due to necessary correction of activity data. For the years 2021 to 2022, the activity data on biogas production were taken from the National Emissions Information System (NEIS – see Energy chapter). These data cover all biogas plants in Slovakia. **Table 7.6** is showing changes led to increase or decrease of emissions in this category.

**Table 7.6: Recalculations in the category 5.B.1.a for 2001 – 2022 and comparison of the submissions**

YEAR	5.B.2 Anaerobic digestion at biogas facilities			5.B.2 – CH <sub>4</sub> EMISSIONS		
	kt dm			kt		
	2024	2025	%	2024	2025	%
2001	56.21	73.87	31.4%	0.04	0.06	31.4%
2002	73.87	73.61	-0.4%	0.06	0.06	-0.4%
2003	73.61	77.87	5.8%	0.06	0.06	5.8%
2004	77.87	85.75	10.1%	0.06	0.07	10.1%
2005	85.75	91.34	6.5%	0.07	0.07	6.5%
2006	91.34	99.86	9.3%	0.07	0.08	9.3%
2007	99.86	117.14	17.3%	0.08	0.09	17.3%

YEAR	5.B.2 Anaerobic digestion at biogas facilities			5.B.2 – CH <sub>4</sub> EMISSIONS		
	kt dm			kt		
	2024	2025	%	2024	2025	%
2008	117.14	132.95	13.5%	0.09	0.11	13.5%
2009	132.95	141.66	6.5%	0.11	0.11	6.5%
2011	141.66	219.67	55.1%	0.11	0.18	55.1%
2011	219.67	399.09	81.7%	0.18	0.32	81.7%
2012	399.09	749.98	87.9%	0.32	0.60	87.9%
2013	749.98	1 372.03	82.9%	0.60	1.10	82.9%
2014	1372.03	1 796.88	31.0%	1.10	1.44	31.0%
2015	1796.88	1 795.72	-0.1%	1.44	1.44	-0.1%
2016	1795.72	1 829.76	1.9%	1.44	1.46	1.9%
2017	1829.76	1 901.41	3.9%	1.46	1.52	3.9%
2018	1901.41	1 774.35	-6.7%	1.52	1.42	-6.7%
2019	1774.35	1 628.87	-8.2%	1.42	1.30	-8.2%
2020	1628.87	1 529.10	-6.1%	1.30	1.22	-6.1%
2021	1529.10	1 410.77	-7.7%	1.22	1.13	-7.7%
2022	1523.38	1 063.59	-30.2%	1.22	0.85	-30.2%

Ad. 3: Emissions of all GHG for the category Waste Incineration – Industrial waste were recalculated in this submission due to inclusion of the waste incinerated in the clinical waste incinerators. Therefore, GHG emissions increased in biogenic and non-biogenic categories of ISW incineration in time series. Revised data on GHG emissions and comparison is provided in the [Table 7.7](#).

**Table 7.7: Recalculations of the category 5.C for 1990 – 2022 and comparison of the submissions**

YEAR	5.C.1.a - GHG EMISSIONS - biogenic			5.C.1.b - GHG EMISSIONS – non-biogenic		
	Gg CO <sub>2</sub> eq.			Gg CO <sub>2</sub> eq.		
	2025	2024	%	2025	2024	%
1990	9.1470	3.7855	241.63%	25.2257	15.4933	162.82%
1991	9.1377	3.7814	241.65%	25.1579	15.4450	162.89%
1992	9.1018	3.7633	241.86%	25.1461	15.4513	162.74%
1993	9.1765	3.7969	241.68%	25.2295	15.4820	162.96%
1994	9.1752	3.7958	241.72%	25.2442	15.4942	162.93%
1995	9.0083	3.7170	242.35%	25.1212	15.4740	162.34%
1996	9.1235	3.7656	242.28%	25.4182	15.6529	162.39%
1997	9.3635	3.8801	241.32%	25.5863	15.6755	163.22%
1998	8.7850	3.6039	243.76%	25.2709	15.6926	161.04%
1999	8.7671	3.5885	244.31%	25.5849	15.9451	160.46%
2000	9.8249	4.0629	241.82%	27.1869	16.7138	162.66%
2001	9.1249	4.1282	221.04%	28.5625	19.1684	149.01%
2002	13.0402	9.0870	143.50%	85.0737	75.3543	112.90%
2003	11.4024	7.2883	156.45%	55.5212	46.5317	119.32%
2004	14.9643	9.8715	151.59%	74.9426	63.7338	117.59%
2005	11.4272	7.6895	148.61%	55.5772	47.4427	117.15%
2006	12.5982	10.7298	117.41%	72.4926	68.3212	106.11%



YEAR	5.C.1.a - GHG EMISSIONS - biogenic			5.C.1.b - GHG EMISSIONS – non-biogenic		
	Gg CO <sub>2</sub> eq.			Gg CO <sub>2</sub> eq.		
	2025	2024	%	2025	2024	%
2007	6.6800	4.1810	159.77%	25.1183	20.2398	124.10%
2008	9.4865	7.1102	133.42%	42.2045	37.4116	112.81%
2009	7.6063	5.3373	142.51%	29.0994	24.7473	117.59%
2010	9.0079	6.5253	138.05%	36.1776	31.3040	115.57%
2011	9.2341	6.0606	152.36%	30.6278	24.7023	123.99%
2012	6.6081	4.9003	134.85%	24.6386	21.4245	115.00%
2013	6.0339	4.5184	133.54%	27.0041	23.8628	113.16%
2014	6.0684	4.4387	136.72%	23.5155	20.3229	115.71%
2015	8.3031	7.0024	118.58%	33.3298	30.7953	108.23%
2016	4.4109	3.5348	124.79%	10.0604	8.5721	117.36%
2017	4.0945	3.6004	113.72%	9.0270	8.2055	110.01%
2018	5.9001	5.1505	114.55%	19.6067	18.2355	107.52%
2019	5.4798	5.0386	108.76%	19.1388	18.3267	104.43%
2020	4.9153	4.4149	111.33%	15.6326	14.7326	106.11%
2021	4.0873	3.5281	115.85%	9.0819	8.1467	111.48%
2022	4.7135	3.9869	118.23%	12.5647	11.2986	111.21%

**Ad 4:** The recalculations were based on the different MCFs used for cesspools. Our previous year calculations for CH<sub>4</sub> production used an MCF value = 0.4 (lower value in the range of line for septic tank in Table 6.3 updated in 2019 Refinement to the 2006 IPCC Guidelines for NGGI). However, the operation of cesspools in Slovakia (and everywhere in the world) works in such a way that the contents (raw wastewater from household) of cesspools are regularly pumped off at intervals of 1 – 1.5 months. During that time, methane production cannot develop to the same extent as in septic tanks, where the sludge emptied interval is about 1-1.5 years. After consultations with several experts, adjusted values of MCF for septic tanks to the conditions in the cesspools were implemented. Due to the ratio of the intervals of withdrawing the contents of cesspools compared to the discharge of septic tanks (10 times more often), the recommended MCF = 0.5 (for septic tanks) was reduced to MCF = 0.05 (for cesspools).

After consultation of this procedure with several experts in methane production and wastewater treatment (Slovak University of Technology, Slovak Water Companies), and also verification with several literature sources, where it is clearly defined that the production in a properly operated cesspool (accumulation storage tank with SRT = 30-40 days) is significantly lower than in a septic tank (tank for mechanical pre-treatment of wastewater with SRT = 300-400 days).

It should be added that the operation of septic tanks as individual methods of wastewater treatment is prohibited, only their operation as a stage of pre-treatment before the biological treatment stage, e.g. before the constructed wetlands, is allowed, while the number of such treatment plants in Slovakia is minimal.

Subsequently, the contents of the cesspools are legally transported to nearby WWTPs (it is included in the measurement of BOD<sub>5</sub> at the input to the WWTP) – an expert estimate of about 45%. Another about 30% of the content of cesspools (illegally) enters rivers and the remaining about 25% of the total content of cesspools reaches the soils. MCFs are used for these parts of wastewater streams in accordance with Table 6.3 of the updated in 2019 Refinement to the 2006 IPCC Guidelines.

Changes and the introduction of new default factors as well as the application of new calculation procedures led to significant changes in the resulting emissions in domestic wastewater sectors as well as in total emissions production. With the change of methane emissions, also implied methane emission factors were changed. In following tables, these changes are recorded.

**Table 7.8: Recalculations of the 5.D.1 and 5.D categories and comparison of the submissions for methane emissions**

YEAR	CH <sub>4</sub> emissions in 5.D.1 – Domestic WW			Total CH <sub>4</sub> emissions		
	kt		%	kt		%
	2024	2025		2024	2025	
1990	13.21	5.21	-60.58%	14.50	6.50	-55.21%
1991	13.13	5.12	-60.98%	14.36	6.35	-55.77%
1992	13.09	5.09	-61.14%	14.27	6.26	-56.12%
1993	13.05	5.05	-61.33%	14.17	6.17	-56.49%
1994	13.14	5.14	-60.88%	14.17	6.17	-56.48%
1995	13.00	4.99	-61.57%	13.92	5.92	-57.44%
1996	12.96	4.96	-61.71%	13.74	5.74	-58.24%
1997	12.96	4.96	-61.73%	13.69	5.69	-58.45%
1998	13.01	5.01	-61.48%	13.75	5.75	-58.20%
1999	12.95	4.95	-61.75%	13.65	5.65	-58.63%
2000	12.87	4.88	-62.12%	13.67	5.68	-58.48%
2001	12.86	4.87	-62.16%	13.61	5.62	-58.72%
2002	12.81	4.90	-61.79%	13.51	5.60	-58.58%
2003	12.59	4.75	-62.25%	13.32	5.48	-58.83%
2004	12.31	4.55	-63.02%	12.92	5.16	-60.08%
2005	12.06	4.38	-63.68%	12.52	4.84	-61.32%
2006	11.89	4.29	-63.91%	12.25	4.65	-62.06%
2007	11.78	4.26	-63.83%	12.13	4.61	-61.98%
2008	11.57	4.05	-64.99%	11.93	4.41	-63.02%
2009	11.47	4.03	-64.84%	11.85	4.41	-62.74%
2010	11.37	4.01	-64.70%	11.74	4.38	-62.68%
2011	11.15	3.88	-65.24%	11.45	4.17	-63.55%
2012	10.91	3.71	-65.97%	11.19	3.99	-64.35%
2013	10.78	3.66	-66.01%	11.05	3.94	-64.37%
2014	10.36	3.32	-67.92%	10.61	3.57	-66.33%
2015	10.61	3.65	-65.58%	10.85	3.89	-64.11%
2016	10.42	3.54	-66.00%	10.66	3.79	-64.48%
2017	10.34	3.55	-65.70%	10.58	3.78	-64.27%
2018	10.29	3.57	-65.28%	10.48	3.77	-64.04%
2019	10.12	3.53	-65.13%	10.33	3.74	-63.84%
2020	9.96	3.49	-64.97%	10.14	3.67	-63.81%
2021	9.56	3.38	-64.65%	9.75	3.57	-63.37%
2022	9.38	3.26	-65.23%	9.54	3.42	-64.16%

## 7.4. CATEGORY-SPECIFIC IMPROVEMENTS AND IMPLEMENTATION OF RECOMMENDATIONS

No UNFCCC review was organised in 2024 and all previous recommendations from the UNFCCC review were implemented in 2023 submission (see [Chapter 7.4](#) of the SVK NIR 2023). However, sectoral experts implemented in this submission (2025) several methodological changes led to major improvements of the waste inventory. Discussion with European waste expert during voluntary review process in the second half of 2024 led to recalculations in 5.a and 5.D categories. The recalculations reflected the IPCC 2019 Refinement to the IPCC 2006 Guidelines, changes were implemented in all

aspects and influenced the inventory on category and gas level. More information can be found in individual chapters of this document.


## 7.5. SOLID WASTE DISPOSAL (CRF 5.A)

Emissions from Solid waste disposal sites (SWDS) are the major emissions source in the Waste sector. Methane emissions are estimated separately for municipal solid waste and non-municipal (industrial) solid waste disposal using IPCC Waste Model. Emissions of CO<sub>2</sub> influencing national total are not occurring in this category as burning waste on landfills is prohibited by law. The unmanaged waste disposal site was not occurring in the Slovak Republic during the reported period.

Total methane emissions in category CRF 5.A were 41.52 Gg (1 162.62 Gg of CO<sub>2</sub> eq.) in 2023 as is shown in [Table 7.1](#). Emissions from landfilling have fallen more sharply in recent years. This results from an active waste policy in the Slovak Republic in the sense of the waste hierarchy according to Article 4 of the Waste Directive. Since 2010, landfilled MSW has decreased from 1.412 million tonnes to 0.939 million tonnes (-34%). Due to the impact of separate collection of paper + food + garden waste, the amount of landfilled MSW with DOC > 0 has also decreased very significantly in this period from the original 829,000 t to the current 355,000 t (-58%). "Net" emissions (after deducting oxidised methane and methane used for electricity generation) for 2023 are 8% lower than in 2018 when the maximum level of emissions from landfilling was reached.

In accordance with the European Landfill Directive (1999/31 EC), Slovak waste legislation also distinguishes between three classes of landfills (= SWDS). Landfills for inert waste are not a source of GHG emissions and waste landfilled for this class of landfills has not been included in the emission calculations. Landfill emissions were calculated separately for municipal waste (MSW) and separately for industrial waste (ISW) as is shown in [Table 7.9](#). MSW share on landfilled waste by 37%, followed by energy industry waste (34%), wastes from waste management facilities (13%), construction waste (8%) and waste packages by 4%. These five groups (20+10+19+17+15) represented 96% of landfilled waste.

**Figure 7.4:** Major groups of landfilled solid waste (ŠÚ SR) in tons in Slovakia in recent years



**Table 7.9:** Activity data from the total SWDS in Slovakia (MSW + ISW) in particular years

YEAR	TOTAL SWDS	MUNICIPAL SOLID WASTE			INDUSTRIAL SOLID WASTE				
		GROUP 20	MSW to SWDS	Share	GROUP 1-19	ISW to SWDS	Share	ISWDS DOC > 0	Share
		tons		%	tons		%	tons	%
2005	1 417 993	1 558 283	1 226 586	78.7%	9 346 816	2 888 366	30.9%	191 407	2.0%
2006	1 509 768	1 623 302	1 259 613	77.6%	12 879 757	5 646 833	43.8%	250 154	1.9%

YEAR	TOTAL SWDS	MUNICIPAL SOLID WASTE			INDUSTRIAL SOLID WASTE				
		GROUP 20	MSW to SWDS	Share	GROUP 1-19	ISW to SWDS	Share	ISWDS DOC > 0	Share
		tons		%	tons		%	tons	%
2007	1 582 192	1 668 660	1 294 853	77.6%	9 252 161	4 261 633	46.1%	287 339	3.1%
2008	1 599 325	1 772 456	1 350 862	76.2%	9 683 380	3 215 530	33.2%	248 463	2.6%
2009	1 597 757	1 745 450	1 349 267	77.3%	6 808 199	2 675 101	39.3%	248 491	3.6%
2010	1 620 725	1 808 506	1 411 543	78.1%	7 814 887	2 483 878	31.8%	209 183	2.7%
2011	1 577 264	1 766 990	1 320 073	74.7%	8 605 496	2 875 331	33.4%	257 191	3.0%
2012	1 488 803	1 750 775	1 297 480	74.1%	7 016 588	2 803 452	40.0%	191 323	2.7%
2013	1 419 773	1 744 429	1 201 906	68.9%	8 216 667	3 797 353	46.2%	217 867	2.7%
2014	1 358 482	1 830 167	1 210 043	66.1%	7 324 208	2 620 480	35.8%	148 439	2.0%
2015	1 457 463	1 888 456	1 303 845	69.0%	8 782 522	2 707 543	30.8%	153 618	1.8%
2016	1 433 301	1 953 478	1 289 895	66.0%	8 717 765	2 499 439	28.7%	143 406	1.7%
2017	1 426 781	2 136 952	1 312 787	61.4%	10 115 259	2 517 432	24.9%	113 994	1.2%
2018	1 350 732	2 325 178	1 250 280	53.8%	10 142 462	2 093 797	20.6%	100 452	1.1%
2019	1 292 728	2 369 725	1 198 249	50.6%	10 037 942	1 666 717	16.6%	94 479	1.0%
2020	1 279 209	2 596 725	1 121 159	43.2%	10 516 841	1 832 869	17.4%	158 050	1.2%
2021	1 136 311	2 705 327	987 539	36.5%	9 943 797	1 470 103	14.8%	148 772	1.5%
2022	1 116 891	2 597 457	982 662	37.8%	10 593 124	1 720 048	16.2%	134 229	1.2%
2023	1 059 209	2 560 971	938 898	36.7%	11 005 377	1 519 287	13.8%	120 311	1.0%

### 7.5.1. Municipal Waste Disposal Sites (Managed)

The first legislation governing the disposal of waste in Slovakia was adopted in 1991, followed by implementing regulations in 1992. The Act No 239/1991 stipulated basic requirements for the operation of waste disposal sites and Governmental Regulation No 606/1992 in the Annex 5 defined three classes of waste disposal sites and technical requirements for their construction. New legislative regulation on solid waste management and disposal entered into force on 1<sup>st</sup> July 2001 in the process of harmonisation with the EU legislation. The Act No. 223/2001 Coll. and Decree of the Ministry of Environment No. 283/2001 Coll. contain new instruments for waste disposal minimisation, monitoring of waste sites and landfill gas generation. The importance to increase the share of recycled waste resulted in the adoption of the Act No. 79/2015 on waste, which introduces the extended responsibility of producers (mix packages) and transfers organisation and financing waste recycling schemes from the state to producer responsibility organisation. This change indicates an increase of waste diverted from disposal.

Currently in the Slovak Republic, municipalities are obliged to introduce and ensure the implementation of separate collection for the separate collection of classical components of MSW, i.e. paper and cardboard, glass, plastics and metals, and biodegradable municipal waste. Long-term monitoring of separate collection of MSW shows an increasing trend in the number of separated components. According to the officially published data from the ŠÚ SR, there has been a year-on-year increase in the rate of sorted municipal waste collection. For example, between 2011 and 2023, the amount of separated garden waste increased from 89 276 t to 348 287 t, separated food waste from 1 855 t to 62 175 t and separated paper from 44 719 t to 236 300 t. It is therefore logical to assume that the percentage of these three components in landfilled mixed municipal waste will also decrease.

Decreasing trend in landfilling is visible in the last decade; however, the total municipal waste production increased and represents more than 472 kg/capita/year. In addition, the share of MSW ending on landfills is decreasing compared to about 88% in 1995 represents 37% in 2023 according to the ŠÚ SR and MŽP SR data.

---

At the time, there are almost 64 non-hazardous waste (NNO) landfills operating in Slovakia, which dispose of municipal and industrial waste in SWDS. Nowadays, all of them were operating as anaerobic sites (CRF 5.A.1.a). Methane recovery takes place at only 8 sites, mostly for energy generation at the SWDS receiving municipal solid waste. Before adopting legislation regulating waste management in 1991, municipal solid waste had been disposed mostly in an uncontrolled manner.

Development of engineered, controlled landfills, including gas collection systems, started in 1993 and old dumps as a disposal destination were gradually replaced over the following decade. It takes some time until a landfill cell is filled, closed and gas generation starts in the landfill body. Thus, the first attempts to flare landfill gas were introduced in 2004.

Law does not allow burning waste on SWDS, neither is it part of operation practice. Fires, which rarely occur on landfills, are considered as emergencies and are extinguished as soon as possible.

Following the IPCC 2006 GL methodology, emissions from the SWDS should be estimated separately for MSW and non-MSW what is industrial solid waste. The CRF tables provide emissions reporting from these two sources together, but data are presented as disaggregated to the MSW and ISW ([Table 7.10](#)).

### Methodological issues

Methane emissions from MSW disposed to SWDSs were calculated using the IPCC 2006 Waste Model. Tier 2 approach is used for emission estimations, using default parameters and country-specific activity data. The IPCC 2006 Waste Model was set to option "Waste by Composition" because the composition of municipal solid waste was modelling including the impact of waste separation.

**Methane Generation Rate (k)** - defines how fast waste decomposes. IPCC default k-rates are estimated as a function of climate zone, which is characterised by mean annual temperature (MAT) and the ratio of mean annual precipitation and potential evapotranspiration (MAP/PET). Slovakia belongs to the temperate climate zone, because even the warmest parts of Slovakia have MAT around 10°C.

Slovakia falls into a climate area where precipitation exceeds evaporation, although some southern areas of the country fall into a precipitation shadow with the opposite trend.

On the other hand, "k" is also depending on the operation of site. Common praxis in Slovakia, mostly in summer months, is backwards recirculation of landfill leachate into the site to support biodegradability of waste and vaporisation. This praxis lowers the costs on the treatment of this landfill waste liquid and this quantity can be higher than rainfall (in summer 50-90 mm/month). Estimation of k-parameter only from the climatic zone and rainfall can lead to an underestimation of real value of this parameter.

Therefore, "k" values in the sense of IPCC 2006 GL (Table 3.3) for the wet climate zone were used in the calculations.

**Degradable Organic Carbon (DOC)** - this parameter identifies organic carbon in waste, which is accessible to biochemical decomposition. DOC of municipal solid waste was estimated from the MSW composition in an IPCC model taking into account changes in composition due to changes of fuel used for heating and changes due to separation of recyclable and compostable materials. The DOC firstly growing due to increasing of biodegradable fraction in the MSW, then decreasing due to diversion of recyclable and compostable waste from landfilled waste.

The content of DOC in MSW began to rise in the late 1990s after a change in the social system and with an increasing living standard. This was mainly reflected in the increased share of food and packaging (paper) in the MSW. The turning point came around 2010, when, in accordance with the Environmental Kuznets Curve theory, the growing environmental awareness of the population began to manifest itself and the DOC value began to decline. Despite the significant growth in the production of municipal waste after 2013, the separate collection of usable components is increasingly being promoted, and a smaller share of MSW ends up in landfills every year (a decrease from 88% to 37%). In recent years, new Mechanical Biological Treatment facilities for the treatment of mixed municipal waste have also

contributed to the change in the DOC of landfilled MSW (20 03 01). Due to the current ongoing change of the Waste Information System (IS OH) and unavailable data, it was not possible to accurately determine the DOC value for recent years. The data therefore only an expert estimate.

**Methane Correction Factor (MCF)** – this parameter reflects the disposal management practices. Analysis of disposal sites database of the ŠGÚ DŠ by depth, year of creation and deposited volume resulted in the development of the MCF. The trend of MCF reflects the impact of waste legislation, causing continuous replacement of semi-aerobic dumps by controlled anaerobic landfills in the period 1990 – 2009. Based on the statistical research, Slovakia operated many small-scale landfill sites. Very small-scale landfills sites ( $\Sigma W < 5\,000$  t/y) represent around 18% of existing SWDS in Slovakia. The criteria for managed-anaerobic landfills are difficult to follow – so these sites can be categorised as shallow. Conditions on sites can be categorised more as aerobic, than anaerobic. It means, that the MCF = 1.0 is used since 2010 (**Table 7.10**).

**Table 7.10: Development of the Methane Correction Factor (MCF)**

Year	1950	1960	1970	1980	1990	1995	2000	2005	2010 – 2023
MCF	0.54	0.54	0.54	0.56	0.56	0.61	0.74	0.86	1.0

**Oxidation Factor (OX)** – reflects the amount of CH<sub>4</sub> from SWDS that is oxidised in the soil or other material covering the waste. There is no reliable information on past practice in covering disposal sites with soil. Due to a lack of relevant information about the real value of the OX in the landfill in Slovakia, the IPCC 2006 GL (Table 3.2) value OX = 0.1 for managed landfill covered with CH<sub>4</sub> oxidising material was used since 1994.

The oxidation factor (10%) was applied in Slovakia reflecting the ESD review 2020 since 1994. The methane emissions were reduced by the default value of the oxidation factor according to the IPCC, when the first anaerobic landfills began to operate. The estimation of the years 1950 – 1993 are without the oxidation factor (OX = 0).

**Methane Recovery (R)** – means combusting landfill gas generated at SWDS in a flare or energy device. Slovakia reported the amount of CH<sub>4</sub> flared without energy recovery for the years 2006 – 2011. This practise not exists after 2011.

The Regulatory Office for Network Industries (ÚRSO) statistically recorded and published data on electricity generated from the LFG since 2011. The lists of companies who received subsidy for producing electricity from renewable sources, including landfill gas is available. The amount of recovered methane is calculated from electricity produced in MWh and the calorific value of the LFG. Expert judgement is that 50% of the LFG is methane and lower heating value (LHV = 18 MJ/m<sup>3</sup>). Emissions from LFG flared with energy use is provided and reported in CRF Table 1.A.5.a. Increase of methane recovery from landfilling is not expected in the next years due to lowering of subsidies for energy recovery LFG. Conversely, the increasing diversion of biodegradable waste away from landfill is leading to a decline in both the quantity and quality of LFG in existing landfills. This is reflected in the cessation of the use of methane from LFG for electricity generation. In recent years, the number of plants has 8.

After further consultations with the ÚRSO, small corrections were made to the data on the amount of electricity produced in older years (2022) and a unified calculation of the methane used for the entire period under the same combustion conditions was introduced (**Table 7.11**).

**Table 7.11: Correction of the LFG calculation based on the ÚRSO data for the years 2011 – 2023**

YEAR	ELECTRICITY PRODUCTION	LFG FOR ELECTRICITY PRODUCTION	METHANE RECOVERY
	MWh	m <sup>3</sup> /year	tons
2011	6 463	4 421 775	1.579
2012	8 627	5 902 314	2.108

YEAR	ELECTRICITY PRODUCTION	LFG FOR ELECTRICITY PRODUCTION	METHANE RECOVERY
	<i>MWh</i>	<i>m<sup>3</sup>/year</i>	<i>tons</i>
2013	8 831	6 041 884	2.158
2014	11 141	7 622 311	2.722
2015	8 373	5 728 535	2.046
2016	9 946	6 804 731	2.430
2017	10 223	6 994 245	2.498
2018	10 092	6 904 619	2.466
2019	10 480	7 170 760	2.561
2020	10 794	7 387 158	2.637
2021	9 575	7 607 853	2.363
2022	9 087	6 203 912	2.218
2023	8 056	5 580 369	1.995

Data about amount of methane used for energy production have been determined only by calculation so far. MŽP SR does not have records or database about number of landfills where LFG is used for energy production or incinerated on flares. The only source of information on the use of LFG from landfills is the ÚRSO data on the amount of electricity produced from landfill gas. Since 2011, this office has been publishing the amount of electricity produced from LFG (MWh) per year for individual companies. Due to the financial bonus that is paid by the state for this amount of electricity produced from waste, the data on the amount of electricity is relatively closely and strictly monitored and controlled by the ÚRSO. Therefore, it can be considered this information to be accurate and reliable.

In determining the amount of methane used from landfilled waste, a back-calculation was used for common in assessing the landfill in terms of prospects for its energy use, just in reverse order. The amount of electricity actually produced is known and the amount of LFG (or methane) used was recalculated with the following formula:

$$\text{LFG vol.} = \text{EG} * \text{Cf} / \text{LHV} * \text{Ef}$$

where LFG vol. = amount of landfill gas used in m<sup>3</sup>, EG = Electricity generated (MWh), Cf = conversion from MWh to MJ, LHV = Low Heating Value of LFG (18 MJ/m<sup>3</sup>), Ef = Electricity conversion efficiency (30%).

Based on this formula, the amount of LFG processed (m<sup>3</sup>) was calculated and, with the theoretical methane content (50%), the weight of fraction RECOVERY methane for each year of calculation was determined. Comparing these data with data from stationary sources database (NEIS) followed, which contains, among other things, data on the amount of used landfill gas. Comparing these two databases (ÚRSO and NEIS), resulted in conclusions that, especially in the past, not all companies were included in both databases.

**Activity data** – Total MSW disposed on landfills is used as activity data for estimation of methane emissions from the SWDS annually. Additionally, the overall MSW balance is used for verification of these activity data. The ŠÚ SR published data on MSW generation and disposal only since 1993. Although this creates a timeline of 26 years, additional historical data had to be generated for the use of the FOD method. Analysis of MSW generation data shows a large difference in MSW generation in the years 1992 – 1994, compared to 1995 – 2011. This can be explained by a “learning period” when waste generators were getting familiar with the new system of data recording. Therefore, these “inflated” data were excluded from methane emissions estimation and replaced by interpolated data, as is explained below. It may be interesting, that similar, but smaller “inflation” of data appears also in the period 2002 – 2005, when the EU Waste Classification System was introduced in Slovakia. Extrapolation of data back to 1950 was made by correlating annual waste per capita to index of real wage (IRW), which is defined as nominal wages index corrected for changes in purchasing power measured by the

consumer price index (nominal wage index/consumer price index). The ŠÚ SR and before 1993, the Statistical Office of the ČSSR has been publishing this index since 1948.

When assessing the amount of MSW disposed to SWDSs, the key factor influencing the MSW management practice is operation of only two MSW incinerators (Bratislava and Košice). These two incinerators burned on average 150 Gg of MSW per year in the period 1993 – 2011 (Bratislava 100 Gg/yr, Košice 50 Gg/yr) and 185-210 Gg of MSW (period 2011 – 2020). According to data published in the yearbooks of the Statistical Office of the Slovak Republic, the amount of MSW waste incinerated for the years 2010 – 2021 never reached more than 10% of the total MSW production in Slovakia.

An overview of activity data for the entire timeline is shown in [Table 7.12](#). Emissions from municipal waste disposed to SWDSs were estimated from unmanaged disposal sites, created before 1993, managed landfills developed after 1993 and considering that part of municipal solid waste still could be illegally disposed in shallow unmanaged sites. Waste generated from industrial, agricultural and other non-municipal activities is discussed in separate [Chapter 7.5.2](#). The entire time series were recalculated with the use of the IPCC 2006 GL - Waste Model. Consistency of extrapolation of disposed municipal waste time-series is ensured by using country-specific data on mid-year population and the IRW, both available as continuous time series since 1950. Waste composition changes are derived from a share of household using gas as heating fuel, this parameter was identified in national censuses, which are organized in Slovakia every 10 years. The dependence of municipal waste production in Slovakia on GDP (or HFC = Households Final Consumptions) has already been mentioned in [Chapter 7.5](#).

**Table 7.12: Activity data used for the solid waste disposal sites methane emissions estimation**

YEAR	POPULATION	IRW*/HFC**	MSW	MSW/CAP	MSW TO SWDS	MSW TO SWDS
		GDP/capita	kt	kt/capita/year	%	kt
1950	3 463 446	75.3	385 745	111	100%	385 745
1960	3 994 270	124.7	736 901	184	100%	736 901
1970	4 528 459	158.5	1 061 904	234	100%	1 061 904
1980	4 984 331	194.2	1 432 061	287	90%	1 288 855
1990	5 297 774	194.0	1 520 550	287	90%	1 368 495
1995	5 363 676	159.8	1 268 355	236	88%	1 116 152
2000	5 400 679	not relevant	1 339 491	248	79%	1 055 925
2005	5 387 285	27 276	1 558 263	289	79%	1 226 570
2010	5 431 024	38 286	1 808 506	333	78%	1 411 543
2011	5 398 384	43 155	1 766 990	327	75%	1 320 073
2012	5 407 579	43 388	1 750 775	324	74%	1 297 480
2013	5 413 393	42 671	1 744 429	322	69%	1 201 906
2014	5 418 649	43 441	1 830 167	338	66%	1 210 043
2015	5 423 800	44 857	1 888 456	348	69%	1 303 845
2016	5 430 798	46 504	1 953 478	360	66%	1 289 895
2017	5 437 754	49 370	2 136 952	393	61%	1 312 787
2018	5 445 382	51 399	2 325 178	427	54%	1 250 280
2019	5 452 257	52 891	2 369 725	434	51%	1 198 249
2020	5 459 781	53 280	2 596 725	446	43%	1 121 159
2021	5 434 712	54 850	2 705 327	497	37%	987 539
2022	5 431 056	57 616	2 597 457	478	38%	982 662
2023	5 426 820	55 802	2 560 971	472	37%	938 898

IRW = income real wage, since the year 2000 not relevant, HFC = household final consumption (EUR) – only year 2005 – 2023

### Uncertainties

The default IPCC uncertainties were used and where possible were adjusted to reflect country-specific data. The total uncertainty of emissions from MSW disposal was estimated to  $\pm 30\%$  ([Table 7.13](#)).



**Table 7.13: Uncertainties used in MSW disposal**

ACTIVITY DATA AND EMISSION FACTORS	UNCERTAINTY RANGE
Fraction of MSW sent to SWDS (MSWF)	±30% for waste data in period 1950 – 1994 ±10% for waste data in period 1995 – 2004 ±5% for waste data in period 2005 – 2021
Total uncertainty of waste composition:	±50% for the entire modelled period
Degradable Organic Carbon (DOC):	Default values:
Paper/cardboard	0.400
Textiles	0.240
Food waste	0.150
Garden and Park waste	0.200
Wood waste	0.430
Fraction of Degradable Organic Carbon Dec. (DOCf) = 0.5	±5% (IPCC default values used)
Methane Correction Factor (MCF):	IPCC default values used:
= 1.0	0%
= 0.8	±20%
= 0.4	±30%
Fraction of CH <sub>4</sub> in generated Landfill Gas (F) = 0.5	±5% (IPCC default value used)

### Source-specific recalculations

No methodological recalculations were implemented in this submission. However, analysis of the composition of household waste has been carried out so far and the available results are not representative of the whole of Slovakia, as they mostly concern small municipalities. It is likely that the composition of municipal waste in small municipalities (< 1 000 inhabitants) will be different than in cities (>10 000 inhabitants). In addition, there are significant regional differences in individual production of municipal waste in Slovakia (Trnava region = 657 kg/capita/year, Košice region = 375 kg/capita/year). Therefore, it can be expected that the composition of municipal waste will also differ regionally.


### 7.5.2. Non-municipal Disposal Sites (Industrial)

In the past, industrial waste was landfilled together with municipal waste in common landfills. It was not until 1991, when the First Waste Act was passed, that some large industrial companies built their own landfills to store their industrial waste. After 2001 (the Second Waste Act), there are three classes of landfills in Slovakia – for inert waste (IO), non-hazardous waste (NNO) and hazardous waste (NO). At the vast majority of NNO landfills (approx. 75), municipal and industrial waste (MSW + ISW) is landfilled together. Only a few large industrial companies operate their own NNO landfills for their industrial waste without MSW. However, the number of such landfills is relatively small and only specific wastes from the energy or metallurgical industries, so waste without organically degradable carbon (DOC = 0), are landfilled.

Since 2005, the records of production and waste management according to the EWC have been significantly improved. The data in the information system managed by the MŽP SR ([IS\\_OH](#)) show that there is a change in the composition of landfills for industrial waste. On the other hand, it is necessary to evaluate positively the deviation from landfilling at ISW in recent years. The maximum volumes of landfilled ISW were recorded in the years 2006 – 2011, or shortly after Slovakia's accession to the EU. During this period, the annual quantities of landfilled ISWs ranged from 250 to 300 000 tons of waste. After the 2011 crisis and its repercussions, the amount of landfilled waste decreased in the years 2014 – 2023 on the level of approximately 130 kt. (95 - 150 kt)

This trend in the decrease in the amount and composition of landfilled ISW is also related to the significant decrease in methane emissions produced in recent years. Compared to 1990, methane emissions from ISW waste decreased by -23%. More information on trend is in [Figure 7.5](#).

**Figure 7.5: Share of ISW on methane emissions production**



### Methodological issues

The first data on ISW are from the year 1997, but due to change of waste classification system in 2002, reliable continuous time series started in 2005. Preparation of time series back to 1950 is based on two periods in development of the Slovak economy. The first period, centrally planned economy from 1950 – 1989, is characterised by low environmental standards, little innovations and modernisation. For the second period, economic transformation from 1990 – 2013, is typical rapid modernisation, closing of inefficient and polluting industries and strengthening environmental standards. Such development cannot be described by standard approach correlating waste generation to the GDP as recommended in the IPCC 2006 GL.

Landfilled biodegradable non-MSW was selected from the database based on the EWC, which is maintained by the MŽP SR and published by the ŠÚ SR. This database is updated annually and summarises reports on waste from individual waste generators. All waste types discussed in the IPCC 2006 GL can be identified in the waste database.


Consistency of time series is ensured by using continuous time series of sectoral growth indicators since 1950. Activity data were completely recalculated in this submission. Time series consistency was maintained by replacing data obtained by waste classification used in 1990 – 2005 using extrapolations to avoid discrepancies caused due to differences in waste classification.

The European Waste Catalogue (EWC) contains 19 groups of industrial waste (=ISW) and one group (20) of municipal waste. For the calculation of emissions from ISW landfills, groups of waste that do not contain biodegradable carbon (DOC) and therefore do not produce GHG emissions were excluded. These were groups 01, 06, 09, 10, 11 and 16. Due to administrative complexity, in the next step, those groups of waste were also excluded from the calculations, which in the given year reached a share in the total landfilled waste  $W_i < 0.2\% \sum W_i$ . It was usually waste from groups 05, 07, 08, 12, 13, 14 and 18. Due to their mass representation in landfilled waste, a completely negligible contribution to the total emissions in a given year can be expected. From the remaining 6 groups of waste (02, 03, 04, 15, 17 and 19), individual types of waste were selected in accordance with the IPCC methodology. It was summarized by weight into seven types of waste according to the main degradable component: Food, Wood, Paper, Textile, C + D waste, Mix\_Package and Sludge. Waste from greenery (Garden) was finally also excluded from the calculations, as it landfilled proportion was very low (approximately 500 to 1 000 t/y). An overview of individual types of landfilled ISW is provided in [Table 7.14](#). and on [Figure 7.6](#).

**Table 7.14: DOC and k-rate parameters used in IPCC Waste Model for ISW**

WASTE TYPE	DOC	k	REFERENCE	MAIN WASTE (EWC)
Food	0.15	0.185	IPCC default	groups 02 02, 02 03 and 02 06
Garden and Park	0.20	0.100	IPCC default	groups 02 01 and 19 05
Paper / Cardboard	0.40	0.060	IPCC default	groups 03 03 07+8, 09 01 07+8, 15 01 01 and 19 12 01
Textiles	0.24	0.060	IPCC default	groups 04 01, 15 01 09, 15 02 02 and 19 12 08
Wood	0.43	0.030	IPCC default	groups 03 01, 15 01 03, 17 02 01 and 19 12 06+7
Sludge	0.355	0.185	IPCC default	groups 19 08 05 and 19 08 11-14
C+D waste	0.05	0.030		group 17 09 03+4
Mix_Package	0.10	0.060		group 15 01 06+10

**Figure 7.6: An overview of individual types of landfilled ISW in tons in 2005 – 2023**



### Uncertainties

Uncertainties related to activity data for ISW are particularly significant for the period 1950 – 1990. In accordance with the IPCC 2006 GL (Chapter 3.6), data on the amount of landfilled ISW for this period were only estimated based on the GDP growth and the industrial production index. For the period 1991 to 2004, there are already better statistics on the production and management of industrial waste. However, the records are according to the old (national) waste catalogue, which was not fully compatible with the current EWC. Since 2005, the data have been used from the documents on waste management of the ŠÚ SR and the MŽP SR. During the detailed verification process, discrepancies were found between these two databases in recent years. These discrepancies did not reach 3% and did not have a significant impact on the estimation.

Periods 1950 – 1990, 1991 – 2004 and 2005 – 2023 can be characterised by changes in legislation and information systems. Due to the calculation of emissions by the FOD method, total emissions are spread over a longer period according to the half-time of decay. It should be noted, that the actual composition of the ISW for the 1950 – 1990 period is estimated with a high uncertainty. However, as already stated in [Chapter 7.2](#), the half-time of decay for most types of these wastes (with the exception of wood) according to IPCC 2006 GL (Table 3.4) is from 4 to 12 years for Slovakia. This means that waste deposited in landfills before 1995 produces zero emissions nowadays, assuming standard conditions for the degradation of organic carbon in the landfill.

[Table 7.14](#) shows, that the weight of landfilled waste fraction C + D and Mix\_Package is much more significant than other types of waste (Paper, Textile or Wood). These two types of waste are

characterized by a relatively high degree of uncertainty on DOC as resulted from their mixed nature. The default IPCC uncertainties were used and adjusted (where possible) to reflect country-specific data or circumstances. The total uncertainty of emissions from disposal of ISW was estimated to be  $\pm 27\%$ .

**Table 7.15: Uncertainties for non-MSW disposal**

ACTIVITY DATA AND EMISSION FACTORS	UNCERTAINTY RANGE
Amount of disposed ISW	$\pm 50\%$ for waste data in period 1950 – 2004 $\pm 5\%$ for waste data in period 2005 – 2021
Degradable Organic Carbon (DOC) =	Default values:
Paper/cardboard	0.40
Textiles	0.24
Food waste	0.15
Wood waste	0.43
Sludge	0.355
C+D waste	0.05
Mix_Package waste	0.10
Fraction of Degradable Organic Carbon Dec. (DOCf) = 0,5	$\pm 5\%$ (IPCC default value was used)
Methane Correction Factor (MCF)	IPCC default values used:
= 1.0	+0%
= 0.8	$\pm 20\%$
= 0.4	$\pm 30\%$
Fraction of CH <sub>4</sub> in generated Landfill Gas (F) = 0.5	$\pm 5\%$ IPCC default values used
k-rate =	Default values:
Paper/cardboard	0.06
Textiles	0.06
Food waste	0.185
Wood waste	0.03
Sludge	0.185
C+D waste	0.03
Mix_Package waste	0.06

### Source-specific recalculations

No recalculations of ISW disposal were implemented in this submission.

## 7.6. BIOLOGICAL TREATMENT OF SOLID WASTE (CRT 5.B)

Waste Framework Directive 2008/98/EC requires the Member States to reduce the disposal of biodegradable waste in landfills. The EU directive was transposed into the Slovak legislation in the Act No 223/2001, Art, 18 (4) m), which stipulates that disposal of biologically degradable waste from parks and gardens together with the MSW is banned in the Slovak Republic from January 2006. There is a range of private and municipal companies, which provide composting of municipal and agricultural waste. **Table 7.16** shows an overview of municipal and industrial composting. With the support of the EU and Governmental grants, the number of municipalities composting waste is growing fast. While 24% of municipalities participated in waste composting in 2002, this number increased to more than 90%. According to the EUROSTAT data 60 kg per capita of biologically degradable waste was recycled in 2023 in comparison with 2005, representing an increase of more than 100% to the 2005 and an increase by 21% compared to the previous year.

### 7.6.1. Composting (CRT 5.B.1)

The most common is windrow composting. More sophisticated technologies, like anaerobic treatment or mechanical-biological treatment (MBT) plants, are not used. Data on composting are disaggregated into composting of MSW reported in the CRT table 5.B.1.a and composting of non-MSW reported in the CRT table 5.B.1.b.

**Table 7.16:** The overview of municipal and industrial composting in 1990 – 2023

YEAR	MSW (CRT 5.B.1.a)			NON-MSW (CRT 5.B.1.b)		
	WASTE TREATED	CH <sub>4</sub>	N <sub>2</sub> O	WASTE TREATED	CH <sub>4</sub>	N <sub>2</sub> O
	<i>kt (dm)</i>	<i>Gg</i>		<i>kt (dm)</i>	<i>Gg</i>	
1990	8.00	0.08	0.00	251.60	2.52	0.15
1995	14.18	0.14	0.01	251.60	2.52	0.15
2000	14.54	0.15	0.01	251.60	2.52	0.15
2005	51.25	0.51	0.03	231.66	2.32	0.14
2010	83.43	0.83	0.05	231.42	2.31	0.14
2011	90.05	0.90	0.05	261.02	2.61	0.16
2012	95.55	0.96	0.06	290.62	2.91	0.17
2013	97.53	0.98	0.06	247.94	2.48	0.15
2014	94.57	0.95	0.06	291.24	2.91	0.17
2015	114.88	1.15	0.07	374.00	3.74	0.22
2016	119.69	1.20	0.07	285.30	2.85	0.17
2017	161.36	1.61	0.10	306.95	3.07	0.18
2018	184.41	1.84	0.11	313.42	3.13	0.19
2019	208.97	2.09	0.13	262.04	2.62	0.16
2020	301.89	3.02	0.18	315.77	3.16	0.19
2021	332.27	3.32	0.20	272.28	2.72	0.16
2022	335.43	3.35	0.20	398.47	3.98	0.24
2023	350.04	3.50	0.21	300.89	3.01	0.18

### 7.6.2. Methodological Issues

The default tier 1 approach from the 2006 IPCC GL was used for emission estimations in this category. Emissions from composting were estimated separately for MSW and ISW.

Default IPCC emission factors for dry weight of waste were used:

- Emission factor 10 g CH<sub>4</sub>/kg of DM waste treated;
- Emission factor 0.6 g N<sub>2</sub>O/kg of DM waste treated.

Activity data in the wet stage was taken from the publication “Waste in the Slovak Republic” and converted to dry matter for reporting purposes in 2023. The second set of activity data was taken from the Water Research Institute – responsible for collecting information regarding the recovery of sewage sludge. The activity data are consistent with the category 5.D Wastewater Treatment and Discharge. Historical activity data of composted municipal solid waste are published since 1992. The missing data for 1990 and 1991 were extrapolated with linear extrapolation.

The data on sewage sludge composting are available since 2003. The latest activity data for wastewater treatment sludge is not in a format compatible with the data series published after 2003, as the European waste catalogue methodology was implemented in 2003. Therefore, emissions from sludge for the period 1990 – 2002 are considered as not estimated. Data on industrial waste composting were collected and published since 1997. No clear trend could be identified, as data vary ±50%, thus the average of the years 2002 – 2013 was used for linear extrapolation.

### 7.6.3. Anaerobic Digestion at Biogas Facilities (CRT 5.B.2)

Anaerobic digestion of organic waste accelerates the natural decomposition of organic material without the presence of oxygen by maintaining optimal values of temperature, moisture content and pH. The generated methane is used to produce heat or electricity. Fugitive emissions of methane from anaerobic digestion due to unintentional leaks, process malfunctions or other unexpected events are reported in

the Waste sector. According to the 2006 IPCC methodology, 0 to 10% of fugitive methane emissions originated from digestion. Generated CO<sub>2</sub> emissions are of biogenic origin and is reported in Energy sector.

### Methodological issues

The default tier 1 approach from the 2006 IPCC GL was used for emission estimations in this category. Emissions from anaerobic digesters were estimated from total amount of recovery biogas.

Default IPCC emission factor for wet weight of waste were used:

- Emission factor 0.8 g CH<sub>4</sub>/kg of wet waste treated;
- Emissions of N<sub>2</sub>O emissions were assumed as negligible due to notation key NA was used.

Currently, comprehensive data on biogas stations in Slovakia are not available. Activity data on biogas was obtained directly from the National Emission Information System. The operators provided data for the years 2001 to 2023. The amount of biogas produced from the NEIS database was available and from the calculated average consumption of feedstock for the production of a unit amount of biogas (5.7 t/ths.m<sup>3</sup>) since 2001.

**Table 7.17: The overview of municipal and industrial anaerobic digestion in 2001 – 2023**

YEAR	MSW (CRT 5.B.2.a)			Non-MSW (CRT 5.B.2.b)		
	WASTE TREATED	CH <sub>4</sub>	N <sub>2</sub> O	WASTE TREATED	CH <sub>4</sub>	N <sub>2</sub> O
	kt (dm)	Gg		kt (dm)	Gg	
2001	NO	IE	NA	73.87	0.06	NA
2002	NO	IE	NA	73.61	0.06	NA
2003	NO	IE	NA	77.87	0.06	NA
2004	NO	IE	NA	85.75	0.07	NA
2005	NO	IE	NA	91.34	0.07	NA
2006	NO	IE	NA	99.86	0.08	NA
2007	NO	IE	NA	117.14	0.09	NA
2008	NO	IE	NA	132.95	0.11	NA
2009	NO	IE	NA	141.66	0.11	NA
2010	NO	IE	NA	219.67	0.18	NA
2011	NO	IE	NA	399.09	0.32	NA
2012	NO	IE	NA	749.98	0.60	NA
2013	NO	IE	NA	1 372.03	1.10	NA
2014	NO	IE	NA	1 796.88	1.44	NA
2015	NO	IE	NA	1 795.72	1.44	NA
2016	NO	IE	NA	1 829.76	1.46	NA
2017	NO	IE	NA	1 901.41	1.52	NA
2018	NO	IE	NA	1 774.35	1.42	NA
2019	NO	IE	NA	1 628.87	1.30	NA
2020	NO	IE	NA	1 529.10	1.22	NA
2021	NO	IE	NA	1 410.77	1.13	NA
2022	NO	IE	NA	1 063.59	0.85	NA
2023	NO	IE	Na	1 193.54	0.95	NA

#### 7.6.4. Uncertainties and Time-series Consistency

The default IPCC uncertainties were used and adjusted (where possible) to reflect country-specific data or circumstances. Uncertainties were calculated using the IPCC 2006 GL default method and values.

Emissions from biological treatment of waste were estimated to have  $\pm 60\%$  uncertainties as is shown in [Table 7.18](#). The highest uncertainty come from CH<sub>4</sub> and N<sub>2</sub>O emission factors.

**Table 7.18: Uncertainties for biological treatment of waste**

ACTIVITY DATA AND EMISSION FACTORS	UNCERTAINTY RANGE
Amount of composted municipal waste	$\pm 10\%$ for waste all data
Amount of composted non-MSW	$\pm 10\%$
Emission factor for CH <sub>4</sub>	4 (0.03-8)
Emission factor for N <sub>2</sub> O	0.24 (0.06-6)

### 7.6.5. Category-specific Recalculations

Emissions of CH<sub>4</sub> for category 5.B.2 – Anaerobic Digestion at Biogas Facilities were recalculated in this submission due to changes in activity data on production of biogas of other waste. The revision was carried out due to an error in the trend of activity data, which was shifted by one year. For the years 2021 to 2022, the activity data on biogas production were taken from the National Emissions Information System (NEIS – see Energy chapter). These data cover all biogas plants in Slovakia. Most visible changes are visible in 2001 and 2022 particular years (-30% and +88%). The revision of activity data lead to change of CH<sub>4</sub> emissions in this category. More information is available in [Table 7.6](#). Industrial waste composting was not recalculated. [Table 7.19](#) shows the overview of the type of communal waste composting in 2022.

**Table 7.19: The overview of type industrial composted waste in 2023**

CODE OF INDUSTRIAL WASTE	PERCENTAGE SHARE OF WASTE
Wastes from geological exploration, extraction, treatment and further processing of minerals and stone	0%
Wastes from agriculture, horticulture, forestry, hunting and fishing, aquaculture and food production and processing	20%
Wastes from wood processing and from the production of paper, board, pulp, lumber and furniture	26%
Wastes from the leather, fur and textile industries	0%
Wastes from organic chemical processes	0%
Wastes from MFSU of paints, varnishes and enamels, adhesives, sealants and printing inks	0%
Wastes from inorganic chemical processes	2%
Wastes from the photographic industry	0%
Wastes from thermal processes	0%
Wastes from chemical surface treatment of metals and coating of metals and other materials; wastes from non - ferrous hydrometallurgical processes	1%
Wastes from shaping, physical and mechanical treatment of metal and plastic surfaces	0%
Wastes from oils and liquid fuels other than edible oils	0%
Waste organic solvents, coolants and propellants	17%
Waste packaging, absorbents, cleaning cloths, filter material and protective clothing not otherwise specified	2%
Wastes not otherwise specified in this catalogue	1%
Construction and demolition wastes, including excavated soil from contaminated sites	0%
Wastes from health or veterinary care or related research other than catering and restaurant wastes not arising from direct medical care	0%
Wastes from off-site treatment plants, off-site waste water treatment plants and drinking water and industrial water treatment plants	30%

---

## 7.7. WASTE INCINERATION AND OPEN BURNING OF WASTE (CRT 5.C)

Incineration of waste and open burning of waste produces mainly CO<sub>2</sub>, in smaller amount also N<sub>2</sub>O and CH<sub>4</sub> emissions. Methane emissions may occur in case of incomplete incineration of waste. This category covers incineration of waste in dedicated incinerators and co-incineration facilities.

Open burning of waste is prohibited by law and handled as an emergency in Slovakia. Thus, no emissions were estimated for the category Open Burning of Waste (CRT 5.C.2).

Activity data for emissions estimation of waste incineration were disaggregated into waste incineration with and without energy utilisation. Each group was further divided into biogenic waste incineration and non-biogenic waste incineration. Emissions from waste incineration with energy utilisation are reported in the Energy sector, subcategory 1.A.1.a.iv (other fuels). Emissions from waste incineration without energy utilisation are reported in the Waste sector (5.C).

### 7.7.1. Waste Incineration (CRT 5.C.1)

Incineration of waste is an accepted practice in the Slovak Republic. It is regulated following EU waste legislation. After a period of modernisation of waste incineration, ones that are more modern replaced smaller and non-compliant facilities.

The following facilities for waste incineration were in operation in 2023 according to [ENVIROPORAL](#):

Two large MSW incinerators with energy utilisation;

- Five ISW incinerators (three of them with energy utilisation);
- One clinical waste incinerator without energy utilisation;
- One incinerator for rendering plant residues;
- Five facilities co-incinerating ISW (cement and lime kilns).

The estimation of emissions from waste incineration was reviewed to increase coordination between the Waste and Energy. There are two key outputs from this review:

- Emissions from the incineration of municipal and industrial waste with energy recovery are estimated and reported in the Energy sector. The increasing trend of waste-derived fuel import for the cement, lime and chemical industries is recognised.
- Emission factor for methane used in the Energy sector is now used also in the Waste sector.
- Correction of previously used notation key "IE" to "NO" in the categories 5.C.1.1.a and 5.C.1.2.a took place due to the fact, that there is no municipal waste incinerated without energy use.

Total GHG non-biogenic and biogenic emissions reported in category 5.C from waste incineration without energy recovery were 10.03 Gg of CO<sub>2</sub> eq. in 2023. The share of emissions in this category originated from the biogenic waste incineration (0.66 Gg of bio-CO<sub>2</sub>). Disaggregation of other waste (non-MSW, clinic and other) to biogenic and non-biogenic waste is shown in [Table 7.20](#).



**Table 7.20: Activity data and emissions from waste incineration without energy recovery reported in the Waste sector in particular years**

YEAR	EMISSIONS FROM ISW INCINERATION WITHOUT ENERGY RECOVERY							
	BIOGENIC – OTHER (CRT 5.C.1.a)				NON-BIOGENIC – OTHER (CRT 5.C.1.b)			
	Amount	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	Amount	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
	kt	Gg			kt	Gg		
1990	7.11	2.4233	0.2364	0.0004	19.59	6.6820	0.6520	0.0011
1995	6.99	2.3972	0.2324	0.0004	19.48	6.6854	0.6482	0.0011
2000	7.63	2.6066	0.2538	0.0004	21.11	7.2124	0.7023	0.0012
2005	7.81	3.5918	0.2755	0.0005	38.00	17.4699	1.3399	0.0022
2010	6.29	2.5974	0.2254	0.0004	25.26	10.4329	0.9052	0.0015
2011	6.64	2.5518	0.2350	0.0004	22.02	8.4635	0.7793	0.0013
2012	4.74	1.8243	0.1682	0.0003	17.67	6.8015	0.6271	0.0010
2013	4.13	1.8380	0.1475	0.0002	18.50	8.2273	0.6602	0.0011
2014	4.25	1.7582	0.1516	0.0003	16.45	6.8123	0.5873	0.0010
2015	5.77	2.4101	0.2071	0.0003	23.15	9.6759	0.8316	0.0014
2016	3.36	1.1330	0.1153	0.0002	7.67	2.5838	0.2629	0.0004
2017	3.21	0.9987	0.1089	0.0002	7.09	2.2016	0.2400	0.0004
2018	4.28	1.6241	0.1503	0.0003	14.22	5.3971	0.4996	0.0008
2019	3.95	1.5125	0.1395	0.0002	13.79	5.2823	0.4872	0.0008
2020	3.61	1.3277	0.1261	0.0002	11.48	4.2226	0.4012	0.0007
2021	3.20	1.0150	0.1080	0.0002	7.10	2.2553	0.2400	0.0004
2022	3.55	1.2400	0.1222	0.0002	9.47	3.3050	0.3256	0.0005
2023	1.72	0.6601	0.0583	0.0001	6.20	2.3850	0.2106	0.0004

### MSW CRT 5.C.1.a(b).i (Biogenic and Non-Biogenic)

Activity data, as well as the detailed methodology for this source, is reported in the Energy sector, as there is no MSW incineration without energy utilisation in the Slovak Republic.

The amount of incinerated MSW is published by the ŠÚ SR since 1993. There are two large municipal waste incinerators in the country, in Bratislava and Košice. The MSW incinerator in Bratislava was put into operation in 1978 with a significant modernisation in 2002. Currently installed capacity is 135 Gg/y, the incinerator can be characterised as a continuously operated stoker. The MSW incinerator in Košice with a capacity of 80 Gg/yr was put in full operation in 1992, modernised to achieve compliance with emission standards in 2005 and reconstructed (boiler replacement and electricity generation) in 2014. Both incineration plants generate heat (steam) and electricity. For this reason, the CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions from MSW incineration are included completely in the Energy sector, category 1.A.1.a Public electricity and heat production.

Activity data on incinerated MSW are based on input from individual incinerators. No municipal waste was incinerated without energy recovery.

### Uncertainties

The default IPPC uncertainties for activity data consistent with the Energy sector were used.

### Source-specific recalculations

Please see [Chapter 7.7.1](#) for recalculations.

## Non-MSW CRT 5.C.1.a(b).ii (Biogenic and Non-Biogenic)

The non-MSW category has undergone significant changes since 1990. The key drivers of these changes were stricter legislation, the new standards (EU approximation) and the commercialisation of waste services. This led to replacing small incineration units in factories and hospitals by regional incinerators. In addition, existing large incinerators were modernised to comply with the new standards or were decommissioned.

From the total non-MSW incinerators and co-incineration plants, only a few have incineration without energy use and can be reported here. There are seven facilities incinerating hospital waste and other waste (not categorised). Sludge from industrial waste treatment was reported in this category back to the year 2012 (no sewage sludge was incinerated without energy recovery). Amounts of various types of incinerated waste included in this category are in [Table 7.21](#).

**Table 7.21:** Activity data of included types of waste from waste incineration without energy recovery reported in the waste sector in particular years

YEAR	WASTE TYPES*																
	02	03	04	05	06	07	08	09	11	12	13	14	15	16	17	18	19
Unit	kilotons																
1990	0.00	0.01	0.02	0.17	NO	NO	NO	0.00	NO	NO	0.49	NO	NO	5.51	1.55	5.51	1.55
1995	0.00	0.01	0.02	0.16	NO	NO	NO	0.00	NO	NO	0.52	NO	NO	5.45	1.53	5.45	1.53
2000	0.00	0.01	0.02	0.18	NO	NO	NO	0.00	NO	NO	0.53	NO	NO	5.93	1.67	5.93	1.67
2005	0.01	0.00	0.00	0.20	0.10	0.43	0.15	NO	NO	0.02	0.82	0.00	0.58	5.25	7.05	5.25	0.22
2010	0.00	0.00	NO	NO	NO	0.03	0.08	0.00	NO	NO	0.11	0.11	0.32	3.54	5.56	3.54	0.00
2011	0.00	0.00	NO	0.01	0.00	0.09	0.02	0.00	0.00	0.00	0.11	0.06	0.23	4.50	3.69	4.50	0.01
2012	0.01	NO	NO	NO	0.02	0.02	0.04	NO	NO	NO	0.06	0.04	0.79	2.71	3.42	2.71	0.00
2013	0.00	0.00	NO	0.20	0.00	0.01	0.14	0.01	NO	NO	0.12	0.07	0.08	2.58	3.84	2.58	NO
2014	0.01	NO	NO	0.00	0.00	0.02	0.05	0.01	NO	NO	0.08	0.12	0.07	2.65	3.33	2.65	NO
2015	0.00	NO	NO	NO	NO	0.11	0.02	0.00	NO	0.00	0.14	0.00	0.04	3.29	5.21	3.29	0.00
2016	0.00	NO	NO	NO	NO	0.02	0.00	0.00	NO	0.00	0.15	0.02	0.06	2.82	0.21	2.82	NO
2017	0.00	NO	NO	NO	NO	0.01	0.01	0.00	NO	0.00	0.15	0.02	0.53	2.55	0.00	2.55	0.00
2018	0.00	0.01	NO	0.00	0.00	0.01	0.02	0.00	0.05	0.04	0.17	0.01	0.14	2.87	2.32	2.87	0.15
2019	0.01	0.00	NO	0.00	0.00	0.01	0.02	0.00	0.04	0.12	0.13	0.01	0.15	2.47	2.51	2.47	0.12
2020	0.00	0.00	NO	0.00	0.00	0.02	0.02	0.00	0.08	0.13	0.12	0.02	0.12	2.49	1.65	2.49	0.12
2021	0.00	NO	NO	0.00	0.00	0.01	0.00	0.00	0.03	0.08	0.14	0.02	0.10	2.53	0.18	2.53	0.18
2022	0.00	NO	NO	0.00	0.00	0.01	0.01	0.00	0.02	0.08	0.20	0.00	0.16	2.82	0.67	2.82	0.07
2023	0.00	NO	NO	0.00	0.00	0.08	0.02	0.01	0.03	0.24	0.17	0.04	0.15	1.44	0.43	1.44	0.08

\*types of waste are following European waste catalogue classification established in Commission Decision 2000/532/EC

## Methodological issues

Emissions from non-MSW are estimated by the IPCC 2019 Refinement, tier 2a approach using country specific data on waste generation and composition. Emissions of CO<sub>2</sub> were estimated using the amount of waste incinerated divided into groups of waste ([Table 7.22](#)), for each one, the specific parameters such as dry matter content, fossil carbon fraction, oxidation factor and degradable components were determined using *Equation 5.1* of IPCC GL 2006 in Chapter 5.2.1.1. Then the calculations were repeated for selected waste groups containing non-biogenic waste to estimate emissions of non-biogenic origin and biogenic waste to estimate emissions of biogenic origin.

**Table 7.22: Parameters to calculate emissions of CO<sub>2</sub>**

WASTE TYPE*	UNIT	DRY MATTER	C-FRACTION	FOSSIL C-FRACTION	FCF	OXIDATION FACTOR	DOC
01	%	0.9	0.04	0.03	0.03	1	0.01
02		0.625	0.29	0	0	1	0.29
03		0.9	0.41	0.01	0.01	1	0.4
04		0.8	0.4	0.16	0.16	1	0.24
05		1	0.8	0.8	0.8	1	0
06		1	0.8	0.8	0.8	1	0
07		1	0.8	0.8	0.8	1	0
08		1.00	0.8	0.8	0.8	1	0
09		0.9	0.04	0.03	0.03	1	0.01
10		0.9	0.04	0.03	0.03	1	0.01
11		0.9	0.04	0.03	0.03	1	0.01
12		0.9	0.04	0.03	0.03	1	0.01
13		1	0.8	0.8	0.8	1	0
14		1	0.8	0.8	0.8	1	0
15		0.9	0.04	0.03	0.03	1	0.01
16		0.9	0.04	0.03	0.03	1	0.01
17		1	0.24	0.2	0.2	1	0.04
18		0.6	0.4	0.24	0.24	1	0.16
19 without ind. sludge		0.9	0.04	0.03	0.03	1	0.01
Industrial sludge	0.22	0.8	0.91	0.71	1	0.09	

\*types of waste are following European waste catalogue classification established in Commission Decision 2000/532/EC

Data on non-MSW incineration are available from 2005 in the NEIS database (more information in the Energy sector). Data for the period 1990 – 2004 were extrapolated using surrogate data, the trend of the impact of air pollution on the forests. Data for sewage sludge incinerated were taken from the calculation as it was confirmed by the producer of the statistics (VÚVH-Water Research Institute), that there is no sewage sludge incinerated without energy recovery. Industrial sludge data are collected by the MŽP SR. Historical data for wastewater treatment sludge is not in a format compatible with the data after 2002, as in this year Slovakia implemented the European Waste Catalogue methodology. Therefore, emissions from sludge for the period 1990 – 2001 are considered as not estimated.

Activity data allow disaggregation into incineration with and without energy use appropriately. The same activity data were used for GHG inventory and Air pollutants inventory. Consistency of the time series was ensured by using the same activity data source for the whole time series. For the estimation of emissions of CH<sub>4</sub> and N<sub>2</sub>O, the tier 1 method was used using country specific data on waste generation. The emission factor for batch type incineration – stoker (Table 5.3, Chapter 5.4.2 of the IPCC 2006 GL) was used to estimate CH<sub>4</sub> emissions. For N<sub>2</sub>O emissions, the emission factor was taken for Industrial waste from the IPCC 2006 GL (Table 5.6, Chapter 5.4.3).

### Uncertainties

The default IPCC uncertainties for activity data were used. The total uncertainty of emissions from the incineration of waste was estimated to be ±45%.

**Table 7.23: Uncertainties for waste incineration**

ACTIVITY DATA AND EMISSION FACTORS	UNCERTAINTY RANGE
Incinerated waste	±5%
Dry matter content ( <i>dm</i> )	±11%
Carbon fraction (CF)	±20%
Oxidation factor	±10%

ACTIVITY DATA AND EMISSION FACTORS	UNCERTAINTY RANGE
<b>EMISSION FACTORS: Calculated as average</b>	
CO <sub>2</sub>	±32%
CH <sub>4</sub>	±50%
N <sub>2</sub> O	±100%

### Category-specific recalculations

Emissions of CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> for the category Waste Incineration – industrial waste was recalculated in this submission due to inclusion of waste incineration in clinical waste incinerators. The waste composition was taken from the NEIS database, where the operators reports all the types of fuels used for incineration ([Table 7.7](#)).

Sewage sludge was taken from the calculation after discussion with the activity data producer (VÚVH), which confirmed that sewage sludge is incinerated only with energy recovery (in biogas facilities reported in the Energy sector).

#### 7.7.2. Open Burning of Waste (CRT 5.C.2)

Open burning of waste is prohibited by the law in the Slovak Republic; therefore, this category is reported as not occurring.

## 7.8. WASTEWATER TREATMENT AND DISCHARGE (CRT 5.D)

Wastewater can be a direct source of methane (CH<sub>4</sub>) when treated or disposed anaerobically (digesters, latrines, lagoons...) or indirect source, when dissolved CH<sub>4</sub> enters aerated systems (activated sludge systems, rivers, lakes...). Wastewater can also be a direct source of nitrous oxide (N<sub>2</sub>O) when advanced treatment systems (biological nitrogen removal) are used at centralised aerobic WWTPs. Indirect source of N<sub>2</sub>O is also produced when treated or untreated wastewater with nitrogen content enters natural water sources (rivers, lakes...). This 5.D category reported emissions (CH<sub>4</sub> and N<sub>2</sub>O) from domestic (5.D.1) and industrial wastewater (5.D.2), which are generated during wastewater treatment processes or after discharging treated or untreated wastewater to the watercourses.

In the Slovakian inventory of the category 5.D, in line with the 2019 Refinement to the IPCC GL, CH<sub>4</sub> and N<sub>2</sub>O direct emissions from modern wastewater treatment plants (WWTPs) and CH<sub>4</sub> and N<sub>2</sub>O indirect emissions from treated and untreated wastewater discharge in the environment are included. CO<sub>2</sub> emissions were not estimated, as they are of biogenic origin.

The typical distribution of wastewater pollution pathways for domestic and industrial wastewater in Slovakia in the year 2023 is presented on [Figure 7.6](#). According to this figure a total generated wastewater is divided to two basic streams: collected and uncollected wastewater. Collected wastewater is further divided into domestic wastewater ([Chapter 7.8.1](#)) from households but also from small urban industrial companies mixed and discharged together into the city sewer system and then treated at the central WWTP. Balance data (flow, COD, BOD<sub>5</sub>, SS, N, P...) from these WWTPs are available annually from the SHMI database. The second collected wastewater stream is industrial wastewater separately discharged and treated at 66 large industrial WWTPs (see [Chapter 7.8.2](#)), from which we also have input and output pollution data (flow, COD, BOD<sub>5</sub>, SS, N, P...).

For the purposes of calculating the balance of GHGs from the both collected wastewater group, we have all the necessary balance data for the individual line groups ([Figure 7.6](#)):

A – treated discharge from individual industrial WWTPs

B – discharge from collected but untreated streams of industrial wastewater

C – discharge from collected but untreated streams of domestic (municipal) wastewater

D - treated discharge from domestic (municipal) WWTPs

T – emission production from treatment process in the frame of WWTP.

The situation is a bit more complicated with uncollected wastewater. We have minimal information about the amount and method of management of this wastewater stream. Uncollected wastewater is divided into two main streams: individual domestic wastewater management by accumulation in cesspools (line F on [Figure 7.7](#)) and individual wastewater treatment in small (<50 p.e.) domestic wastewater treatment plants (line H on [Figure 7.7](#)) The third negligible stream of wastewater ends up in latrines, dry toilets or other unspecified streams (line G on [Figure 7.7](#)).

It must be noticed that cesspools, according to Slovak water legislation, that are just short-term retention tanks dislike septic tanks, are in use in Slovakia. An important part of material from these tanks are collected and transported by trucks to WWTP (estimate 45% - see [Figure 7.7](#) stream from F-box to aerobic treatment of collected wastewater), whereas the remaining fraction is discharged to soils (25%) or rivers (30%), see E- and F-lines on [Figure 7.7](#).

For the purposes of calculating the balance of GHG emissions from the uncollected wastewater group, we estimate the necessary balance data for the line groups ([Figure 7.7](#)) on the base of specific population pollution – 60 g BOD<sub>5</sub>/cap. day):


E – untreated (illegal) discharge from cesspools to rivers

F – untreated (illegal) discharge from cesspools to soil

G – discharge to latrines

H - treated discharge from small domestic WWTPs and direct emissions from treatment process (< 50p.e.).

**Figure 7.7: The typical balance of wastewater pathways for domestic and industrial wastewater in Slovakia**



Total methane emissions from wastewater treatment sector were 3.325 Gg in 2023 and this value was produced dominantly from domestic WW (95.85%). Compared to the previous year, methane emissions continue slowly to decrease (2.4%), which is caused mainly by lower amounts of the population connected to cesspool tanks, which are the dominant producer of methane from Slovak wastewater.

Total N<sub>2</sub>O emissions from wastewater treatment were 0.353 Gg in 2023, which represents relatively stable emissions production between 2019 – 2023 years. In the long term, there is a slight increase in N<sub>2</sub>O emissions, which is mainly due to the increase in the share of large WWTP with nitrogen removal and its subsequent emission into the air within the terms of 2019 Refinement to the 2006 IPCC Guidelines for NGGI. In the industrial WWTPs relatively very small but a continuously decreasing trend of N<sub>2</sub>O emissions is recorded in all monitored years. **Table 7.24** shows trends of emissions from domestic and industrial wastewater during the last years.


**Table 7.24: GHG emissions in individual categories in wastewater handling in 1990 – 2023**

YEAR	DOMESTIC WASTEWATER				INDUSTRIAL WASTEWATER			
	BOD IN OUTPUT*	CH <sub>4</sub>	NITROGEN IN OUTPUT*	N <sub>2</sub> O	COD IN OUTPUT*	CH <sub>4</sub>	NITROGEN IN OUTPUT*	N <sub>2</sub> O
	Gg							
1990	108.76	5.208	54.15	0.2486	46.75	1.286	4.435	0.035
1995	79.65	4.994	39.60	0.1912	33.81	0.930	3.669	0.029
2000	73.13	4.877	34.45	0.1807	29.04	0.798	2.905	0.023
2005	59.20	4.379	22.09	0.2101	16.88	0.464	1.902	0.015
2010	51.41	4.014	19.60	0.2220	13.39	0.368	1.671	0.023
2015	43.81	3.651	17.56	0.2440	8.81	0.242	0.745	0.016
2016	41.53	3.541	17.53	0.2515	8.90	0.245	0.829	0.018
2017	39.55	3.547	17.73	0.2701	8.48	0.233	0.788	0.029
2018	38.54	3.571	17.21	0.2730	7.18	0.198	0.624	0.013
2019	37.66	3.530	17.34	0.3074	7.48	0.206	0.595	0.013
2020	37.04	3.489	17.25	0.3217	6.59	0.181	0.536	0.012
2021	35.36	3.382	15.97	0.2988	6.89	0.189	0.624	0.011
2022	34.69	3.264	16.35	0.3116	5.68	0.156	0.514	0.010
2023	34.30	3.187	15.73	0.3432	5.02	0.138	0.496	0.010


\*In output means all sources from collected and uncollected wastewater participated on CH<sub>4</sub> or N<sub>2</sub>O production.

The distribution of methane and N<sub>2</sub>O emissions from domestic and industrial wastewater in Slovakia is presented on **Figures 7.8** and **7.9**.

**Figure 7.8: Distribution of the methane emissions (in Gg) from domestic and industrial wastewater**



**Figure 7.9:** Distribution of the N<sub>2</sub>O emissions (in Gg) from domestic and industrial wastewater



### 7.8.1. Domestic Wastewater (CRF 5.D.1)

In 2023, 71.55% of the Slovak population was connected to public sewage systems and the rest is using cesspool tanks or individual treatment systems. Wastewater collection and treatment in Slovakia is developing toward modern, advanced WWTPs. Totally 782 domestic and municipal WWTPs treat yearly about 445 Mil. m<sup>3</sup> of wastewater and 76.6% is treated in tertiary level (with nitrogen and phosphorus removal).

All these treatment plants operate in a standard technological line, where raw wastewater undergoes mechanical pre-treatment and then enters the biological stage in the aerobic mode of activated sludge. The result of the process is treated wastewater that goes to the surface recipient - rivers. Balance data (flow, COD, BOD<sub>5</sub>, SS, N, P..) from these WWTPs are available annually from the SHMI database.

A by-product of the treatment process is excess sludge, which subsequently undergoes a stabilization process. The legislation and practice in wastewater treatment in Slovakia require that sewage sludge must be stabilised directly at the wastewater treatment plant (e.g. Act No 188/2003 Coll. requires that only stabilised sewage sludge can be further processed outside of WWTP i.e. on compost, in incinerator etc.). Thus, according to the Slovak Technical Norm 75 6401 "Sewage Treatment Plants" the small and middle size domestic WWTPs (about 730 plants with capacity obviously lower than 10 000 p.e.) are used an aerobic sludge stabilisation (long term aeration) with sludge retention time (SRT) higher than 25 d. During the stay at WWTPs, this sludge does not go into a state of anaerobic digestion (methanation), i.e. the formation of methane is negligible here.

The largest domestic WWTPs (ca. 50 WWTPs each with a capacity of more than 10 000 p.e.) generate about 76% of total Slovak sewage sludge production. These large WWTPs are processing excess sludge by anaerobic digestion (in digester towers) way with biogas (CH<sub>4</sub> + CO<sub>2</sub>) production. The produced biogas is captured in a gas holder, from where it is discharged for combustion in boilers. The result of combustion is either only the heat produced for heating the boiler (approx. 30 WWTPs) or the production of heat and electricity (23 WWTPs) in cogeneration units (CHP). The produced heat is used to heat the digesters and the produced electricity is dissipated into the public electricity grid. The entire biogas system at WWTP is strictly monitored and is maximally protected against leaks of unburned biogas into the air. Therefore, it can be stated that free biogas leaks into the air are under the threshold of significance at large WWTPs, which is why they are not even included in this chapter. All data on produced biogas, heat and electricity from domestic WWTPs biogas are reported in [Chapter 7.6.3](#).

The amount of produced (aerobic and anaerobic) sludge from small and large WWTPs and their further process ways are summarised in [Table 7.25](#).

**Table 7.25: Distribution of the sludge from domestic WWTPs (data from the WRI)**

YEAR	TOTAL GENERATED	TOTAL USE	DIRECT AGR. LAND APPLIC.	COMPOSTED	INCINER.	LANDFILLED	TEMPORARY STORED ON-SITE
	tons						
1990	55 000	45 207	-	-	-	-	-
1995	55 000	45 207	-	-	-	-	-
2000	56 279	35 358	-	-	-	13 796	7 125
2005	56 360	39 120	-	-	-	8 530	8 710
2010	54 760	48 063	923	47 140	-	16	6 681
2015	56 242	51 602	NO	34 689	16 913	1 709	2 932
2016	53 054	45 738	NO	34 695	11 043	2 359	4 957
2017	54 517	46 654	NO	34 416	12 238	2 636	5 227
2018	55 929	44 659	NO	32 982	11 677	2 451	8 819
2019	54 832	45 149	NO	32 217	12 932	2 296	7 387
2020	55 519	48 490	NO	36 562	11 928	2 302	4 727
2021	54 764	50 042	NO	37 289	12 753	456	4 266
2022	55 049	43 835	NO	33 509	10 326	1 540	9 674
2023	56 420	46 747	NO	31 050	15 697	490	9 183

Total methane emissions from domestic wastewater were 3.187 Gg in 2023. The main contributions to these emissions have retention tanks (cesspools) with 1.601 Gg as well as treatment processes with 1.518 Gg in 2023, which summarily represents about 92% of methane emissions from domestic wastewater (**Tables 7.26**).

**Table 7.26: Summary of methane emissions from the domestic WW by pathways in particular years (wastewater stream letters C-D-E-F-G-H correspond with streams on Figure 7.7, T – represents methane emissions streams both from treatment processes in centralised and small domestic WWTPs).**

PATHWAY	DOMESTIC WW TREATED AND UNTREATED	TREATMENT PROCESS IN WWTPS	UNTREATED DISCHARGE FROM CESSPOOLS TO RIVERS OR SOILS		IN AIR FROM CESSPOOLS TANKS	REST/ UNCATERO RISED	DISCHARGE FROM DOMESTIC WWTPs
	C+D	T	E - rivers	E - soils	F	G	H
MFC	0.035	0.03	0.035	0.1	0.05	0.1	0.035
YEAR	CH <sub>4</sub> in Gg						
1990	1.084	0.824	0.295	0.842	1.264	0.900	0
1995	0.502	1.274	0.294	0.841	1.263	0.819	0
2000	0.412	1.380	0.294	0.841	1.263	0.687	0
2005	0.190	1.306	0.283	0.807	1.212	0.578	0.002
2010	0.114	1.369	0.241	0.690	1.150	0.445	0.006
2011	0.113	1.369	0.239	0.682	1.137	0.332	0.006
2012	0.101	1.246	0.236	0.675	1.124	0.323	0.007
2013	0.103	1.284	0.234	0.667	1.112	0.258	0.007
2014	0.088	1.032	0.231	0.660	1.099	0.206	0.008
2015	0.083	1.399	0.228	0.652	1.087	0.194	0.008
2016	0.066	1.391	0.226	0.645	1.074	0.132	0.008
2017	0.058	1.503	0.223	0.637	1.062	0.056	0.009
2018	0.055	1.579	0.220	0.630	1.049	0.029	0.009
2019	0.056	1.589	0.216	0.618	1.030	0.011	0.010
2020	0.057	1.584	0.212	0.607	1.011	0.008	0.010
2021	0.054	1.563	0.203	0.580	0.966	0.006	0.010




PATHWAY	DOMESTIC WW TREATED AND UNTREATED	TREATMENT PROCESS IN WWTPS	UNTREATED DISCHARGE FROM CESSPOOLS TO RIVERS OR SOILS		IN AIR FROM CESSPOOLS TANKS	REST/UNCATEGORISED	DISCHARGE FROM DOMESTIC WWTPs
	<i>C+D</i>	<i>T</i>	<i>E - rivers</i>	<i>E - soils</i>	<i>F</i>	<i>G</i>	<i>H</i>
<i>MFC</i>	<i>0.035</i>	<i>0.03</i>	<i>0.035</i>	<i>0.1</i>	<i>0.05</i>	<i>0.1</i>	<i>0.035</i>
<b>YEAR</b>	<i>CH<sub>4</sub> in Gg</i>						
<b>2022</b>	0.047	1.470	0.201	0.574	0.956	0.005	0.010
<b>2023</b>	0.053	1.518	0.197	0.468	0.936	0.005	0.011

The new calculation of CH<sub>4</sub> emissions according to IPCC 2019 Refinement caused significant changes in both partial and total emission values. The IPCC 2019 Refinement (Table 6.3 updated) set many new MCF default values for the calculation procedure, as was stated in previously year report (2022). This year was recalculated CH<sub>4</sub> emissions from cesspools. The recalculations were based on the different MCFs used for septic tanks and cesspools. Our previous calculations for CH<sub>4</sub> production used an MCF value = 0.4 (lower value in the range of line for septic tank in Table 6.3 updated in 2019 Refinement to the 2006 IPCC Guidelines for NGGI). However, the operation of cesspools in Slovakia (and everywhere in the world) works in such a way that the contents of cesspools are regularly exhausted at intervals of 1 – 1.5 months. During that time, methane production cannot develop to the same extent as in septic tanks, where the sludge emptied interval is about 1-1.5 years. After consultations with several experts, we proceeded to adjust the MCF for septic tanks to the conditions in the cesspool. As stated in the previous text, the production of methane from cesspools in this year's inventory is divided into four separate streams and the MCF corresponding to them:

- MCF = 0.05 for methane production direct in cesspool tanks: SRT in cesspools is ca. 10 times shorter compared to septic tanks (1-1.5 year/1-1.5 months) then MCF for septic tanks/10 = 0.5/10 = 0.05
- MCF = 0.035 for methane production from the part cesspool stream which is legally transported to nearby WWTPs as part of treated wastewater in centralised WWTP (as recommended in Table 6.3 (updated) of the 2019 Refinement to the 2006 Guidelines for discharge to aquatic environment other than reservoirs, lakes and estuaries). This part of cesspool wastewater (as TOW) is already included in influent and effluent pollution data form individual WWTPs in yearly database from SHMI, i.e. is not separately evaluated in methane emissions
- MCF = 0.35 for methane production of wastewater from cesspools to rivers (as recommended in Table 6.3 (updated) of the 2019 Refinement to the 2006 Guidelines for discharge to aquatic environment other than reservoirs, lakes and estuaries). This part of production is integrated as a new column (E – rivers in Table 7.31).
- MCF = 0.1 for methane production of wastewater from cesspools to soil. This value of MCF is compromising estimation from 2019 Refinement to the 2006 GL (Table 6.3. Updated) and 2006 IPCC GL Volume 5, Chapter 4, Table 4.1.). This part of production is integrated as a new column (E – soils in Table 7.31).

All these cesspool pollution data (TOW) are calculated on the base of estimated population connected on cesspools with specific population equivalent (p.e. = 60 gBOD<sub>5</sub>/cap. day).

**Figure 7.10: Distribution of the domestic wastewater methane emissions (in Gg)**




As can be seen from [Table 7.26](#) and from [Figure 7.10](#), methane emissions from effluents into rivers decreased significantly. A similar trend is evident also in methane emissions from cesspools, even though it is still the largest source of methane from wastewater. A slightly increasing trend can be observed for emissions arising during the treatment process on WWTPs. This is due to the gradual increasing of connected population to public sewerage systems and to new WWTPs.

Total N<sub>2</sub>O emissions from domestic wastewater treatment were 0.343 Gg. The minority of N<sub>2</sub>O emissions is generated both from WWTPs untreated (0.0224 Gg) and treated discharges (0.0521 Gg), on the other hands, dominant producer is treatment process on WWTPs with 75.3% on total emissions ([Table 7.27](#)).

**Table 7.27: Summary of N<sub>2</sub>O emissions from the domestic WW by pathways in particular years**

YEAR	UNTREATED DISCHARGE AND RETENTION TANKS	DIRECT FROM TREATMENT PROCESS TO AIR	TREATED DISCHARGE	TOTAL
	N <sub>2</sub> O in Gg			
1990	0.1288	0.0000	0.1198	0.2486
1995	0.0762	0.0000	0.1150	0.1912
2000	0.0693	0.0148	0.0693	0.1807
2005	0.0417	0.1089	0.0595	0.2101
2010	0.0320	0.1334	0.0566	0.2220
2011	0.0319	0.1311	0.0561	0.2191
2012	0.0303	0.0978	0.0465	0.1746
2013	0.0291	0.1227	0.0523	0.2041
2014	0.0269	0.0826	0.0400	0.1495
2015	0.0263	0.1626	0.0552	0.2440
2016	0.0258	0.1704	0.0553	0.2515
2017	0.0252	0.1862	0.0587	0.2701
2018	0.0246	0.1920	0.0564	0.2730
2019	0.0246	0.2256	0.0573	0.3074
2020	0.0244	0.2400	0.0573	0.3217
2021	0.0236	0.2252	0.0500	0.2988
2022	0.0231	0.2346	0.0539	0.3116
2023	0.0224	0.2686	0.0521	0.3432

**Figure 7.11: Distribution of the domestic wastewater N<sub>2</sub>O emissions (in Gg)**



As is it evident from [Table 7.27](#) and from [Figure 7.11](#), nitrous oxide emissions from both treated and untreated effluents into rivers decreased continuously. A significantly increasing trend can be observed for emissions arising from the biological treatment process. This is due to the gradual implementation of nitrification and denitrification processes in new as well as existing WWTPs. It is a new parameter in N<sub>2</sub>O emission reports, which completely change emission trends in the historical content. Similar to what was stated with methane emissions, a new calculation of N<sub>2</sub>O emissions according to IPCC 2019 Refinement caused significant changes in both partial and total emission values.

### Methodological issues

The IPCC 2019 Refinement to the 2006 IPCC 2006 GL method was accommodated to reflect new available data and observed trends in wastewater management. Known influent and effluent BOD from all individual domestic WWTPs (evidence database from SHMÚ) was used in emissions estimation from WWTPs instead of calculating a difference between theoretical total organics on input (TOW from population equivalent) and organic component removed with sludge (evidence data from VÚVH). At present, we still feel a lack of information about individual treatment systems (cesspools and domestic WWTPs), so emissions for these systems have been calculated on the basis of the estimated number of inhabitants using these systems.

N<sub>2</sub>O emissions calculation is based on the IPCC 2019 Refinement, but due to the increased number of advanced WWT plants, recommended nitrogen removal by nitrification/denitrification had to be included in the calculation. The effectiveness of N removal in WWTPs was adjusted according Table 6.10c (IPCC 2019 Refinement). According to the information from the VÚVH, measurements of nitrogen content in sludge was provided also in 2023 (46.7 k/kg TS).

Default parameters and emission factors from the IPCC 2019 Refinement were used for CH<sub>4</sub> and N<sub>2</sub>O emissions estimation of domestic wastewater. Default value 0.6 kg CH<sub>4</sub>/kg BOD was used for the maximum CH<sub>4</sub> producing capacity (B<sub>0</sub>). Default value 0.035 for methane correction factor (MCF) was used for all pathways except for retention tanks (cesspools) where MCF=0.05 was applied. MCF for direct emissions from treatment processes was used 0.03.

Identification of wastewater pathways is based on population using individual pathways. Estimation of CH<sub>4</sub> emissions from collected domestic wastewater is based on BOD measured data on really generated discharged pollution to watercourses from public sewers. Emissions of CH<sub>4</sub> from cesspool tanks, dry toilets, small domestic WWTPs and from untreated discharge from public sewers were estimated based on population and BOD<sub>5</sub> per person per day (60 g – country-specific value).

## Uncertainties

The default uncertainties based on the IPCC 2019 Refinement were used and adjusted (where possible) to assess emissions estimation and to reflect country-specific data or circumstances. The calculation of methane emissions was based on real pollution data (BOD<sub>5</sub>) at the output of existing WWTPs. Emissions from individual treatment systems (septic tanks) were defined on an estimate of the number of inhabitants connected to these facilities. However, the operation of these individual installations is outside the central evidence and therefore emissions from this group are burdened with very high uncertainty.

The list of the most significant emission factors and their uncertainty range is given in [Table 7.28](#). To define the total uncertainty of emissions for methane or N<sub>2</sub>O is relatively complicated, as the total uncertainty should be defined as the conjunction of the all-individual uncertainties entering into the final emission calculation. Based on expert estimates and discussions, a value of ±15% was defined as the overall uncertainty for methane emissions and a value of ±25% was defined for N<sub>2</sub>O emissions.

**Table 7.28: Uncertainties for the category of domestic wastewater treatment**

EMISSION FACTORS AND ACTIVITY DATA	UNCERTAINTY RANGE
<b>Emission factors</b>	
<u>For methane calculation:</u>	
EF <sub>1</sub> (kg CH <sub>4</sub> /kg BOD) = 0.6 (default value)	±10%
MCF for treated and untreated system = 0.1 (default value)	±10%
MCF for cesspools systems = 0.05 (default value)	±20% (temperature depend)
<u>For N<sub>2</sub>O calculation:</u>	
N <sub>2</sub> O Emission factor effluent = 0.005 (default value)	±10%
<b>Activity data</b>	
<u>For methane calculation:</u>	
TOW from operational WWTPs influent and effluent (SHMÚ data)	±10% (sampling and analytical errors)
BOD per person and day (for septic tanks) = 60 g/person per day	±30%
Human population distribution (collected, uncollected)	±5%
<u>For N<sub>2</sub>O calculation:</u>	
N <sub>eff</sub> from real WWTPs influent and effluent (SHMÚ data)	±10% (sampling and analytical errors)
Protein annual consumption (ŠÚ SR data)	±5%
N in sludge (VÚVH data)	±10% (sampling and analytical errors)

## Category-specific recalculations

Estimation of CH<sub>4</sub> and N<sub>2</sub>O emissions from domestic wastewater sector were calculated using a new calculation methodology according to the 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories already in previous year report. There were many changes on individual level in the calculated CH<sub>4</sub> and N<sub>2</sub>O emission data caused by a change of the default MCF and EF values for discharge from treated and untreated systems, for wastewater treatment systems emissions, by efficiency of BOD<sub>5</sub>, COD, N<sub>tot</sub> removal in primary, secondary and tertiary treatment systems, etc. This year were recalculations realised only by changing of MCF for cesspools as above stated.

Detailed comments and final comparison of the previous inventory and re-calculated time-series are presented in the [Chapter 7.3](#). Recalculation data for methane emissions from domestic wastewater are recorded in the [Table 7.8](#).

### 7.8.2. Industrial Wastewater (CRF 5.D.2)

In 2023, 66 individual plants were included in the group of industrial wastewater treatment plants, which processed a total of 91.45 million m<sup>3</sup> of wastewater with a total annual pollution of 39.57 kt COD (903,400 p.e.). In addition, part of the industrial wastewater (from smaller industrial plants) was discharged into the municipal sewers and treated at municipal wastewater treatment plants together with collected domestic wastewater from households (emissions from this part of industrial wastewater is incorporate in [Chapter 7.8.1](#). Domestic wastewater)

Water consumption for industrial purposes and resulting pollution discharge of wastewater have significantly decreased in the period 1990 – 2015, but during last year's 2018 – 2023 there was a relatively stable wastewater production and slight decrease of organic pollution (as COD and Ntot in [Table 7.29](#)).

Similar to domestic WWTPs, industrial plants also output treated water and excess sludge. Input and output data from all these plants are available every year (flow, COD, BOD, SS, N, P...) from SHMI database. Based on these data, emissions (CH<sub>4</sub> and N<sub>2</sub>O) from industrial WWTPs are calculated. In six from 66 WWTPs, the wastewater is (partially) treated in an anaerobic reactor (IC-reactor, mixed), the biogas produced is burned and used to heat the reactors. Similar to domestic wastewater treatment plants, the entire biogas system at these WWTP is strictly monitored and is maximally protected against leaks of unburned biogas into the air. Therefore, it can be stated that free biogas leaks into the air are negligible at these WWTPs, which is why they are not even mentioned in this chapter. All data on produced biogas and heat in industrial WWTPs are reported in [Chapter 7.6](#).

Until 2001, produced and removed industrial treatment sludge was reported as "NE" in the CRF table 5.D.2. In the reflection of the discussion during the UNFCCC review 2019, data about sludge production and disposal ways from industrial wastewater treatment (back from the year 2005) were processed based on the ŠÚ SR and the "IS Odpady", which is a database of waste production operated by the MŽP SR. For the purposes of this submission, the actual values of industrial sludge production have been used ([Table 7.29](#)).

**Table 7.29: Distribution of the sludge from industrial WWTPs since 2005 (data from the ŠÚ SR)**

YEAR	TOTAL GENERATED	TOTAL USE	DIRECT AGR. LAND APPLIC.	COMPOSTED	INCINER.	LANDFILLED	TEMPORARY STORED ON-SITE
	tons						
2005	10 307	5 577	2 231	1 037	1 501	785	24
2010	25 571	19 769	1 102	6 369	1 228	11 058	13
2015	11 485	7 500	813	3 248	2 496	898	45
2016	13 651	12 200	1 134	3 353	2 021	5 641	50
2017	22 211	15 538	362	3 460	1 206	1 063	9 447
2018	49 669	40 461	287	3 520	3 307	1 006	32 341
2019	12 935	9 393	49	3 361	2 663	1 327	1 993
2020	32 599	28 611	1	3 893	1 326	6 445	16 946
2021	20 724	10 992	1	3 191	1 013	6 034	750
2022	14 240	10 046	1	1 905	925	5 598	1 618
2023	16 130	10 179	5	4 419	1 417	1 657	2 683

Total methane emissions were estimated to be 0.1381 Gg and total N<sub>2</sub>O emissions were 0.0101 Gg from industrial wastewater treatment in 2023. The pathways A and B ([Figure 7.6](#)) are included in the estimation of methane emissions. [Table 7.30](#) shows the activity data and resulting emissions estimation.


**Table 7.30: GHG emissions from industrial wastewater treatment in particular years**

YEAR	TOTAL ORGANIC PRODUCT	NITROGEN IN EFFLUENT	CH <sub>4</sub>	N <sub>2</sub> O
	kt DC - COD	Gg		
1990	46.746	4.435	1.286	0.0348
1995	33.814	3.669	0.930	0.0288
2000	29.035	2.905	0.798	0.0228
2005	16.880	1.902	0.464	0.0149
2010	13.386	1.671	0.368	0.0227
2011	10.747	1.463	0.296	0.0221
2012	10.080	1.283	0.277	0.0233

YEAR	TOTAL ORGANIC PRODUCT	NITROGEN IN EFFLUENT	CH <sub>4</sub>	N <sub>2</sub> O
	<i>kt DC - COD</i>	<i>Gg</i>		
2013	9.919	1.041	0.273	0.0140
2014	9.072	0.836	0.249	0.0261
2015	8.811	0.745	0.242	0.0160
2016	8.899	0.829	0.245	0.0177
2017	8.480	0.788	0.233	0.0290
2018	7.184	0.624	0.198	0.0128
2019	7.477	0.594	0.206	0.0128
2020	6.590	0.536	0.181	0.0120
2021	6.888	0.624	0.189	0.0107
2022	5.676	0.514	0.156	0.0098
2023	5.021	0.496	0.138	0.0101


The [Figure 7.12](#) confirms significantly decrease of methane emissions both from treated as well as untreated industrial wastewater. In contrast to domestic wastewater, industrial wastewater does not consider the process of methane formation at a WWTP with activated sludge (default EF = 0 in Table 6.8. Updated in the 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories). Especially, the proportion of untreated WW is already almost negligible, which indicates a high level of industrial wastewater treatment quality.

**Figure 7.12:** Distribution of the industrial wastewater methane emissions (in Gg)



Even with N<sub>2</sub>O emissions, a significant decrease is observed in the items treated and untreated wastewater. Since 2009, balance data on the input and output of industrial WWTPs have been available. Based on these data, the emissions produced in the process of nitrification and denitrification were also calculated (similarly as in domestic wastewater chapter). These data also have a decreasing tendency, with the exception of the years 2014 and 2017, when there were extremely high TN loads in wastewater. However, the slight decrease in N<sub>2</sub>O emissions continues ([Figure 7.13](#)).

**Figure 7.13:** Distribution of the industrial wastewater N<sub>2</sub>O emissions (in Gg)



### Methodological issues

The new methodology recommended by the IPCC 2019 Refinement to the 2006 IPCC 2006 GL has been used for estimating emissions from industrial wastewater already in previous year report. The same methodology was used in this year also. COD values from individual industrial WWTPs reported by the SHMÚ were used in methane emissions estimation. It is assumed that the use of the reported COD data will provide better results than estimation according to the methodology provided in Chapter 6.4.1.3 of the IPCC 2019 Refinement to the IPCC 2006 GL. Treated and untreated pollution (as COD) from industrial wastewater discharged into rivers by separate industrial sewers were considered here as a methane emissions source. Industrial wastewater discharged to public sewers is included in domestic wastewater. Default value 0.25 kg CH<sub>4</sub>/kg COD of maximum CH<sub>4</sub> producing capacity (B<sub>0</sub>) was used. Default value 0.11 of methane correction factor (MCF) for both pathways was used (old methodology used MCF = 0.1). It is expected if anaerobic treatment of industrial wastewater was used (only three facilities), that all methane from this treatment was burned (with or without energy utilization). Unlike domestic wastewater, the new calculation methodology does not consider methane production in the activated sludge process (EF = 0).

In compare to the old methodology, the IPCC 2019 Refinement provides a slightly changed methodology for the estimation of N<sub>2</sub>O emissions from industrial wastewater. Slovakia currently collects information on produced and discharged pollution from all sources. The SHMÚ and ŠÚ SR started to publish data on nitrogen discharged to watercourses from 2009. These data allowed to develop a simple model, which estimates N<sub>2</sub>O emissions generated from the treated and untreated discharge of industrial wastewater. For emissions estimation from industrial wastewater, default emission factors based on the IPCC 2019 Refinement were used. Default value 0.005 kg N<sub>2</sub>O-N/kg N was used. Data on discharged nitrogen are available only for the period 2009 – 2023. A good correlation (0.92) was identified between the discharged TN and COD. COD was used for extrapolation of missing TN activity data in the period 1990 – 2008. Extrapolations were done separately for treated and untreated discharge.

N<sub>2</sub>O emissions from nitrification-denitrification processes were realized only for the period 2009 – 2022, when real input and output for N-loads were available. Default EF = 0.016 kg N<sub>2</sub>O/kg TN was used to calculate emissions from the cleaning process (according to IPCC 2019 Refinement).

### Uncertainties

The default uncertainties based on the IPCC 2019 Refinement to the IPCC 2006 GL were used to assess methane and N<sub>2</sub>O emissions estimation and also to reflect country-specific data or circumstances. The calculation of methane and N<sub>2</sub>O emissions was based on real pollution data (COD and TN) at the

effluent of all existing industrial WWTPs. Data on the proportion of treated and untreated industrial water were also available.

The list of the most significant emission factors and their uncertainty range is given in [Table 7.30](#). To define the total uncertainty of industrial wastewater emissions for methane or N<sub>2</sub>O is relatively complicated, as the total uncertainty should be defined as the conjunction of the individual uncertainties entering into the final emission calculation. Based on expert estimates and also based on recommendation data from the IPCC 2019 Refinement (Table 6.13 was updated), a different value of the overall uncertainty for methane N<sub>2</sub>O emissions was defined.

**Table 7.30: Uncertainties for the category of industrial wastewater treatment**

EMISSION FACTORS AND ACTIVITY DATA	UNCERTAINTY RANGE
Emission factors	
<u>For methane calculation:</u>	
Maximum CH <sub>4</sub> producing capacity (B <sub>0</sub> )	±30%
EF <sub>j</sub> (kg CH <sub>4</sub> /kg COD) = 0.25 (default value)	±10%
MCF for treated and untreated system = 0.1 (default value)	±10%
<u>For N<sub>2</sub>O calculation:</u>	
N <sub>2</sub> O Emission factor effluent = 0.005 (default value)	±30%
Activity data	
<u>For methane calculation:</u>	
Human population (P)	±5%
TOW from real industrial WWTPs effluent (SHMÚ data)	±20% (sampling and analytical errors)
<u>For N<sub>2</sub>O calculation:</u>	
N <sub>eff</sub> from real WWTPs influent and effluent (SHMÚ data)	±25% (sampling and analytical errors)

### Source-specific recalculations

Due to slight changes in the recommended calculation procedures (according to IPCC 2019 Refinement), there were changes in the resulting values of individual parameters as well as in the resulting total values. A new calculation for both methane and nitrogen emissions were carried out in the industrial wastewater sector. A detailed comparison of the previous inventory submission for time-series and new data for methane emissions are shown in [Table 7.8](#). A detailed comparison of the previous inventory submission for time-series and new data for N<sub>2</sub>O emissions are shown in [Table 7.9](#). The actual values of methane and nitrous oxide emissions for the industrial wastewater sector are shown in [Table 7.30](#) and in [Figure 7.11](#) and [Figure 7.12](#).

## 7.9. MEMO ITEMS (CRT 5.F)

The IPCC Waste Model provides estimates on stored carbon and overview of this parameter is shown in [Table 7.31](#), disaggregated to municipal solid waste and non-municipal solid waste. (Note: These data were not inserted in the CRT table 5.F, as this table requires CO<sub>2</sub> emissions, but SWDS are generating CH<sub>4</sub>. The main contradiction is that long-term stored carbon remains as carbon. Emissions in these categories were recalculated based on the changes in waste disposal composition (see [Chapter 7.5](#)).

**Table 7.31: Accumulated Long-term stored C in SWDS in particular years**

YEAR	ACCUMULATED STORED C	ANNUAL CHANGE IN STORED C	ANNUAL CHANGE IN STORED C IN HWP WASTE
	Gg		
1990	1 043.18	47.67	35.48
1995	1 244.80	42.18	28.26
2000	1 512.78	54.61	32.62
2005	1 852.94	74.89	45.90



YEAR	ACCUMULATED STORED C	ANNUAL CHANGE IN STORED C	ANNUAL CHANGE IN STORED C IN HWP WASTE
	Gg		
2010	2 295.50	97.29	59.94
2011	2 382.54	87.04	52.40
2012	2 465.10	82.57	48.73
2013	2 541.06	75.95	44.82
2014	2 617.18	76.12	44.94
2015	2 698.17	81.00	47.68
2016	2 774.17	76.00	44.85
2017	2 849.79	75.62	43.88
2018	2 920.18	70.38	39.98
2019	2 987.92	67.74	37.55
2020	3 052.49	64.57	36.23
2021	3 097.31	44.82	22.47
2022	3 139.07	41.77	20.75
2023	3 177.35	38.28	19.00

<b>CHAPTER 8. OTHER (CRT 6)</b> .....	<b>427</b>
<b>CHAPTER 9. INDIRECT CO<sub>2</sub> AND NITROUS OXIDE EMISSIONS</b> .....	<b>427</b>
<b>CHAPTER 10. RECALCULATIONS AND IMPROVEMENTS</b> .....	<b>427</b>
10.1. Explanations and Justifications for Recalculations.....	427
10.2. Implications for Emission Levels.....	428
10.3. Recalculations, Including in Response to the Review Process, and Planned Improvements to the Inventory .....	430
<b>CHAPTER 11. INFORMATION ON CHANGES IN NATIONAL SYSTEM</b> .....	<b>434</b>
<b>CHAPTER 12. INFORMATION ON MINIMIZATION OF ADVERSE IMPACTS             IN ACCORDANCE WITH ARTICLE 3, PARAGRAPH 14</b> .....	<b>434</b>
<b>REFERENCES</b> .....	<b>436</b>
<b>ANNEX 1. KEY CATEGORIES</b> .....	<b>443</b>
<b>ANNEX 2. ASSESSMENT OF COMPLETENESS</b> .....	<b>461</b>
<b>ANNEX 3. ASSESSMENT OF UNCERTAINTY</b> .....	<b>471</b>
<b>ANNEX 4. QUALITY ASSURANCE/QUALITY CONTROL PLAN</b> .....	<b>490</b>
<b>ANNEX 5. ENERGY BALANCE OF THE ŠÚ SR FOR THE YEAR 2023</b> .....	<b>494</b>

---

## CHAPTER 8. OTHER (CRT 6)

Slovakia does not report any emissions under the sector Other.

## CHAPTER 9. INDIRECT CO<sub>2</sub> AND NITROUS OXIDE EMISSIONS

The CO<sub>2</sub> resulting from the atmospheric oxidation of CH<sub>4</sub>, CO and NMVOC is referred to as indirect CO<sub>2</sub>. The IPCC 2006 GL and the IPCC 2019 Refinement provide a method how the CO<sub>2</sub> inputs from the atmospheric oxidation of NMVOC in industry can be calculated.

Indirect CO<sub>2</sub> emissions from these processes were estimated and are included in the IPPU sector. Indirect emissions were estimated in the category 2.D – Non-energy products from fuels and solvent use for the first time in this submission as reported for the time series. More information can be found in [Annex A4.4](#) of [Chapter 4](#).

Indirect N<sub>2</sub>O emissions in the Agriculture sector address nitrous oxide (N<sub>2</sub>O) emissions that result from the deposition of the nitrogen emitted as NO<sub>x</sub> and NH<sub>3</sub>. N<sub>2</sub>O is produced in soils through the biological processes of nitrification and denitrification. Indirect N<sub>2</sub>O emissions from manure management are reported in CRT table 3.B(b) as a part of N<sub>2</sub>O emissions in this category. These indirect emissions result from volatile nitrogen losses, that occur during manure collection and storage and which are diffused into the surrounding air. Nitrogen losses begin at the point of excretion in houses and other animal production areas and continue through on-site management in storage and treatment systems. Indirect N<sub>2</sub>O emissions from these sources are included in the categories 3.B(b).5 – Indirect N<sub>2</sub>O emissions from manure management.

Indirect N<sub>2</sub>O emissions from managed soils are reported in CRT table 3.D.2 – Indirect N<sub>2</sub>O emissions from managed soils as a part of N<sub>2</sub>O emissions. These emissions are from the following sources:

- the volatilization of N (as NH<sub>3</sub> and NO<sub>x</sub>) following the application of synthetic and organic N fertilizers and /or urine and dung deposition from grazing animals,
- and the subsequent deposition of the N as ammonium (NH<sub>4</sub><sup>+</sup>) and oxides of N (NO<sub>x</sub>) on soils and waters, and the leaching and runoff of N from synthetic and organic N fertilizer additions, crop residues, mineralization /immobilization of N associated with loss/gain of soil C in mineral soils through land use change or management practices, and urine and dung deposition from grazing animals.

No indirect emissions are reported in the Energy, LULUCF and Waste sectors.

## CHAPTER 10. RECALCULATIONS AND IMPROVEMENTS

### 10.1. Explanations and Justifications for Recalculations

The main driver for recalculations in the 2025 greenhouse gas inventory submission of the Slovak Republic has been the updating of the COPERT software, methodologies and the new input data. The significance of the sources based on the results of the key category and uncertainty analyses are considered when prioritising improvements to be made in the inventory calculations. No UNFCCC review was performed in 2024. The recalculations made since the previous inventory submission (2024) are described also in the appropriate sectoral [Chapters 3-7](#). The list of the major recalculations with the

short descriptions made in the 2025 submission is summarized in [Tables 10.3](#) and [10.4](#). No recommendation from the EU ESD inventory reviews (2024) have been addressed by the TERT.

## 10.2. Implications for Emission Levels

Reflecting the QA/QC activities for improving the emissions inventory of GHG and recommendations provided by the experts during the review process for inventory submissions under the UNFCCC, the experts involved in the National Inventory System of the Slovak Republic proposed the recalculations of the several subsectors and categories. The recalculations and reallocations of emissions are based on updated or revised methodologies (for agricultural and LULUCF activities and for waste categories), updated statistical information (waste and households) or based on harmonization between GHG and air pollutant input data (for the IPPU sector in solvents use). The recalculations listed in [Tables 10.1](#) and [10.2](#) were provided in CTR tables 2025, version 2 against previous inventory submission from September 15, 2024 version 2 with and without the LULUCF sector. The [Table 10.3](#) presents list of performed recalculations with the short summarizing description (detailed information is provided in the sectoral chapters in this document). Total GHG emissions without LULUCF and with indirect emissions increased after recalculations made in 2025 submission for the year 1990 by 0.08%, and for the year 2022 decreased by 0.46% ([Table 10.1](#)). Regarding total GHG emissions with LULUCF and with indirect emissions, GHG emissions decreased in 2025 submission by 0.58% for the year 2022 ([Table 10.2](#)). This comparison used the GWP taken from [AR5](#).

**Table 10.1:** Comparison of the GHG emissions trend without LULUCF and with indirect emissions in 2024 and 2025 submissions

NATIONAL GHG INVENTORY WITHOUT LULUCF WITH INDIRECT EMISSIONS			
YEAR	Submission 2024 v2	Submission 2025 v0.3	2025 v0.3/2024 v2
	Gg of CO <sub>2</sub> eq.		%
1990	73 455.32	73 513.34	100.08%
1991	64 199.80	64 177.97	99.97%
1992	58 768.01	58 198.55	99.03%
1993	55 182.38	54 642.99	99.02%
1994	52 665.58	52 095.68	98.92%
1995	53 180.07	52 617.26	98.94%
1996	53 033.33	52 465.61	98.93%
1997	52 826.91	52 244.35	98.90%
1998	52 092.98	51 505.00	98.87%
1999	50 838.56	50 282.12	98.91%
2000	48 904.08	48 360.51	98.89%
2001	51 134.01	50 631.62	99.02%
2002	49 866.18	49 323.51	98.91%
2003	50 096.54	49 648.66	99.11%
2004	50 723.59	50 318.75	99.20%
2005	50 682.48	50 336.41	99.32%
2006	50 706.80	50 266.70	99.13%
2007	48 888.15	48 398.28	99.00%
2008	49 369.45	48 797.00	98.84%
2009	45 145.99	44 567.68	98.72%
2010	45 888.76	45 476.25	99.10%
2011	44 812.42	44 373.20	99.02%
2012	42 442.53	42 014.11	98.99%
2013	42 119.84	41 690.86	98.98%
2014	40 103.47	39 688.50	98.97%

NATIONAL GHG INVENTORY WITHOUT LULUCF WITH INDIRECT EMISSIONS			
YEAR	Submission 2024 v2	Submission 2025 v0.3	2025 v0.3/2024 v2
	Gg of CO <sub>2</sub> eq.		%
2015	40 842.50	40 455.60	99.05%
2016	41 278.70	41 006.56	99.34%
2017	42 401.89	42 146.51	99.40%
2018	42 218.93	42 004.85	99.49%
2019	39 910.63	39 659.93	99.37%
2020	37 176.89	36 957.76	99.41%
2021	41 206.13	40 920.57	99.31%
2022	37 052.21	36 880.44	99.54%

Figure 10.1: Comparison of the recalculated GHG emissions trend without LULUCF and with indirect emissions in 2024 and 2025 submissions for 1990 – 2022 in Gg of CO<sub>2</sub> eq.





Table 10.2: Comparison of the recalculated GHG emissions trend with LULUCF and with indirect emissions in 2024 and 2025 submissions for 1990 – 2022 in Gg of CO<sub>2</sub> eq.

NATIONAL GHG INVENTORY WITH LULUCF AND WITH INDIRECT EMISSIONS			
YEAR	Submission 2024 v2	Submission 2025 v0.3	2025 v0.3/2024 v2
	Gg of CO <sub>2</sub> eq.		%
1990	64 562.80	64 620.81	100.09%
1991	54 480.23	54 463.38	99.97%
1992	48 419.10	47 854.33	98.83%
1993	45 044.28	44 509.63	98.81%
1994	43 087.65	42 522.16	98.69%
1995	44 149.65	43 590.22	98.73%
1996	44 094.12	43 529.37	98.72%
1997	44 099.60	43 519.59	98.68%
1998	42 299.24	41 713.36	98.61%
1999	41 885.22	41 330.77	98.68%
2000	39 981.55	39 439.59	98.64%
2001	42 890.50	42 389.16	98.83%
2002	41 144.36	40 602.60	98.68%
2003	41 950.09	41 502.90	98.93%
2004	42 569.21	42 165.17	99.05%

NATIONAL GHG INVENTORY WITH LULUCF AND WITH INDIRECT EMISSIONS			
YEAR	Submission 2024 v2	Submission 2025 v0.3	2025 v0.3/2024 v2
	Gg of CO <sub>2</sub> eq.		%
2005	46 417.70	46 072.14	99.26%
2006	43 348.38	42 908.64	98.99%
2007	41 976.54	41 487.01	98.83%
2008	43 551.22	42 979.04	98.69%
2009	39 556.75	38 978.91	98.54%
2010	41 184.39	40 772.59	99.00%
2011	39 746.09	39 307.55	98.90%
2012	36 275.51	35 847.59	98.82%
2013	35 182.73	34 754.13	98.78%
2014	35 348.12	34 933.33	98.83%
2015	35 608.23	35 220.94	98.91%
2016	35 966.00	35 693.32	99.24%
2017	37 197.65	36 941.68	99.31%
2018	37 987.62	37 772.80	99.43%
2019	34 916.16	34 664.75	99.28%
2020	29 997.91	29 777.95	99.27%
2021	33 994.72	33 707.75	99.16%
2022	29 826.47	29 653.26	99.42%

Figure 10.2: Comparison of the recalculated GHG emissions trend with LULUCF and with indirect emissions in 2024 and 2025 submissions for 1990 – 2022 in Gg of CO<sub>2</sub> eq.



### 10.3. Recalculations, Including in Response to the Review Process, and Planned Improvements to the Inventory

**UNFCCC review:** No UNFCCC review had been taking place in 2023. The status of implementation for the 2022 recommendations is described in [Table A4.3](#).

**EU ESR review:** The requirements for the Union review of the national inventory data submitted by Member States are set out in the regulation (EU) 2018/842 (ESR) and the regulation 2018/1999 (Governance). The initial, annual review 2024 concerning Member States' inventories for the year 2022 was carried out as planned during the spring 2024. Second step of the review of Slovakia was not necessary in the review cycle 2024 due to no issue was found. The reviewers raised several issues

---

during the first step of the 2024 ESR review which leads to no recommendation and were resolved during the first step of the review.

**Recalculations:** In term to further improvements of the GHG emissions inventory, the NS SR made recalculations for the 2025 submission. These recalculations are listed in [Table 10.3](#) below. Focus is on the main issues planned by the sectoral experts in the short and long term perspective, especially in the categories prioritised with the key impact on GHG emissions (for example national parameters applied in the agriculture). Major recalculations in sectors are in the Agricultural sector, mostly in rabbits and LULUCF – HWP, other recalculations are connected with the Copert software update, new input data and updating methodologies. Changes in methodological approach connecting to implementation of the IPCC 2019 Refinement to the 2006 IPCC Guidelines were developed and prepared under the project funded from EU grand successfully implemented in the years 2022 – 2023. More information can be found on the [website](#): PROJECT EMISSIONS – Preparation of methodology and improvement of emission inventories and emission projections.

In line with the project EMISSIONS, new archiving system was introduced into NS SR. Calculation IS MESAP serves as database of data, store system and enables data validation, uncertainty calculations and preparation of visualisation and graphs for different presentation purposes and deliverables.

The status of recommendations including planned improvements can be found in [Annex 4, Table A4.3](#) of this document, but also directly in the sectoral chapters.

**Table 4:** List of recalculations in March 15, 2025 submission (version 0.3) against September 15, 2024 submission (version 2) with short explanation

RECALCULATED CATEGORY (SUBMISSION 2024 v2 VERSUS SUBMISSION 2025 v0.3)		YEARS	GHG AFFECTED	EXPLANATION
<b>1. ENERGY SECTOR</b>				
1.A.3.b	Road transport	2013-2022	CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O	Recalculation is based on an update of the model to a newer version. The update involves correction of several emission factors, parameters, calculations, and adding new vehicle categories.
1.A.4.b	Households - Biomass	2021-2022	CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O	Biomass consumption was recalculated due to new statistical inputs on households
1.B.1.b	Charcoal production	2022	CH <sub>4</sub>	The recalculation is based on correction of calculation.
1.B.1.a	Coal mining and handling	1990-2022	CO <sub>2</sub> , CH <sub>4</sub>	The recalculations is a result of change of emission factors from underground mining. Slovakia switched from CIAB to IPCC emission factors. Also the calculation of emissions from abandoned mines was corrected.
1.B.2.c	Venting and flaring	1990-2022	CH <sub>4</sub>	Correction of calculation resulted in recalculation of emissions.
1.B.2.b	Gas post-meter	1990-2022	CO <sub>2</sub> , CH <sub>4</sub>	Reconstruction of time-series for appliances based on gas distribution resulted in recalculation of emissions.
<b>2. INDUSTRIAL PROCESSES AND PRODUCT USE SECTOR</b>				
2.D.3.d	Urea Catalytic Converters	2013-2022	CO <sub>2</sub>	Software update of the COPERT model resulted in the corrections to several emission factors, and the addition of new vehicle categories.
<b>3. AGRICULTURE</b>				
3.B.1.b	Non-dairy cattle	1990-2022	N <sub>2</sub> O	Fixed inconsistency for the activity data in the share of pasture on total AWMS for the animal subcategory (heifers) of Non-dairy cattle. No impact on trend and marginal impact on level of emissions.
3.B.3.a	Breeding swine Market swine	1990-2022	N <sub>2</sub> O	Update of the method for the calculation of nitrogen excretion rate for the swine categories.
3.B.4.h.i	Rabbits	1990-2022	CH <sub>4</sub> , N <sub>2</sub> O	Addition of new emissions source from rabbits.
3.D.1.b	Organic N fertilizers	1990-2022	N <sub>2</sub> O	Recalculation based on the recommendation during the review on identified irregularities between nitrogen volatilized as NH <sub>3</sub> and NO <sub>x</sub> . It will have impact in the 3.D.1.b Organic N fertilizers and 3.D.2 Indirect N <sub>2</sub> O emissions from managed soils categories.
3.D.1.b.iii	Other organic fertilizers	1990-2022	N <sub>2</sub> O	Recalculation based on the implementation of updated data on the N content in different types of fertilizers included in the category 3.D.1.b.iii Other organic fertilizers. The issue has been identified during QA process in cooperation with Central Controlling and Testing Institute in Agriculture.
3.D.2	Indirect N <sub>2</sub> O emissions from managed soils	1990-2022	N <sub>2</sub> O	Recalculation based on the recommendation during the review on identified irregularities between nitrogen volatilized as NH <sub>3</sub> and NO <sub>x</sub> . It



RECALCULATED CATEGORY (SUBMISSION 2024 v2 VERSUS SUBMISSION 2025 v0.3)		YEARS	GHG AFFECTED	EXPLANATION
				will have impact in the 3.D.1.b Organic N fertilizers and 3.D.2 Indirect N <sub>2</sub> O emissions from managed soils categories
<b>4. LULUCF</b>				
4(III)	Direct & indirect N <sub>2</sub> O emissions from N mineralization/immobilization	1991-2022	N <sub>2</sub> O	Calculation error (incorrected formula used) of indirect N <sub>2</sub> O emissions from N mineralization/immobilization.
4.G	Harvested Wood Products	2021-2022	CO <sub>2</sub>	Correction of input data.
<b>5. WASTE</b>				
5.A	Solid Waste Disposal – 5.A.1.a Anaerobic	2011-2022	CH <sub>4</sub>	Recalculation based on revision MSW composition (% share of paper + garden + food) based on consideration of recycling share.
5.A	Solid Waste Disposal – 5.A.1.a Anaerobic	2022	CH <sub>4</sub>	Recovery disposal gas was wrongly reported by the external organisation for 2022 (ÚRSO), data was corrected and lowered.
5.A	Solid Waste Disposal – 5.A.1.a Anaerobic	2010-2019	CH <sub>4</sub>	Activity data for waste disposal was updated by the Statistical Office of the Slovak Republic, minor changes.
5.A	Solid Waste Disposal – 5.A.1.a Anaerobic	2000-2022	CH <sub>4</sub>	Activity data for waste disposal was updated according real data from the disposal companies, approved by MŽP SR, minor changes 5-10%.
5.B.2.b	Composting of the Municipal Waste - 5.B.2.b. Other waste	2001-2022	CH <sub>4</sub>	This recalculation is connected with the correction of activity data of digestion in 2001 – 2022. The revision of new data is connected with data refinement provided by the NEIS.
5.C	Waste Incineration without Energy Use: 5.C.1.1.b (biogenic) and 5.C.1.2.b (non-biogenic)	1990-2022	CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O	Emissions of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O were recalculated for all-time series 1990 – 2022 due inclusion of the waste incinerated in the clinical waste incinerators. These recalculations increased biogenic as well as non-biogenic GHG emissions in equivalents.
5.D.1	Domestic Wastewater	1990-2022	CH <sub>4</sub>	Recalculations of methane emissions based on the implementation of different MCFs used for methane emission in individual retention tanks - cesspools. Explained different methane production in septic tanks and cesspools.
5.F	5.F.1 –Long-term C Storage in WDS	1990-2022	CO <sub>2</sub>	Recalculations are connected with recalculations in 5.A category for SWDS and parameters.
5.F	5.F.2 – Annual Change in Total Long-term C Storage 5.F.3 - Annual Change in Total Long-term C Storage in HWP Waste	2000-2022	CO <sub>2</sub>	Recalculations are connected with recalculations in 5.A category for SWDS and parameters.

---

## CHAPTER 11. INFORMATION ON CHANGES IN NATIONAL SYSTEM

The regular update of the SVK NS with all qualitative and quantitative indicators is provided in the NIDs and was provided in the First Biennial Transparency Report published in December 2024, Eight National Communication of the SR on Climate Change, published in February 2023 and in the Fifth Biennial Report in 2023.

There were no significant changes in the arrangement of the National System of the Slovak Republic during inventory preparation year 2024. National System description is provided in [Chapter 1.2](#).

However, several changes occurred during the year 2024 in the expert team due to including trainees and newcomers into the internal team of SVK NS. However, the SVK NS is continuing in the process of strengthening capacity among the national system in line with the improvement and prioritization plans. The uncertainties calculations were previously based on external cooperation, now (since the year 2021), an internal expert is responsible for all sectors across inventory. In addition, a new expert was involved in the cropland category to strengthen new calculations on land-based matrix and new expert was involved into agricultural team.

In addition, voluntary review undergone in categories fugitive emissions, SWDS and wastewater. The aim of the voluntary review was to verify changes in methodology connected with the implementation of the IPCC 2019 Refinements. The recalculations are described in the appropriate chapters of this document.

During previous years, the several new institutions were involved in the inventory, among others in transport (Control and Testing Body for road vehicles), Ministry of Transport of the Slovak Republic – Section of Buildings (for buildings energy balance mostly focusing of residential heating and cooling), State Nature Protection Body (for wetlands identification), new internal (SHMÚ) expert on emission projections and emissions estimation in household sector.

Figure and Tables in [Chapter 1.2.5](#) provide more information on actual structure and functions of the SVK NS and changes.

## CHAPTER 12. INFORMATION ON MINIMIZATION OF ADVERSE IMPACTS IN ACCORDANCE WITH ARTICLE 3, PARAGRAPH 14

Implementation of increasingly stringent environmental regulations and economic policies, which penalize further use of environmentally harmful substances, technologies and might be associated with a range of side effects. It is not excluded that some of possible adverse economic effects will affect some developing and least developed countries having less means for adequate remedial response measures. The magnitudes of these potential impacts are typically given by the stringency of adopted measures, selection of the particular policy instrument, size and strength of the implementing economy relative to the world markets and the actual macroeconomic set up of the affected developing countries.

---

In this chapter was updated based on the [5<sup>th</sup> Biennial Report of Slovakia to the UNFCCC](#) and identified potential channels of how domestically implemented environmental policies in the Slovak Republic might have exercised any impact on third countries. Furthermore, any existing evidence about the potential magnitudes of these effects is highlighted. Similarly, the activities in particular those related to the development aid of the Slovak Republic implemented in order to minimize the negative consequences caused by these policies are described in this chapter. The aim is to meet our commitments under the Kyoto Protocol in respect with transparent reporting on potential adverse social, environmental and economic impacts particularly on developing countries.

### Economic Impacts

Although the Slovak economy has decarbonized significantly in the last thirty years, further decarbonisation is needed. Slovakia went through a period of abrupt decarbonisation in the 1990s and 2000s that was caused by the changing structure of the economy, and technology improvements. Regardless of the improvements achieved so far, further decarbonisation is needed to contribute to the EU-wide decarbonisation goals in 2030 – decrease greenhouse gases by 55% compared to 1990 levels. This equals to abating an additional 6.3 million tonnes of CO<sub>2</sub> equivalent annually by 2030 (approximately 15% of current gross emissions). To model the most cost-effective path of decarbonisation, the first Slovak marginal abatement cost curve (MACC) was constructed. MACC compares various decarbonisation measures from all sectors of the economy by their price for a ton of CO<sub>2</sub> equivalent abated, and their abatement potential in 2030. Three emission-reduction goals were identified as follow: 55%, 67%, and 76% based on the MACC. These goals together with needed levers are discussed below in turn. Slovakia is close to achieving the EU-wide "Fit for 55" target to reduce emissions by 55% (6.3 Gg of CO<sub>2</sub> equivalents) in 2030 compared to the 1990 levels. While there is not yet an official target for Slovakia, a 55% reduction is achievable at a societal net cost (including public and private spending) of 2.7 billion EUR by 2030, via cost-effective levers below 30 EUR per ton of CO<sub>2</sub> equivalent (many of which have a negative price). Nevertheless, these levers are individually small and require complex implementation efforts across many stakeholders. Therefore, Slovakia should aim also beyond the 55% target and implement additional levers. Electrification of the steel sector is the key in the push for decarbonisation beyond the "Fit for 55" target. Currently the most polluting industry, it has many levers available that enable deep decarbonisation even without implementing carbon capture and storage (CCS). Electrification and efficiency improvements of the steel sector can abate in total of 6.2 Gg of CO<sub>2</sub> equivalents per year, additional levers across industries before the CCS could abate 1.7 Gg of CO<sub>2</sub> equivalents by 2030. The societal net cost would reach approximately 5 billion EUR by 2030. In total, this would lead to a 67% decrease compared to 1990. Reaching the full 2030 decarbonisation potential requires significant CCS investments. The key lever beyond 14.2 Gg of CO<sub>2</sub> equivalents abatement is the carbon capture and storage technology implemented across key point emitters to capture their remaining emissions. However, investing in CCS is CAPEX-intensive and would require significant political and societal efforts, including implementing supporting regulations. Total abatement compared to 1990 after implementing all the available levers would be 76% at a societal cost of over 13.5 billion EUR. Slovakia has a low-carbon electricity mix and expected electricity oversupply to support decarbonisation. Slovak low emissions intensity electricity creates suitable conditions for decarbonisation via electrification of the key sectors (e.g. transport and steel) as it will not result in significant secondary GHG emissions. With the decommissioning of Nováky and Vojany coal power plants, and the opening of nuclear power plants Mochovce 3 & 4, Slovakia will decarbonize its electricity generation even further (achieving ~90 ton of CO<sub>2</sub> equivalent/GWh) and will secure sufficient electricity supply to fulfil an increased demand from decarbonisation levers (e.g. electric arc furnaces). The MACC was constructed before the Russian invasion of Ukraine, but its conclusions remain relevant. The invasion motivated the EU to rapidly reduce dependence on Russian fossil fuels by increasing energy efficiency, which is fully in line with the measures suggested by this study. Importantly, as outlined in the REPowerEU plan, the EU climate targets are not jeopardized by the new geopolitical situation. The study was prepared in a joint collaboration of Value for Money Department, Ministry of Finance (ÚHP),

---

Institute for Environmental Policy (IEP), and Boston Consulting Group (BCG) during October and November 2021. The work was conducted via a joint project team composed of the authors of this study. During the MACC modelling, the authors used various internal and external benchmarks (including BCG proprietary databases and tools).

## REFERENCES

### GENERAL

2006 IPCC Guidelines for National Greenhouse Gas Inventories

2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories

2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands

2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol

Statistical Office of the Slovak Republic (updated every year).

Gera, M.; Bastak, I.; Gerova, E.; Kremler, M.; Szemesova, J. Climatologic Data Downscaling with Dynamical Adaptation for Purpose of Pollutant Spreading Analysis EMS Annual Meeting- ECAS 2008, 8<sup>th</sup> Annual Meetin of the European Meteorological Society, 7<sup>th</sup> European Conference on Applied Climatology" (ECAS), 29.9.-3.10.2008, Amsterdam, Abstracts, Vol. 5, 2008, ISSN 1812-7053.

Good Practices in the Slovak Republic with decoupling of the GHG Emission Growth Rate from Economic Development Slovak Hydrometeorological Institute, Ministry of the Environment, Bratislava, July 2008.

Report on the Estimation of Assigned Amount Units under the Kyoto Protocol revised version according FCCC/IRR/2007/SVK from July 2007, MZP SR and SHMU, Bratislava 2007.

The Fourth National Communication of the SR on Climate Change and Report on Demonstrable Progress of the SR to Achieve Commitments under the Kyoto Protocol the Ministry of Environment of the SR, Slovak Hydrometeorological Institute, Bratislava 2005.

The Fifth National Communication of the SR on Climate Change the Ministry of Environment of the SR, Slovak Hydrometeorological Institute, Bratislava 2009.

The Sixth National Communication of the SR on Climate Change the Ministry of Environment of the SR, Slovak Hydrometeorological Institute, Bratislava 2013.

The Seventh National Communication of the SR on Climate Change the Ministry of Environment of the SR, Slovak Hydrometeorological Institute, Bratislava 2018.

The Eight National Communication of the SR on Climate Change and the Fifth Biennial Report of the SR, Bratislava 2023.

IPCC Fifth Assessment Report, Chapter 8, 2014.

### CHAPTER 3 – ENERGY

2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories.

Energy 2023, Statistical Office of Slovak Republic (2024) ISBN: 978-80-8121-918-4.

COPERT 5 Computer programme to calculate emissions from road transport - User manual.

Szemesová, J.; Zemko, M.; Petraš, M.; Frankovič, B.: Assessment of the statistical survey on heating systems and fuels consumption in households. Slovenská štatistika a demografia 3/2021.

---

Szemesová, J.; Jonáček, Z.; Horváth, J.; Zetochová, L.; Labovský, J.: Improving the allocation of road transport emissions in AEA module and coherence between AEA and PEFA modules. Conference on Air protection. Proceedings 2021.

Horváth, J, Szemesová J. and Zetochová, L.: Štatistické odchýlky v údajoch o motorových palivách. Slovenská štatistika a demografia 1/2021.

Horváth, J., Mačala, J. Szemesová, J., Zetochová, L. Scrapping or not scrapping: About whether grants for scrapping and electromobility are effective. Air protection proceedings 2018.

Eva Krtková, Vladimír Danielik, Janka Szemesová, Klára Tarczay, Gábor Kis-Kovács and Vladimír Neuzil, Non-Energy Use of Fuels in the Greenhouse Gas Emission Reporting, Atmosphere 2019, 10, 406; doi:10.3390/atmos10070406.

Ntziachristos, L., Samaras, Z. EMEP/EEA air pollutant inventory guidebook 2023 on Passenger cars, light commercial vehicles, heavy duty vehicles including buses and motor cycles. European Environmental Agency 2019.

Horváth, J.; Szemesová, J.; Zetochová, L.: Roadmap to zero carbon road transport, Proceedings of the 24<sup>th</sup> International Transport and Air Pollution (TAP) Conference. 2022.

Horváth, J.; et. al: Results of the Eurostat project: New methodology for the Allocation of Emissions from Transport. International Conference Air Protection, High Tatras, 2022.

Jonáček, Z.; et. al.: Integrated Project LIFE AQI - Statistical Survey on the Use of Combustion Devices and Fuels Consumption in Households for the year 2022. International Conference Air Protection, High Tatras, 2022.

Horváth, J.; Szemesová, J.; Jonáček, Z.; Transport emission footprint in the Slovak economy, Proceedings of the 25<sup>th</sup> International Transport and Air Pollution (TAP) and the 3<sup>rd</sup> Shipping & Environment (S&E) Conference, 2024.

Lintnerová, O.: Geológia kaustobilitov. Uhlie a uhľovodíky. Archivované 2012-01-19 na Wayback Machine Uni, 2009.

#### **CHAPTER 4 – INDUSTRIAL PROCESSES**

Tomlein, P.; Tomlein M.: Refrigerants Logging and Reporting - Abstract from the IIR Conference on Sustainable Refrigeration and Heat Pump Technology, Stockholm Sweden 2010.

Tomlein, P.: Data collection system in Slovakia and why we consider Articles 5 and 18 of the F-gas proposal to be relevant, Presentation 2013.

Tomlein P., Tomlein Mi., Tomlein Ma.: Reporting System Based on Company Certification in Refrigeration Industry, Abstract, 2012.

Tomlein, P.: Inspection System on cooling circuits, in: IIR Compressors, Papiernička, 2004.

Tomlein, P.: Refrigerants logging and reporting, in: IIR Conference on Sustainable Refrigeration and Heat Pump Technology, Stockholm, Sweden, 2010.

#### **CHAPTER 5 – AGRICULTURE**

The Intergovernmental Panel on Climate Change, Agriculture, 2006 IPCC Guidelines for National Greenhouse Gas Inventory, Vol.4 Agriculture, Forestry and Other Land Use Methodology, Chapter 11 N<sub>2</sub>O emissions from managed soils, and CO<sub>2</sub> emissions from lime and urea application, Institute for Global Environmental Strategies, Hayama, Japan ISBN4-87888-032-4, 2006.

Green Report 2023 (Zelená správa 2023). The Ministry of Agriculture and Rural Development of the Slovak Republic 2023.

---

List of Livestock to the 31.12.2023, (Súpis hospodárskych zvierat k 31.12.2023). Statistical Office of the Slovak Republic.

Crop yields data for crops and vegetables in 2021 (Definitívne údaje o úrode poľnohospodárskych plodín a zeleniny v SR za rok 2020). Statistical Office of the Slovak Republic.

Animal production, sales of agricultural products from the primary production and crop balance in 2020 (Živočíšna výroba, predaj výrobkov z prvovýroby a bilancia plodín za rok 2020). Statistical Office of the Slovak Republic.

Nitrates Directive (Dusičnanová smernica). Ministry of Environment of the Slovak Republic.

The animal production, sales of primary production and crop balance. Statistical Office of the Slovak Republic.

Bielek, P.: Nitrogen in Agricultural Soils of the SR. Research institute of soil fertility Bratislava, 1998, ISBN: 80-85361-44-2 (in Slovak).

EUROSTAT: European Statistics code of practice, 2011, ISBN 978-92-79-21679-4, doi: 10.2785/18474

Bezák, P.: Guidelines for the Sewage Sludge Application (Príručka pre aplikáciu kalu z čistiarní odpadových vôd do poľnohospodárskej pôdy) (in Slovak). Bratislava 2010.

Tonhauzer, K.; Zetochová, L.: Comparison of the FAO GHG database and the Slovak greenhouse gas inventory activity data and emissions in agriculture. Meteorologický časopis, 20, 1, s. 3-9. ISSN 1335-339X, 2017.

Tonhauzer, K.; Tonhauzer, P.; Szemesová, J.; Šiška, B.: Estimation of N<sub>2</sub>O emissions from the agricultural soils and determination of nitrogen leakages. Atmosphere. Land-Atmosphere Interactions: Biogeophysical and Biogeochemical Feedbacks, 2020, Zv. 11 (6).

Brouček, J.; Brestenský V.; Palkovičová Z.: Differences in amounts of methane emissions from enteric fermentation from Slovak ewe farming between 2015 and 2016. Ecology & Safety, ISSN 1314-7234, Volume 12, 2017.

Petrikovič, P.; Heger, J.; Sommer A.: Nutrition for Pigs. Research Institute of animal production, 2005, ISBN 80-88872-45-6 in Slovak.

Strauch, D.; Baader, W.; Tietjen, C.: Waste from agricultural production. Ulmer Eugen Verlag, ISBN-978-3800143283, 1995, in German.

Perikovič, P.; Sommer, A.: Nutrition for Cattle. Research Institute for animal production. ISBN: 80-88872-21-9, 2002, in Slovak.

Jančová, M.: Growing and Utilization of Grassland and Clover grassland on Arable Land of Foothill and Mountain Areas. National Agriculture and Food Centre ISBN 978-80-89800-09-4, 2016, in Slovak.

Torma, S.; Vilček, J.: Postharvest residues of sugar beet and their role in the nutrient cycle. Listy Cukrovarnické a Reparské; Prague Vol. 133, Is. 9-10, p. 285-287, 2017, in Slovak.

FAOSTAT: Crops and livestock products, Fertilizers by Nutrient, Milk Production. Food and Agriculture Organization of the United Nations.

IFASTAT: Fertilizer consumption - statistical information on fertilizer & raw materials. International Fertilizers Association.

PLIS: Body weight, Pregnancy of animals, Information System about Breeds.

Lošák, T.; Dostál, J.: Digestát vo výžive a hnojení repky, Mendel university in Brno.

Gáborík, Š.: Dusíkaté hnojenie po zbere obilnín, The Central Control and Testing Institute in Agriculture.

---

Brestenský, V.; a kol. 2015. Chov hospodárskych zvierat. [Online] Národné poľnohospodárske a potravinárske centrum - Výskumný ústav živočíšnej výroby Nitra, 2015. [Dátum: 13. Marec 2019.] [http://www.vuzv.sk/pdf/chov\\_hz.pdf](http://www.vuzv.sk/pdf/chov_hz.pdf). ISBN 978-80-89418-41-1.

Brestenský, V.; Botto, L.: 2015. Odstraňovanie a skladovanie hospodárskych. Nitra : Agroinštitút Nitra, 2015.

PPA, 2015. Guidelines for the support for selected non-project measures, Bratislava, The Agricultural Paying Agency

## CHAPTER 6 – LULUCF

Green Report 2023 ([Zelená správa 2023](#)). The Ministry of Agriculture and Rural Development of the Slovak Republic.

Barančíková, G.; Makovníková, J.; Skalský, R.; Tarasovičová, Z.; Nováková, M.; Halás, J.; Koco, Š.; Gutteková, M. 2013: Changes in organic carbon pool in agricultural soils and its different development in individual agro-climatic regions of Slovakia. *Agriculture (Poľnohospodárstvo)*, vol. 59, 2013, no. 1, pp. 9–20.

Barančíková, G.; Makovníková, J.; Halas, J. 2016: Effect of land use change on soil organic carbon. *Agriculture (Poľnohospodárstvo)*, vol. 62, 1, pp. 10-18.

Bielek, P.; Jurčová, O. 2010: Metodika bilancie pôdnej organickej hmoty a stanovenie potreby organického hnojenia poľnohospodárskych pôd. Bratislava: VÚPOP, 2010. 145 s. ISBN 978-80-89128-80-8.

Cienciala, E.; Černý, E.; Tatarinov, F.; Apltauer, J.; Exnerová, Z. 2006: Biomass functions applicable to Scots pine. *Trees* 20:483–495.

Cienciala, E.; Apltauer, J.; Exnerová, Z.; Tatarinov, F. 2008: Biomass functions applicable to oak trees grown in central-european forestry. *Journal of Forest Science* 54:109–120.

Cienciala, E.; Palán, Š. 2014: Methodological background for quantification of CO<sub>2</sub> originating from changes in the pool of harvested wood products. Report for the Czech Ministry of the Environment, 26 pp (In Czech).

Halaj, J.; Petráš, R. 1998: Rastové tabuľky hlavných drevín/Yield tables of main tree species. Bratislava, Slovak Academic Press, 325 p.

Kobza, J.; Barančíková, G.; Cepková, V.; Došeková, A.; Fulajtár, E.; Houšková, B.; Makovníková, J.; Matúšková, L.; Medveď, M.; Pavlenda, P.; Schlosserová, J.; Styk, J.; Vojtáš, J. 2002: Soil monitoring of the Slovak republic - present state and development of monitored soil properties 1997-2001. Bratislava, Soil Science and Conservation Research Institute, 178 pp. (in Slovak).

Kobza, J.; Barančíková, G.; Čumová, L.; Dodok, R.; Hrivňáková, K.; Makovníková, J.; Náčiniaková - Bezáková, Z.; Pálka, B.; Pavlenda, P.; Schlosserová, J.; Styk, J.; Širáň, M.; Tothová, G. 2009: Soil monitoring of the Slovak republic - present state and development of monitored soil properties 2002 – 2006, Bratislava, Soil Science and Conservation Research Institute, 200 s. (in Slovak).

Kobza, J.; Barančíková, G.; Dodok, R.; Hrivňáková, K.; Makovníková, J.; Pálka, B.; Pavlenda, P.; Schlosserová, J.; Styk, J.; Širáň, M. 2014: Soil monitoring of the Slovak republic - present state and development of monitored soil properties 2007 – 2012. Bratislava, Soil Science and Conservation Research Institute, 249 s. (in Slovak).

Lehtonen, A.; Cienciala, E.; Tatarinov, F.; Mäkipää, R. 2007: Uncertainty estimation of biomass expansion factors for Norway spruce in the Czech Republic. *Annals of Forest Science* 64(2): p. 133-140.

- 
- Pajtík, J.; Konôpka, B.; Lukáč, M. 2011: Individual biomass factors for beech, oak and pine in Slovakia: a comparative study in young naturally regenerated stands. *Trees* 25, p. 277-288.
- Pavlenda, P. 2008: Quantification of carbon stocks in forest soils. In: Kobza, J. (Ed.): Piate pôdoznalecké dni. Pôda - národné bohatstvo. Zborník z medzinárodnej konferencie, Bratislava: VÚPOP, 2013, s. 243-250 (in Slovak).
- Pavlenda, P.; Capuliak, J.; Pavlendová, H. 2015: Soil Organic Carbon and Land Use Change: Can We Detect Soil Density Increase? *Pedometrics* 2015. 14-18 September, IUSS, University of Cordoba, Spain, p. 184.
- Pavlenda, P.; Priwitzer, T.; Capuliak, J.; Šebeň, V.; Bošela, M.; Pajtík, J.; Raši, R.; Pavlendová, H. 2016: Proposal for improvement of input data and procedures in the national methodology of greenhouse gases inventory in the LULUCF sector. Applicable deliverable of the project APVV-0243-11 Evaluation and modelling of carbon stocks in forest ecosystems for GHG inventory in the country (C-FORLAND), NFC, 20 pp. (in Slovak).
- Petráš, R.; Pajtík, J. 1991: Sústava česko-slovenských objemových tabuliek drevín./The system of Czecho-Slovak tree species volume tables. *Lesnícky časopis*, 37, (1), p. 49-56.
- Požgaj, A.; Chovanec, D.; Kurjatko, S.; Babiak, M., 1993: The Structure and properties of wood. *Príroda*, Bratislava, 468 pp. 1993, (in Slovak).
- Priwitzer, T.; Pajtík, J.; Konôpka, B.; Ištoňa, J.; Pavlenda, P. 2008: Quantification of carbon stocks in the model ecosystems of first age class. *Lesn. Čas. Forestry Journal*, 54, Supplement, 1, 2008, p. 69 – 79. (in Slovak with English summary).
- Priwitzer, T.; Pajtík, J.; Konôpka, B.; Ištoňa, J.; Pavlenda, P. 2009: Carbon allocation in forest ecosystem – case studies from young pine stands. *Lesn. Čas. – Forestry Journal*, 55(3): p. 239 – 250, (in Slovak with English summary).
- Raši, R.; Cienciala, E.; Priwitzer, T.; Palán, Š.; Pavlenda, P. 2015: [Bilancia uhlíka v drevných produktoch na Slovensku](#) (Carbon balance in harvested wood products in Slovakia), *Lesn. Cas. For. J.*, 61(2015) 101–106.
- Stanová, V., (ed) 2000: Rašeliniská Slovenska. DAPHNE – Inštitút aplikovanej ekológie, Bratislava, 194.
- Šály, R. 1998: Pedology and microbiology, 1998, TU Zvolen, Zvolen, in Slovak.
- Šebeň, V. 2017: Národná inventarizácia a monitoring lesov SR 2015-2016. Informácie, metódy, výsledky. (National forest inventory and monitoring SR 2015-2016. Information, methods, results) *Lesnícke štúdie* 65/2017. Národné lesnícke centrum – Lesnícky výskumný ústav Zvolen, 256 p. (in Slovak).
- Šebík, L.; Polák, L. 1990: The theory about wood production. Bratislava, *Príroda*: 322 pp., 1990, in Slovak.
- Šmelko, Š. 2000: Dendrometry. TU Zvolen, Zvolen, 399 p., 2000, in Slovak.
- Šmelko, Š.; Scheer, L.; Petráš, R.; Ďurský, J.; Fabrika, M. 2003: Measurement of forest and wood. Ústav pre výchovu a vzdelávanie pracovníkov LVH SR, Zvolen, 239 p. (in Slovak).
- Šmelko, Š.; Šebeň, V.; Bošela, M.; Merganič, J.; Jankovič, J. 2008: National forest inventory and monitoring of the Slovak republic 2005-2006. Basic concept and selected summary information, NFC-FRI Zvolen, Zvolen, 15 p.
- Šmelko, Š.; Šebeň, V.; Priwitzer, T. 2011: Determination of dry wood density and carbon stock in aboveground dendromass from sampling forest inventory in mountain forest ecosystem of the Biosphere Reserve Poľana. *Lesn. Čas. – Forestry Journal*, 57(2): p. 87–95.



---

Tóth, G.; Jones, A.; Montanarella, L. (Eds.) 2013: LUCAS TOPSOIL SURVEY - methodology, data and results. Luxembourg: Publications Office of the European Union, 2013. ISBN 978-92-79-32542-7.

Torma, S.; Vilček, J. 2017: Rastlinné zvyšky poľnohospodárskych plodín po zbere úrody – zdroj organickej hmoty. In: Moderná mechanizácia, roč. 20, č.3, s. 22-23. ISSN 1335-6178.

Wirth, C.; Schumacher, J.; Schulze, E. D. 2004: Generic biomass for Norway spruce in central Europe. A meta-analysis approach toward prediction and uncertainty estimation. *Tree Physiology* 24:121–139.

Wutzler, T.; Wirth, C.; Schumacher, J. 2008: Generic biomass functions for Common beech (*Fagus sylvatica* L.) in Central Europe - predictions and components of uncertainty. *Canadian Journal of Forest Research*, 38(6), 1661-1675.

## CHAPTER 7 – WASTE

MŽP SR Stratégia obmedzovania ukladania biologicky rozložiteľných odpadov na skládky odpadov Bratislava, XII. 2010.

Báreková, A.: Skúsenosti z realizácie projektu odpadového hospodárstva pre regionálne združenie obcí, TVIP 2014 Hustopoče IV.2014.

Priatel'ia Zeme SPZ Kompostovanie – Príručka o zbere a zhodnocovaní biologických odpadov, VII.2005.

Kaufman, P.: Metodika analýzy zloženia komunálneho odpadu, SPU Nitra 2009.

Veverka, M.; Lešinský, D.: Analýza stavu a vývoja nakladania s komunálnym odpadom na Slovensku. CEPTA Zvolen 05/2013.

Wagnerová, E.; Uriček, D.: Kogenerace - plyn z TKO ako zdroj energie. *Alternatívni energie* 5/2005.

MŽP SR Program odpadového hospodárstva SR na roky 2016 – 2020, Bratislava, X. 2015.

Gera, M.; Szemesova, J.: Uncertainty Analysis for Estimation of Landfill Emissions and Data Sensitivity for Input Variation, 2nd International Workshop on Uncertainty in Greenhouse Gas Inventories, International Institute for Applied Systems Analysis A-2361 Laxenburg, Austria, 27.-28.10. 2007.

Slovak Environmental Agency, Database of Waste Management Centre, MŽP Department of waste management, 2008.

Szemesová J.; Gera M.: Uncertainty Analysis for Estimation of Landfill Methane Emissions Contribution to Geophysics and Geodesy, SAS, Vol 37, No 3, 251 – 265, 2007.

Szemesová, J.; Gera, M.: Emission estimation of solid waste disposal sites according to the uncertainty analysis methodology, *Bioclimatology and Natural Hazards*, ISBN 978-80-228-17-60.

Szemesová J.; Gera M. Uncertainty analysis for estimation of landfill emissions and data sensitivity for the input variation, *Climatic Change* DOI 10.1007/s10584-010-9919-1, 2010.

Bodík, I.; Hutňan, M.; Sedláček, S.; Lazor M.: Možnosti a perspektívy zvyšovania produkcie bioplynu na komunálnych ČOV s prídavkom externých substrátov, 2009. (Potential and perspectives for increase of biogas generation in municipal WWTP by addition external substrates), in Slovak language only.

Štátny Geologický ústav Dionýza Štúra: Register Skládok odpadov, State Geological Institute of Dionýz Štúr: Disposal sites Registry.

Štatistický Úrad SR: Štatistické ročenky SR, ČSSR, ČSFR za roky 1950 – 2015 (Statistical office of the Slovak Republic: Statistical Yearbooks of Slovakia, Czechoslovak Socialist Republic and Czech and Slovak Federative Republic Waste Yearbook 1992 -2015).

Bodík, I.; Kubaská, M.: Municipal Sewage Sludge Management in the Slovak Republic, Actual Status and Perspectives, *Journal of Residuals Science & Technology*, Vol. 10, No. 4 October 2013.

---

Drahovská, D.; Kozáková, K.; Miháliková, M.; Kohút, M.: Pokrok v zbere, odvádzaní a čistení komunálnych odpadových vôd. In: Zborník konferencie Odpadové vody 2020, AČE SR 2020.

Škôrňová, J.; Korpísová, A.: Hodnotenie vypúšťaných odpadových vôd na Slovensku za rok 2019. In: Zborník konferencie Odpadové vody 2020, AČE SR 2020.

Kozáková, K.; Sumegová, L.: Produkcia kalov z komunálnych ČOV a nakladanie s nimi. In: Zborník konferencie Kaly a odpady 2020, AČE SR 2020.

Bodík, I.; Mrafková, L.; Szemesová, J.: Odpadové vody ako zdroj emisií skleníkových plynov (Wastewater, a Source of GHG Emissions). 12<sup>th</sup> Biannual Conference Wastewater 2022.

Bartman et al.: Wastewater Treatment and Discharge. IPCC Guidelines for National Greenhouse Gas Inventories, IPCC Refinements 2019.

Law, Y., Jacobsen, G.E., Smith, A.M., Yuan, Z. & Lant, P. (2013) Fossil organic carbon in wastewater and its fate in treatment plants. *Water Research* 47(14): 5270-5281.

Griffith, D.R., Barnes, R.T. & Raymond, P.A. (2009) Inputs of Fossil Carbon from Wastewater Treatment Plants to U.S. Rivers and Oceans. *Environmental Science & Technology* 43(15): 5647-5651.

---

## ANNEX 1. KEY CATEGORIES

Description of methodology used for identifying key categories:

This Annex describes and completed the methodology used to identify key categories. The level of disaggregation is based on the recommendation in the IPCC 2006 GL.

Key categories analysis for the year 2023 according to Approach 1 and Approach 2 (including uncertainties) (IPCC 2006 GL and IPCC 2019 Refinement) was performed with and without LULUCF by level and trend assessments.

By level assessment Approach 1, 28 key categories with LULUCF and 24 without LULUCF were identified and by level assessment Approach 2, 16 key categories with LULUCF and 19 without LULUCF were identified in 2023.

By trend assessment Approach 1, 32 key categories with LULUCF and 27 without LULUCF were identified and by trend assessment Approach 2, 19 key categories with LULUCF and 25 without LULUCF were identified.

The final results are presented in [Tables A1.1 - A1.4](#) and according to the [ERT recommendation G.5](#) from the draft ARR 2022, the summaries are presented in [Tables A1.5 - A1.6](#).

Analysis for the base year 1990 was performed by level assessment and 30 key categories with LULUCF and 25 without LULUCF were identified by Approach 1 and 20 key categories with LULUCF and 20 without LULUCF were identified by Approach 2.

The results are presented in [Table A1.7](#) and [Table A1.8](#) and the summary is presented in [Table A1.9](#).

More information on key categories and uncertainty assessment can be found in [Chapters 1.2.12](#) and [1.2.13](#) of this Document.

**Table A1.1: Key categories identified using Approach 1 and Approach 2 by level assessment (L1 & L2) with LULUCF in 2023**

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS 2023	LEVEL ASSESSMENT L1	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
2.F.1	Refrigeration and Air conditioning	HFCs	398.92	0.90	0.90	YES
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	1 385.82	3.11	4.10	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	2 783.58	6.26	10.36	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 026.18	4.55	14.91	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	277.03	0.62	15.89	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	3 181.94	7.15	23.04	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	1 899.01	4.27	27.31	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	269.06	0.60	27.91	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	7 538.72	16.94	44.86	YES
1.A.4	Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	213.78	0.48	45.58	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	306.43	0.69	46.27	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 677.32	8.27	54.54	YES
2.A.1	Mineral Industry - Cement Production	CO <sub>2</sub>	1 294.23	2.91	57.77	YES
2.A.2	Mineral Industry - Lime Production	CO <sub>2</sub>	475.06	1.07	58.84	YES
2.A.4	Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	210.09	0.47	59.34	YES
2.B.1	Chemical Industry - Ammonia Production	CO <sub>2</sub>	509.92	1.15	60.49	YES
2.B.8	Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	376.47	0.85	61.41	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	3733.94	8.39	69.80	YES
4.A.1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	-6 671.90	15.00	85.13	YES
4.A.2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	-337.60	0.76	85.88	YES
4.B.1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	-689.34	1.55	87.43	YES
4.G	Harvested Wood Products	CO <sub>2</sub>	-291.90	0.66	88.60	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	12.47	0.78	90.20	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	37.04	2.33	92.74	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	41.52	2.61	95.61	YES
3.B	Manure Management	N <sub>2</sub> O	0.78	0.52	97.47	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	1.74	1.16	98.63	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	1.14	0.77	99.40	YES

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS 2023	LEVEL ASSESSMENT L2	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	2 783.58	0.01	0.02	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	3 181.94	0.02	0.05	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	7 538.72	0.04	0.09	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 677.32	0.01	0.11	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	3 733.94	0.02	0.14	YES
4.A.1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	-6 671.90	0.53	0.68	YES
4.A.2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	-337.60	0.02	0.70	YES
4.B.1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	-689.34	0.05	0.75	YES
4.G	Harvested Wood Products	CO <sub>2</sub>	-291.90	0.01	0.78	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	37.04	0.02	0.82	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	41.52	0.03	0.85	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	7.46	0.01	0.86	YES
3.B	Manure Management	N <sub>2</sub> O	0.78	0.05	0.92	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	1.74	0.03	0.95	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	1.14	0.04	0.99	YES
5.B	Biological Treatment of Solid Waste	N <sub>2</sub> O	0.39	0.01	1.00	YES

**Table A1.2: Key categories identified using Approach 1 and Approach 2 by level assessment (L1 & L2) without LULUCF in 2023**

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS 2023	LEVEL ASSESSMENT L1	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
2.F.1	Refrigeration and Air conditioning	HFCs	398.92	1.11	1.11	YES
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	1 385.82	3.84	5.06	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	2 783.58	7.72	12.77	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 026.18	5.62	18.39	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	277.03	0.77	19.59	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	3 181.94	8.82	28.41	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	1 899.01	5.26	33.68	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	269.06	0.75	34.42	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	7 538.72	20.90	55.33	YES
1.A.4	Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	213.78	0.59	56.22	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	306.43	0.85	57.07	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 677.32	10.19	67.26	YES
2.A.1	Mineral Industry - Cement Production	CO <sub>2</sub>	1 294.23	3.59	71.26	YES
2.A.2	Mineral Industry - Lime Production	CO <sub>2</sub>	475.06	1.32	72.57	YES
2.A.4	Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	210.09	0.58	73.19	YES
2.B.1	Chemical Industry - Ammonia Production	CO <sub>2</sub>	509.92	1.41	74.60	YES
2.B.8	Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	376.47	1.04	75.74	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	3 733.94	10.35	86.09	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	12.47	0.97	88.46	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	37.04	2.88	91.59	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	41.52	3.22	95.09	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	7.46	0.58	95.67	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	0.27	0.02	95.69	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	1.14	0.84	99.41	YES

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS 2023	LEVEL ASSESSMENT L2	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	0.03	0.03	YES
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	0.03	0.03	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	12 861.05	0.08	0.11	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 293.69	0.01	0.12	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	9 028.53	0.07	0.21	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	3 930.58	0.02	0.23	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	4 503.02	0.03	0.26	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 634.43	0.02	0.36	YES
2.B.8	Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	428.80	0.01	0.38	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	4 167.97	0.03	0.42	YES
1.A.4	Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	1.45	0.00	0.46	YES
2.C.1	Metal Industry - Iron and Steel Production	CH <sub>4</sub>	0.49	0.01	0.48	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	111.14	0.08	0.56	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	27.92	0.03	0.60	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	2.60	0.01	0.61	YES
1.A.3.b	Fuel combustion - Road Transportation	N <sub>2</sub> O	0.19	0.00	0.63	YES
3.B	Manure Management	N <sub>2</sub> O	1.96	0.17	0.82	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	2.93	0.07	0.89	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	2.20	0.10	0.99	YES
5.B	Biological Treatment of Solid Waste	N <sub>2</sub> O	0.16	0.01	0.99	YES

**Table A1.3: Key categories identified using Approach 1 and Approach 2 by trend assessment (T1 & T2) with LULUCF in 2023**

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	TREND ASSESSMENT T1	CONTRIB. TO TREND %	CUMULATIVE TOTAL	KEY IN TREND ANALYSIS
2.F.1	Refrigeration and Air conditioning	HFCs	0.00	1.20	1.20	YES
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	0.00	0.87	2.19	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	0.03	8.59	10.78	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	0.01	3.07	13.85	YES
1.A.1	Fuel combustion - Energy Industries - Other Fossil Fuels	CO <sub>2</sub>	0.00	0.43	14.28	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	0.01	2.95	17.23	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	0.01	2.34	19.56	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	0.00	0.53	20.09	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	0.00	0.55	20.64	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	0.07	16.74	37.38	YES
1.A.3.e	Fuel combustion - Other Transportation	CO <sub>2</sub>	0.01	2.33	39.97	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	0.03	8.12	48.21	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	0.02	6.27	54.48	YES
2.A.1	Mineral Industry - Cement Production	CO <sub>2</sub>	0.01	1.96	57.05	YES
2.B.1	Chemical Industry - Ammonia Production	CO <sub>2</sub>	0.00	1.10	58.59	YES
2.B.8	Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	0.00	0.57	59.26	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	0.02	5.74	64.99	YES
4.A.1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	0.03	8.11	73.89	YES
4.A.2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	0.04	9.62	83.50	YES
4.B.1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	0.01	2.39	85.90	YES
4.B.2	Cropland - Land Converted to Cropland	CO <sub>2</sub>	0.00	0.51	86.41	YES
4.C.2	Grassland - Land Converted to Grassland	CO <sub>2</sub>	0.00	0.84	87.24	YES
4.G	Harvested Wood Products	CO <sub>2</sub>	0.01	1.33	88.80	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	0.00	0.47	89.39	YES
1.B.1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	0.00	0.65	90.47	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	0.00	0.78	91.25	YES
1.B.2.c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	0.00	0.65	91.90	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	0.00	1.00	92.92	YES
3.B	Manure Management	CH <sub>4</sub>	0.00	0.64	93.56	YES



IPCC CATEGORY CODE	IPCC CATEGORY	GAS	TREND ASSESSMENT T1	CONTRIB. TO TREND %	CUMULATIVE TOTAL	KEY IN TREND ANALYSIS
5.A	Solid Waste Disposal	CH <sub>4</sub>	0.01	2.47	96.06	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	0.00	0.53	96.59	YES
2.B.2	Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	0.01	1.27	98.40	YES

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	TREND ASSESSMENT T2	CONTRIB. TO TREND %	CUMULATIVE TOTAL	KEY IN TREND ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	0.15	1.63	2.03	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	0.34	3.69	7.96	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	0.20	2.25	10.81	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	0.09	1.01	11.82	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	0.14	1.53	14.26	YES
4.A.1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	2.73	29.88	44.65	YES
4.A.2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	2.19	24.06	68.71	YES
4.B.1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	0.71	7.77	76.47	YES
4.B.2	Cropland - Land Converted to Cropland	CO <sub>2</sub>	0.17	1.84	78.32	YES
4.C.2	Grassland - Land Converted to Grassland	CO <sub>2</sub>	0.28	3.02	81.34	YES
4.G	Harvested Wood Products	CO <sub>2</sub>	0.27	2.94	85.11	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	0.09	1.01	86.33	YES
1.A.4	Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	0.08	0.92	87.25	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	0.08	0.85	89.18	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	0.26	2.85	92.35	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	0.13	1.45	93.80	YES
3.B	Manure Management	N <sub>2</sub> O	0.11	1.24	96.39	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	0.09	0.98	97.37	YES
5.B	Biological Treatment of Solid Waste	N <sub>2</sub> O	0.09	1.04	99.62	YES

**Table A1.4: Key categories identified using Approach 1 and Approach 2 by trend assessment (T1 & T2) without LULUCF in 2023**

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	TREND ASSESSMENT T1	CONTRIB. TO TREND %	CUMULATIVE TOTAL	KEY IN TREND ANALYSIS
2.F.1	Refrigeration and Air conditioning	HFCs	0.01	1.49	1.49	YES
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	0.01	1.83	3.46	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	0.05	13.17	16.63	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	0.01	3.35	19.98	YES
1.A.1	Fuel combustion - Energy Industries - Other Fossil Fuels	CO <sub>2</sub>	0.00	0.52	20.50	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	0.02	4.22	24.72	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	0.02	4.67	29.39	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	0.00	0.64	30.14	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	0.07	19.84	49.99	YES
1.A.3.e	Fuel combustion - Other Transportation	CO <sub>2</sub>	0.01	3.24	53.63	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	0.04	11.40	65.30	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	0.03	7.05	72.34	YES
2.A.1	Mineral Industry - Cement Production	CO <sub>2</sub>	0.01	2.14	75.32	YES
2.B.1	Chemical Industry - Ammonia Production	CO <sub>2</sub>	0.00	1.29	76.99	YES
2.B.8	Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	0.00	0.62	77.73	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	0.02	6.28	84.01	YES
2.C.2	Metal Industry - Ferroalloys Production	CO <sub>2</sub>	0.00	0.51	84.53	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	0.00	0.66	85.88	YES
1.A.4	Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	0.00	0.52	86.41	YES
1.B.1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	0.00	0.96	87.37	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	0.00	1.24	88.61	YES
1.B.2.c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	0.00	0.93	89.54	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	0.01	1.85	91.42	YES
3.B	Manure Management	CH <sub>4</sub>	0.00	0.93	92.35	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	0.01	2.90	95.25	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	0.00	0.65	95.90	YES
2.B.2	Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	0.01	1.79	98.33	YES

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	TREND ASSESSMENT T2	CONTRIB. TO TREND %	CUMULATIVE TOTAL	KEY IN TREND ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	0.04	1.29	1.86	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	0.21	6.64	8.50	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	0.05	1.43	9.93	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	0.09	2.99	13.34	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	0.10	3.08	16.41	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	0.37	11.63	28.61	YES
1.A.3.e	Fuel combustion - Other Transportation	CO <sub>2</sub>	0.06	1.90	30.75	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	0.27	8.39	39.33	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	0.10	3.01	42.34	YES
2.B.8	Chemical Industry - Petrochemical and Carbon Black Production		0.03	0.94	45.10	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	0.14	4.47	49.57	YES
3.H	Urea Application	CO <sub>2</sub>	0.03	1.04	51.28	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	0.12	3.77	55.76	YES
1.A.4	Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	0.10	2.99	58.76	YES
1.B.1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	0.03	0.95	59.71	YES
2.C.1	Metal Industry - Iron and Steel Production	CH <sub>4</sub>	0.05	1.51	62.56	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	0.13	4.21	66.77	YES
3.B	Manure Management	CH <sub>4</sub>	0.04	1.23	68.00	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	0.28	8.91	76.91	YES
5.B	Biological Treatment of Solid Waste	CH <sub>4</sub>	0.15	4.66	81.57	YES
1.A.3b	Fuel combustion - Road Transportation	N <sub>2</sub> O	0.03	1.06	84.10	YES
3.B	Manure Management	N <sub>2</sub> O	0.23	7.22	93.29	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	0.06	1.98	95.27	YES
5.B	Biological Treatment of Solid Waste	N <sub>2</sub> O	0.11	3.34	98.75	YES
5.D	Wastewater Treatment and Discharge	N <sub>2</sub> O	0.02	0.78	99.53	YES

**Table A1.5: Key categories identified using Approach 1 and Approach 2 by level assessment (L1 & L2) with LULUCF in 1990**

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS BASE YEAR	LEVEL ASSESSMENT L1	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	4.53	4.53	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	12 861.05	15.26	19.79	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 293.69	2.72	22.51	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	2 867.64	3.40	25.95	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	9 028.53	10.71	36.67	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	3 930.58	4.66	41.33	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	4 503.02	5.34	46.91	YES
1.A.3.e	Fuel combustion - Other Transportation	CO <sub>2</sub>	1 813.95	2.15	49.51	YES
1.A.4	Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	580.74	0.69	50.20	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	6 852.15	8.13	58.32	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 634.43	4.31	62.64	YES
2.A.1	Mineral Industry - Cement Production	CO <sub>2</sub>	1 464.50	1.74	65.03	YES
2.A.2	Mineral Industry - Lime Production	CO <sub>2</sub>	794.92	0.94	65.97	YES
2.A.4	Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	446.73	0.53	66.51	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	4 167.97	4.94	72.36	YES
4.A.1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	-5 999.27	7.12	80.10	YES
4.A.2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	-2 263.04	2.68	82.79	YES
4.B.1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	-950.94	1.13	83.92	YES
4.B.2	Cropland - Land Converted to Cropland	CO <sub>2</sub>	466.51	0.55	84.47	YES
4.G	Harvested Wood Products	CO <sub>2</sub>	-470.41	0.56	85.72	YES
1.B.1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	28.39	0.94	87.28	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	49.56	1.65	88.94	YES
1.B.2.c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	23.55	0.78	89.72	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	111.43	3.70	93.44	YES
3.B	Manure Management	CH <sub>4</sub>	25.58	0.85	94.29	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	27.92	0.93	95.23	YES
2.B.2	Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	4.05	1.27	97.11	YES
3.B	Manure Management	N <sub>2</sub> O	2.12	0.67	97.79	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	3.03	0.95	98.74	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	2.31	0.73	99.47	YES

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS BASE YEAR	LEVEL ASSESSMENT L2	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	0.01	0.01	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	12 861.05	0.04	0.05	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	2 867.64	0.01	0.07	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	9 028.53	0.03	0.10	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	4 503.02	0.01	0.12	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	6 852.15	0.03	0.16	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	4 167.97	0.02	0.19	YES
4.A.1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	-5 999.27	0.32	0.52	YES
4.A.2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	-2 263.04	0.08	0.60	YES
4.B.1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	-950.94	0.04	0.64	YES
4.B.2	Cropland - Land Converted to Cropland	CO <sub>2</sub>	466.51	0.02	0.67	YES
4.C.2	Grassland - Land Converted to Grassland	CO <sub>2</sub>	-195.77	0.01	0.68	YES
4.F.2	Other land - Land Converted to Other Land	CO <sub>2</sub>	293.10	0.02	0.70	YES
4.G	Harvested Wood Products	CO <sub>2</sub>	-470.41	0.01	0.71	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	13.72	0.01	0.73	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	111.43	0.04	0.78	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	27.92	0.01	0.80	YES
3.B	Manure Management	N <sub>2</sub> O	2.12	0.09	0.90	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	3.03	0.03	0.94	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	2.31	0.05	0.99	YES

**Table A1.6: Key categories identified using Approach 1 and Approach 2 by level assessment (L1 & L2) without LULUCF in 1990**

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS BASE YEAR	LEVEL ASSESSMENT L1	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	5.20	5.20	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	12 861.05	17.52	22.72	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 293.69	3.12	25.84	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	2 867.64	3.91	29.80	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	9 028.53	12.30	42.09	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	3 930.58	5.35	47.44	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	4 503.02	6.13	53.86	YES
1.A.3.e	Fuel combustion - Other Transportation	CO <sub>2</sub>	1 813.95	2.47	56.83	YES
1.A.4	Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	580.74	0.79	57.62	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	6 852.15	9.33	66.96	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 634.43	4.95	71.91	YES
2.A.1	Mineral Industry - Cement Production	CO <sub>2</sub>	1 464.50	1.99	74.65	YES
2.A.2	Mineral Industry - Lime Production	CO <sub>2</sub>	794.92	1.08	75.74	YES
2.A.4	Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	446.73	0.61	76.35	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	4 167.97	5.68	83.07	YES
1.B.1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	28.39	1.08	85.57	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	49.56	1.89	87.48	YES
1.B.2.c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	23.55	0.90	88.38	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	111.43	4.25	92.65	YES
3.B	Manure Management	CH <sub>4</sub>	25.58	0.98	93.62	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	27.92	1.06	94.69	YES
2.B.2	Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	4.05	1.46	96.84	YES
3.B	Manure Management	N <sub>2</sub> O	2.12	0.76	97.62	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	3.03	1.09	98.72	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	2.31	0.83	99.55	YES

IPCC CATEGORY CODE	IPCC CATEGORY	GAS	EMISSIONS/ REMOVALS BASE YEAR	LEVEL ASSESSMENT L2	CUMULATIVE TOTAL	KEY IN LEVEL ANALYSIS
1.A.1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	0.03	0.03	YES
1.A.1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	12 861.05	0.07	0.10	YES
1.A.1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 293.69	0.01	0.12	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	2 867.64	0.02	0.14	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	9 028.53	0.07	0.21	YES
1.A.2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	3 930.58	0.02	0.23	YES
1.A.3.b	Fuel combustion - Road Transportation	CO <sub>2</sub>	4 503.02	0.03	0.26	YES
1.A.3.e	Fuel combustion - Other Transportation	CO <sub>2</sub>	1 813.95	0.01	0.27	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	6 852.15	0.06	0.34	YES
1.A.4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 634.43	0.02	0.35	YES
2.C.1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	4 167.97	0.03	0.41	YES
1.A.4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	13.72	0.03	0.44	YES
1.B.1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	28.39	0.01	0.45	YES
1.B.2.b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	49.56	0.01	0.46	YES
3.A	Enteric Fermentation	CH <sub>4</sub>	111.43	0.08	0.56	YES
3.B	Manure Management	CH <sub>4</sub>	25.58	0.01	0.57	YES
5.A	Solid Waste Disposal	CH <sub>4</sub>	27.92	0.03	0.60	YES
5.D	Wastewater Treatment and Discharge	CH <sub>4</sub>	2.12	0.18	0.81	YES
3.B	Manure Management	N <sub>2</sub> O	3.03	0.07	0.89	YES
3.D.1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	2.31	0.10	0.99	YES
3.D.2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	3 819.21	0.03	0.03	YES

**Table A1.7: Key categories identified using Approach 1 and Approach 2 by level assessment with and without LULUCF in 2023**

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 2023 WITH LULUCF	APPROACH 1 2023 WITHOUT LULUCF	APPROACH 2 2023 WITH LULUCF	APPROACH 2 2023 WITHOUT LULUCF
2.F.1 Refrigeration and Air conditioning	HFCs	YES	YES	NO	NO
1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	YES	YES	NO	NO

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 2023 WITH LULUCF	APPROACH 1 2023 WITHOUT LULUCF	APPROACH 2 2023 WITH LULUCF	APPROACH 2 2023 WITHOUT LULUCF
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	YES	YES	NO	NO
1.A.3.b Fuel combustion - Road Transportation	CO <sub>2</sub>	YES	YES	YES	YES
1.A.4 Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	YES	YES	NO	NO
1.A.4 Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	YES	YES	NO	NO
1.A.4 Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	YES	YES	YES	YES
2.A.1 Mineral Industry - Cement Production	CO <sub>2</sub>	YES	YES	NO	NO
2.A.2 Mineral Industry - Lime Production	CO <sub>2</sub>	YES	YES	NO	NO
2.A.4 Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	YES	YES	NO	NO
2.B.1 Chemical Industry - Ammonia Production	CO <sub>2</sub>	YES	YES	NO	NO
2.B.8 Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	YES	YES	NO	YES
2.C.1 Metal Industry - Iron and Steel Production	CO <sub>2</sub>	YES	YES	YES	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	YES	X	YES	X
4.A.2 Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	YES	X	YES	X
4.B.1 Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	YES	X	YES	X
4.G Harvested Wood Products	CO <sub>2</sub>	YES	X	YES	X
1.A.4 Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	NO	NO	NO	YES
1.B.2.b Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	YES	YES	NO	NO
2.C.1 Metal Industry - Iron and Steel Production	CH <sub>4</sub>	NO	NO	NO	YES
3.A Enteric Fermentation	CH <sub>4</sub>	YES	YES	YES	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	CH <sub>4</sub>	NO	X	NO	X
4.A.2 Forest Land - Land Converted to Forest Land	CH <sub>4</sub>	NO	X	NO	X
5.A Solid Waste Disposal	CH <sub>4</sub>	YES	YES	YES	YES
5.B Biological Treatment of Solid Waste	CH <sub>4</sub>	NO	YES	YES	YES
1.A.3.b Fuel combustion - Road Transportation	N <sub>2</sub> O	NO	NO	NO	YES
3.B Manure Management	N <sub>2</sub> O	YES	NO	YES	YES
3.D.1 Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	YES	YES	YES	YES
3.D.2 Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	YES	YES	YES	YES
5.B Biological Treatment of Solid Waste	N <sub>2</sub> O	NO	NO	YES	YES



**Table A1.8: Key categories identified using Approach 1 and Approach 2 by trend assessment with and without LULUCF in 2023**

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 2023 WITH LULUCF	APPROACH 1 2023 WITHOUT LULUCF	APPROACH 2 2023 WITH LULUCF	APPROACH 2 2023 WITHOUT LULUCF
2.F.1 Refrigeration and Air conditioning	HFCs	YES	YES	NO	NO
1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.1 Fuel combustion - Energy Industries - Other Fossil Fuels	CO <sub>2</sub>	YES	YES	NO	NO
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	YES	NO	NO	NO
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	YES	YES	NO	NO
1.A.3.b Fuel combustion - Road Transportation	CO <sub>2</sub>	YES	YES	YES	YES
1.A.3.e Fuel combustion - Other Transportation	CO <sub>2</sub>	YES	YES	NO	YES
1.A.4 Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.4 Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	YES	YES	YES	YES
2.A.1 Mineral Industry - Cement Production	CO <sub>2</sub>	YES	YES	NO	NO
2.B.1 Chemical Industry - Ammonia Production	CO <sub>2</sub>	YES	YES	NO	NO
2.B.8 Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	YES	YES	NO	YES
2.C.1 Metal Industry - Iron and Steel Production	CO <sub>2</sub>	YES	YES	YES	YES
2.C.2 Metal Industry - Ferroalloys Production	CO <sub>2</sub>	NO	YES	NO	NO
3.H Urea Application	CO <sub>2</sub>	NO	NO	NO	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	YES	X	YES	X
4.A.2 Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	YES	X	YES	X
4.B.1 Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	YES	X	YES	X
4.B.2 Cropland - Land Converted to Cropland	CO <sub>2</sub>	YES	X	YES	X
4.C.2 Grassland - Land Converted to Grassland	CO <sub>2</sub>	YES	X	YES	X
4.E.2 Settlements - Land Converted to Settlements	CO <sub>2</sub>	NO	X	NO	X
4.F.2 Other land - Land Converted to Other Land	CO <sub>2</sub>	NO	X	NO	X
4.G Harvested Wood Products	CO <sub>2</sub>	YES	X	YES	X
1.A.4 Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	YES	YES	YES	YES
1.A.4 Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	NO	YES	YES	YES

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 2023 WITH LULUCF	APPROACH 1 2023 WITHOUT LULUCF	APPROACH 2 2023 WITH LULUCF	APPROACH 2 2023 WITHOUT LULUCF
1.B.1 Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	YES	YES	NO	YES
1.B.2.b Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	YES	YES	NO	NO
1.B.2.c Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	YES	YES	NO	NO
2.C.1 Metal Industry - Iron and Steel Production	CH <sub>4</sub>	NO	NO	NO	YES
3.A Enteric Fermentation	CH <sub>4</sub>	YES	YES	YES	YES
3.B Manure Management	CH <sub>4</sub>	YES	YES	NO	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	CH <sub>4</sub>	NO	X	NO	X
4.A.2 Forest Land - Land Converted to Forest Land	CH <sub>4</sub>	NO	X	NO	X
5.A Solid Waste Disposal	CH <sub>4</sub>	YES	YES	YES	YES
5.B Biological Treatment of Solid Waste	CH <sub>4</sub>	YES	YES	YES	YES
1.A.3.b Fuel combustion - Road Transportation	N <sub>2</sub> O	NO	NO	NO	YES
2.B.2 Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	YES	YES	NO	NO
3.B Manure Management	N <sub>2</sub> O	NO	NO	YES	YES
3.D.1 Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	NO	NO	YES	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	N <sub>2</sub> O	NO	X	NO	X
4.A.2 Forest Land - Land Converted to Forest Land	N <sub>2</sub> O	NO	X	NO	X
4.B.2 Cropland - Land Converted to Cropland	N <sub>2</sub> O	NO	X	NO	X
4.C.2 Grassland - Land Converted to Grassland	N <sub>2</sub> O	NO	X	NO	X
4.E.2 Settlements - Land Converted to Settlements	N <sub>2</sub> O	NO	X	NO	X
4.F.2 Other land - Land Converted to Other Land	N <sub>2</sub> O	NO	X	NO	X
5.B Biological Treatment of Solid Waste	N <sub>2</sub> O	NO	NO	YES	YES
5.D Wastewater Treatment and Discharge	N <sub>2</sub> O	NO	NO	NO	YES

**Table A1.9: Key categories identified using Approach 1 and Approach 2 by level assessment with and without LULUCF in 1990**

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 1990 WITH LULUCF	APPROACH 1 1990 WITHOUT LULUCF	APPROACH 2 1990 WITH LULUCF	APPROACH 2 1990 WITHOUT LULUCF
1.A.1 Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.1 Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.1 Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	YES	YES	YES	YES

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 1990 WITH LULUCF	APPROACH 1 1990 WITHOUT LULUCF	APPROACH 2 1990 WITH LULUCF	APPROACH 2 1990 WITHOUT LULUCF
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.2 Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	YES	YES	NO	YES
1.A.3.b Fuel combustion - Road Transportation	CO <sub>2</sub>	YES	YES	YES	YES
1.A.3.e Fuel combustion - Other Transportation	CO <sub>2</sub>	YES	YES	NO	YES
1.A.4 Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	YES	YES	NO	NO
1.A.4 Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	YES	YES	YES	YES
1.A.4 Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	YES	YES	NO	YES
2.A.1 Mineral Industry - Cement Production	CO <sub>2</sub>	YES	YES	NO	NO
2.A.2 Mineral Industry - Lime Production	CO <sub>2</sub>	YES	YES	NO	NO
2.A.4 Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	YES	YES	NO	NO
2.C.1 Metal Industry - Iron and Steel Production	CO <sub>2</sub>	YES	YES	YES	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	YES	X	YES	X
4.A.2 Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	YES	X	YES	X
4.B.1 Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	YES	X	YES	X
4.B.2 Cropland - Land Converted to Cropland	CO <sub>2</sub>	YES	X	YES	X
4.C.2 Grassland - Land Converted to Grassland	CO <sub>2</sub>	NO	X	YES	X
4.E.2 Settlements - Land Converted to Settlements	CO <sub>2</sub>	NO	X	NO	X
4.F.2 Other land - Land Converted to Other Land	CO <sub>2</sub>	NO	X	YES	X
4.G Harvested Wood Products	CO <sub>2</sub>	YES	X	YES	X
1.A.4 Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	NO	NO	YES	YES
1.B.1 Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	YES	YES	NO	YES
1.B.2.b Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	YES	YES	NO	YES
1.B.2.c Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	YES	YES	NO	NO
3.A Enteric Fermentation	CH <sub>4</sub>	YES	YES	YES	YES
3.B Manure Management	CH <sub>4</sub>	YES	YES	NO	YES
4.A.1 Forest Land - Forest Land Remaining Forest Land	CH <sub>4</sub>	NO	X	NO	X
4.A.2 Forest Land - Land Converted to Forest Land	CH <sub>4</sub>	NO	X	NO	X
5.A Solid Waste Disposal	CH <sub>4</sub>	YES	YES	YES	YES
2.B.2 Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	YES	YES	NO	NO
3.B Manure Management	N <sub>2</sub> O	YES	YES	YES	YES

IPCC SOURCE CATEGORIES	GAS	APPROACH 1 1990 WITH LULUCF	APPROACH 1 1990 WITHOUT LULUCF	APPROACH 2 1990 WITH LULUCF	APPROACH 2 1990 WITHOUT LULUCF
3.D.1 Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	YES	YES	YES	YES
3.D.2 Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	YES	YES	YES	YES

---

## ANNEX 2. ASSESSMENT OF COMPLETENESS

Assessment of completeness is one of the elements of quality control procedure in the inventory preparation on the general and sectoral level. The completeness of the emission inventory is improving from year to year and the updates are regularly reported in the national inventory documents. The completeness check for ensuring time series consistency is performed and the estimation is complete in recent inventory submission (2025).

Several categories are reported as not occurring (NO) due to the not existence of the emission source or activity is not occurring in Slovakia. If the methodology does not exist in the IPCC 2006 GL, the notation key not applicable (NA) was used. Several categories are not estimated (NE) because of emissions are under the threshold. The included elsewhere categories (IE) are listed in CRT table 9 with the explanations and also described in this document in the appropriate sectoral chapters. Lists of information on notation keys used for each sector was prepared, see [Tables A2.1-A2.7](#) below.

Both direct GHGs as well as precursor gases are covered by the inventory of the Slovak Republic. The geographic coverage is complete; the whole territory of the Slovak Republic is covered by the inventory.

**Table A2.1: Notation keys in the Energy sector – combustion of fuels which are not occurring in specific subcategory**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS		
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
1.A. Fuel combustion				1.A.2 Manufacturing industries and construction			
1. A.1. Energy industries				a. Iron and steel			
Peat	NO	NO	NO	Other fossil fuels	NO	NO	NO
a. Public electricity and heat production				Peat	NO	NO	NO
Peat	NO	NO	NO	b. Non-ferrous metals			
1.A.1.a.i Electricity Generation				Other fossil fuels	NO	NO	NO
Liquid Fuels	NO	NO	NO	Peat	NO	NO	NO
Solid Fuels	NO	NO	NO	c. Chemicals			
Other Fossil Fuels	NO	NO	NO	Peat	NO	NO	NO
Peat	NO	NO	NO	d. Pulp, paper and print			
1.A.1.a.ii Combined heat and power generation				Other fossil fuels	NO	NO	NO
Peat	NO	NO	NO	Peat	NO	NO	NO
1.A.1.a.iii Heat plants				e. Food processing, beverages and tobacco			
Liquid Fuels	NO	NO	NO	Other fossil fuels	NO	NO	NO
Peat	NO	NO	NO	Peat	NO	NO	NO
b. Petroleum refining				f. Non-metallic minerals			
Solid fuels	NO	NO	NO	Peat	NO	NO	NO
Other fossil fuels	NO	NO	NO	1.A.2.g.i Manufacturing of machinery			
Peat <sup>(5)</sup>	NO	NO	NO	Solid Fuels	NO	NO	NO
Biomass <sup>(6)</sup>	NO	NO	NO	Other Fossil Fuels	NO	NO	NO
c. Manufacture of solid fuels and other energy industries				Peat	NO	NO	NO
Other fossil fuels	NO	NO	NO	1.A.2.g.ii Manufacturing of transport equipment			
Peat	NO	NO	NO	Other Fossil Fuels	NO	NO	NO
Biomass	NO	NO	NO	Peat	NO	NO	NO
1.A.1.c.i Manufacture of solid fuels				1.A.2.g.iii Mining (excluding fuels) and quarrying			
Liquid Fuels	NO	NO	NO	Other Fossil Fuels	NO	NO	NO
Other Fossil Fuels	NO	NO	NO	Peat	NO	NO	NO
Peat	NO	NO	NO	1.A.2.g.iv Wood and wood products			
Biomass	NO	NO	NO	Solid Fuels	NO	NO	NO
1.A.1.c.ii Oil and gas extraction				Other Fossil Fuels	NO	NO	NO
Solid fuels				Peat	NO	NO	NO
Other Fossil Fuels	NO	NO	NO	1.A.2.g.v Construction			
Peat	NO	NO	NO	Other Fossil Fuels	NO	NO	NO
Biomass	NO	NO	NO	Peat	NO	NO	NO

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS		
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
1.A.2.g.vi Textile and leather				Solid fuels	NO	NO	NO
Other Fossil Fuels	NO	NO	NO	Other fossil fuels	NO	NO	NO
Peat	NO	NO	NO	Biomass	NO	NO	NO
1.A.2.g.viii Other (please specify)				i. Pipeline transport			
Other Fossil Fuels	NO	NO	NO	Liquid fuels	NO	NO	NO
Peat	NO	NO	NO	Solid fuels	NO	NO	NO
1.A.3 Transport				Other fossil fuels	NO	NO	NO
Solid fuels	NO	NO	NO	Biomass	NO	NO	NO
a. Domestic aviation				ii. Other	NO	NO	NO
Biomass	NO	NO	NO	1.A.4 Other sectors			
b. Road transportation				Other fossil fuels	NO	NO	NO
Other liquid fuels	NO	NO	NO	Peat	NO	NO	NO
ii. Light duty trucks				a. Commercial/institutional			
Liquefied petroleum gases (LPG)	NO	NO	NO	Other fossil fuels	NO	NO	NO
Other liquid fuels	NO	NO	NO	Peat	NO	NO	NO
Gaseous fuels	NO	NO	NO	1.A.4.a.i Stationary combustion			
iii. Heavy duty trucks and buses				Other Fossil Fuels	NO	NO	NO
Liquefied petroleum gases (LPG)	NO	NO	NO	Peat	NO	NO	NO
Other liquid fuels	NO	NO	NO	b. Residential			
iv. Motorcycles				Other fossil fuels	NO	NO	NO
Liquefied petroleum gases (LPG)	NO	NO	NO	Peat	NO	NO	NO
Other liquid fuels	NO	NO	NO	1.A.4.b.i Stationary combustion			
Gaseous fuels	NO	NO	NO	Other Fossil Fuels	NO	NO	NO
v. Other	IE, NO	NO	NO	Peat	NO	NO	NO
c. Railways				c. Agriculture/forestry/fishing			
Solid fuels	NO	NO	NO	Other fossil fuels	NO	NO	NO
Gaseous fuels	NO	NO	NO	Peat	NO	NO	NO
Other fossil fuels	NO	NO	NO	i. Stationary			
d. Domestic Navigation				Other fossil fuels	NO	NO	NO
Residual fuel oil	NO	NO	NO	Peat	NO	NO	NO
Gasoline	NO	NO	NO	ii. Off-road vehicles and other machinery			
Other liquid fuels	NO	NO	NO	Liquefied petroleum gases (LPG)	NO	NO	NO
Gaseous fuels	NO	NO	NO	Other liquid fuels	NO	NO	NO
Other fossil fuels	NO	NO	NO	Gaseous fuels	NO	NO	NO
e. Other transportation				Other fossil fuels	NO	NO	NO
Liquid fuels	NO	NO	NO	iii. Fishing	NO	NO	NO

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS		
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O		CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O
Residual fuel oil	NO	NO	NO	Gaseous fuels	NO	NO	NO
Gas/diesel oil	NO	NO	NO	Other fossil fuels	NO	NO	NO
Gasoline	NO	NO	NO	Biomass	NO	NO	NO
Other liquid fuels	NO	NO	NO	Military Diesel Oil			
Gaseous fuels	NO	NO	NO	Solid fuels	NO	NO	NO
Biomass	NO	NO	NO	Gaseous fuels	NO	NO	NO
Other fossil fuels	NO	NO	NO	Other fossil fuels	NO	NO	NO
1.A.5 Other (Not specified elsewhere)				Biomass	NO	NO	NO
a. Stationary (please specify)				1.D.1.a International aviation (aviation bunkers)			
Other				Biomass	NO	NO	NO
Other Fossil Fuels	NO	NO	NO	1.D.1.b International navigation (marine bunkers)			
Peat	NO	NO	NO	Residual fuel oil	NO	NO	NO
b. Mobile (please specify)				Gasoline	NO	NO	NO
Military use Jet Kerosene				Other liquid fuels	NO	NO	NO
Solid fuels	NO	NO	NO	Gaseous fuels	NO	NO	NO
Gaseous fuels	NO	NO	NO	Biomass	NO	NO	NO
Other fossil fuels	NO	NO	NO	Other fossil fuels	NO	NO	NO
Biomass	NO	NO	NO	1.D.2 Multilateral operations	NO	NO	NO
Military Gasoline							
Solid fuels	NO	NO	NO				

**Table A2.2: Notation keys in the Energy sector - categories 1.B.1 and 1.B.2**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CH <sub>4</sub>		CO <sub>2</sub>	
	Recovery/Flaring	Emissions	Emissions	
1.B.1.a Coal Mining and Handling	NO			CH <sub>4</sub> recovery is not occurring in Slovakia from this activity
i. Underground mines	NO			
Mining activities	NO			
Post-mining activities	NO		NO	Emissions not occurring in this subcategory
Abandoned underground mines	NO			Emissions not occurring in this subcategory
Flaring of drained methane or conversion of methane to CO <sub>2</sub>		NO	NO	
Other	NO	NO	NO	



GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CH <sub>4</sub>		CO <sub>2</sub>	
	Recovery/Flaring	Emissions	Emissions	
ii. Surface mines	NO	NO	NO	No surface mines are occurring in Slovakia
Mining activities	NO	NO	NO	
Post-mining activities	NO	NO	NO	
1.B.1.b Solid Fuel Transformation	NO		NO	
1.B.1.c Other	NA	NA	NA	

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>		CH <sub>4</sub>	N <sub>2</sub> O	COMMENT
	Emissions	Amount captured			
1.B.2.a Oil		NO			CO <sub>2</sub> is not captured in Slovakia from this activity
1. Exploration	NO	NO	NO	NO	This activity is not occurring in Slovakia
2. Production		NO			Emissions not occurring in this subcategory
3. Transport		NO			Emissions not occurring in this subcategory
4. Refining/storage		NE			Emissions are not estimated from CO <sub>2</sub> capture
6. Other	NO	NO	NO	NO	No other source exists
1.B.2.b Natural Gas		NO			
1. Exploration	NO	NO	NO	NO	This activity is not occurring in Slovakia
2. Production		NO			
3. Processing		NO			
4. Transmission and storage		NO			
5. Distribution		NO			
6. Other		NA, NO			
1 B.2.c Venting and Flaring		NO			
Venting		NO			
ii. Gas		NO			
iii. Combined	NO	NO	NO		This activity is not occurring in Slovakia

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>		CH <sub>4</sub>	N <sub>2</sub> O	COMMENT
	Emissions	Amount captured			
Flaring		NO			
iii. Combined	NO	NO	NO	NO	This activity is not occurring in Slovakia

**Table A2.3: Notation keys in the Energy sector – combustion of fuels which are IE and NE in specific subcategory**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	
1.A.3 Transport				
b. Road transportation				
Other liquid fuels (please specify)	IE	IE	IE	Emissions from combustion of lubricants in two-stroke engines are included in those of gasoline
i. Cars				
Other liquid fuels (please specify)	IE	IE	IE	The emissions from combustion of lubricants in two-stroke engines are included in those of gasoline.
v. Other (please specify)	IE			Emissions reported in category non-energy products from fuels and solvent use - other (2.D.3)
Urea-based catalysts	IE	IE	IE	Emissions reported in category non-energy products from fuels and solvent use - other (2.D.3)
Diesel Oil	IE			
1. B. 1. b. Fuel transformation				
2. Coke production	IE			
1. B. 2. a. Oil				
5. Distribution of oil products	NE		NE	This activity is occurring in Slovakia but no EFs are available in the IPCC 2006 GL (Not determined for EFs Table 4.2.4) and also not in the 2019 Refinements.
1. B. 2. c. Venting and flaring				
Venting				
i. Oil	IE	IE		Included in appropriate 1.B.2.a categories based on 2019 IPCC Refinement
Flaring				
i. Oil	IE	IE		Included in appropriate 1.B.2.a categories based on 2019 IPCC Refinement

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	
ii. Gas	IE	IE		Included in appropriate 1.B.2.b categories based on 2019 IPCC Refinement

**Table A2.4: Notation keys in the IPPU sector**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	SF <sub>6</sub>	NF <sub>3</sub>	COMMENT
Total Industrial Processes							NO	
2.A. Mineral industry		NO	NO					
2.A.4 Other process uses of carbonates		NO	NO					
2.B Chemical Industry				NO	NO	NO	NO	No F-gases are produced in chemical industry
2.B.3. Adipic acid production	NO		NO					Production of adipic acid is not occurring in Slovakia
2.B.4. Caprolactam, glyoxal and glyoxylic acid production	NO		NO					This production is not occurring in Slovakia
2.B.5. Carbide production		NO						No CH <sub>4</sub> emissions occur
2.B.6. Titanium dioxide production	NO							This production is not occurring in Slovakia
2.B.7. Soda ash production	NO							This production is not occurring in Slovakia
2.B.8. Petrochemical and carbon black production		NA,NO						No CH <sub>4</sub> emissions occur
2.B.9. Fluorochemical production				NO	NO	NO	NO	This production is not occurring in Slovakia
2.B.10. Other (as specified in table 2(I).A-H)	NO	NO	NO	NO	NO	NO	NO	This production is not occurring in Slovakia
2.C Metal Industry				NO		NO	NO	
2.C.3. Aluminium production						NO		No SF <sub>6</sub> emissions occur
2.C.4. Magnesium production	NO			NO	NO	NO		This production is not occurring in Slovakia
2.C.6. Zinc production	NO							This production is not occurring in Slovakia
2.C.7. Other (as specified in table 2(I).A-H)	NO	NO		NO	NO	NO	NO	No sources are occurring in this subcategory
2.D Non-energy Products from Fuels and Solvent Use		NO,NE,NA	NO,NE,NA					Different type of activity data was used for calculation, see NID
2.D.1. Lubricant use		NE	NE					No methodology is available
2.D.2. Paraffin wax use		NE	NE					No methodology is available
2.D.3. Other		NO,NA	NO,NA					No sources are occurring in this subcategory
2.E Electronics Industry			NO	NO	NO	NO	NO	No sources are occurring in this subcategory

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	HFCs	PFCs	SF <sub>6</sub>	NF <sub>3</sub>	COMMENT
2.E.1. Integrated circuit or semiconductor			NO	NO	NO	NO	NO	No sources are occurring in this subcategory
2.E.2. TFT flat panel display			NO	NO	NO	NO	NO	No sources are occurring in this subcategory
2.E.3. Photovoltaics				NO	NO	NO	NO	No sources are occurring in this subcategory
2.E.4. Heat transfer fluid				NO	NO	NO	NO	No sources are occurring in this subcategory
2.E.5. Other (as specified in table 2(III))			NO	NO	NO	NO	NO	No sources are occurring in this subcategory
2.F Product Uses for ODS					NO	NO	NO	These types of gas are not used
2.F.1. Refrigeration and air conditioning					NO	NO	NO	These types of gas are not used
2.F.2. Foam blowing agents					NO	NO	NO	These types of gas are not used
2.F.3. Fire protection					NO	NO	NO	These types of gas are not used
2.F.4. Aerosols					NO	NO	NO	These types of gas are not used
2.F.5. Solvents				NO	NO	NO	NO	No sources are occurring in this subcategory
2.F.6. Other applications				NO	NO	NO	NO	These types of gas are not used
2.G Other Product Manufacture and Use	NO	NO		NO	NO		NO	These types of gas are not used
2.G.1. Electrical equipment				NO	NO		NO	These types of gas are not used
2.G.2. SF <sub>6</sub> and PFCs from other product use					NO	IE		SF <sub>6</sub> emissions are included in G.1 category
2.G.4. Other	NO	NO	NO	NO	NO	NO	NO	No sources are occurring in this subcategory
2.H Other as specified in tables 2(I).A-H and 2(II)	NO	NA	NA	NO	NO	NO	NO	No sources are occurring in this subcategory

**Table A2.5: Notation keys in the Agriculture sector**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	
I. Livestock				
3.C Rice Cultivation		NO		No rice cultivation in Slovakia
3.D Agricultural Soils		NO		
3.D.6. Cultivation of organic soils (i.e. histosols)			NE	Activity is under threshold of significance.
3.D.7. Other			NO	No methodology is available in the IPCC 2006 GL or in the IPCC 2019 RF for N <sub>2</sub> O, CH <sub>4</sub> and N <sub>2</sub> O emissions in this subcategory.
3.E Prescribed Burning of Savannas		NO	NO	No savannahs are occurring in Slovakia.

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	
3.F Field burning of Agricultural Residues		NA, NO	NA, NO	This practise is forbidden by law in Slovakia.
3.I Other Carbon-Containing Fertilizers	NO			No methodology is available in the IPCC 2006 GL or in the IPCC 2019 RF for CO <sub>2</sub> emissions in this subcategory.
3.J Other	NA	NA	NA	No other sources were identified in Slovakia.

**Table A2.6: Notation keys used in the Waste sector**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	EMISSIONS			COMMENT
	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	
5.A Solid Waste Disposal	NO			No CO <sub>2</sub> emissions are reported in waste disposal.
5.A.1. Managed waste disposal sites	NO			NE is reported for amount of CH <sub>4</sub> flared in 2016.
5.A.2. Unmanaged waste disposal sites	NO	NO		Unmanaged waste disposal sites are not occurring in Slovakia
5.A.3. Uncategorized waste disposal sites	NO	NO		No uncategorised sites
5.B Biological Treatment of Solid Waste				No CH <sub>4</sub> emissions are flared as this practise is not occurring in Slovakia
5.B.2. Anaerobic digestion at biogas facilities			NA	Not reported due to a lack of available emission factors. According to the 2006 IPCC Guidelines, Volume 5 Waste page 4.6 Table 4.1 emissions in both basis were assumed as negligible The methane emissions from 5.B.2.a category is included in category 5.B.2.b The methane emissions from 5.B.2 is energy recovered in the category 1.A.5
5.C Incineration and Open Burning of Waste				Biogenic and non-biogenic municipal solid waste incineration is included in energy sector (with energy use incineration, category 1.A.1.a.iv - other fuels).
5.C.2. Open burning of waste	NO	NO	NO	This practise is not occurring in Slovakia.
5.D Wastewater Treatment and Discharge				No CO <sub>2</sub> emissions are reported in wastewater treatment.
5.D.3. Other (as specified in table 6.B)	NO	NO	NO	All sources are included in subcategories 5.D.1 and 5.D.2, therefore no emissions are occurring here.
5.E. Other	NO	NO	NO	No additional emissions sources were identified.

**Table A2.7: Notation keys used in the LULUCF**

GREENHOUSE GAS SOURCE AND SINK CATEGORIES	Net CO <sub>2</sub> emissions/removals	CH <sub>4</sub>	N <sub>2</sub> O	COMMENT
4.B Cropland		NA, NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia CH <sub>4</sub> emissions and removals from drainage and rewetting and other management of organic and mineral soils - this activity not occurring in Slovakia.
4.B.1. Cropland remaining cropland		NA, NO	NO	CH <sub>4</sub> and N <sub>2</sub> O emissions biomass burning not occurring in Slovakia, CH <sub>4</sub> and N <sub>2</sub> O emissions and removals from drainage and rewetting and other management of organic and mineral soils - this activity not occurring in Slovakia. Emissions from histosols are below the threshold, notation key NE was used.
4.B.2. Land converted to cropland		NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia.
4.C Grassland		NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia.
4.C.1. Grassland remaining grassland	NO, NA	NO	NO	CO <sub>2</sub> - tier 1 assumes no change in living biomass, DOM and soil.
4.C.2. Land converted to grassland		NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia.
4.D Wetlands	NO	NO	NO	As permanent surface waters have no carbon stock by definition, no emissions are reported.
4.D.1. Wetlands remaining wetlands	NO	NO	NO	No changes in AD, area remaining constant for reporting period. Wetlands consist of surface waters (watercourses and water bodies).
4.D.2. Land converted to wetlands	NO	NO	NO	No changes in area from and to WE, AD data not exist.
4.E Settlements		NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia.
4.E.1. Settlements remaining settlements	NA, NO	NO	NO	CO <sub>2</sub> - change in living biomass DOM and soil no change. Direct N <sub>2</sub> O emissions from N input not occurring in Slovakia, CH <sub>4</sub> and N <sub>2</sub> O emissions from biomass burning - not occurring in Slovakia.
4.E.2. Land converted to settlements		NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia.
4.F Other Land		NO		CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O emissions biomass burning not occurring in Slovakia.
4.F.2. Land converted to other land		NO		CH <sub>4</sub> emissions biomass burning not occurring in Slovakia.
4.H Other	NO	NO	NO	N <sub>2</sub> O not occurring. This category is not reporting in Slovakia.

## ANNEX 3. ASSESSMENT OF UNCERTAINTY

Chapter 3 of the IPCC 2006 GL (IPCC 2019 Refinements) provides methods for calculation of uncertainty in emissions inventory. As the Slovak Republic reports the results of Approach 1 the reporting is to be carried out using Table 3.2 for uncertainty calculation. The Slovak Republic provide Approach 2 for uncertainty analyses according to the Chapter 3 of the IPCC 2006 GL (IPCC 2019 Refinements) for the complete Energy and IPPU sectors. The methodology and results are described in the appropriate sectoral chapters of this document. Slovakia intends to use hybrid combination of Approaches 1 and 2 in submissions for calculation of total uncertainty of the inventory. **Tables A3.1** and **A3.2** show the result of uncertainty analysis for years 2023 and 1990. According to the ERT recommendation G.3 from the draft ARR 2022, the table provide with the final row with the overall uncertainty levels.

**Table A3.1: Approach 1 uncertainty with LULUCF assessment in 2023 (emissions in Gg of CO<sub>2</sub> eq., uncertainty in %)**

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
2F1	Refrigeration and Air conditioning	F – gases	0.00	398.92	2.10	0.00	2.10	0.00	0.01	0.01	0.00	0.02	0.00
2F2	Foam Blowing Agents	F – gases	0.00	1.73	8.21	0.00	8.21	0.00	0.00	0.00	0.00	0.00	0.00
2F3	Fire Protection	F – gases	0.00	27.20	13.49	0.00	13.49	0.00	0.00	0.00	0.00	0.01	0.00
2F4	Aerosols	F – gases	0.00	10.04	10.08	0.00	10.08	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Liquid Fuels	CO <sub>2</sub>	3 819.21	1 385.82	5.00	3.60	6.16	0.09	0.00	0.02	-0.02	0.15	0.02
1A1	Fuel combustion - Energy Industries - Solid Fuels	CO <sub>2</sub>	12 861.05	2 783.58	2.50	3.60	4.38	0.19	-0.04	0.04	-0.16	0.15	0.05
1A1	Fuel combustion - Energy Industries - Gaseous Fuels	CO <sub>2</sub>	2 293.69	2 026.18	2.50	2.75	3.72	0.07	0.02	0.03	0.04	0.11	0.01
1A1	Fuel combustion - Energy Industries - Other Fossil Fuels	CO <sub>2</sub>	35.61	157.57	5.00	5.00	7.07	0.00	0.00	0.00	0.01	0.02	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO <sub>2</sub>	2 867.64	277.03	5.00	3.60	6.16	0.00	-0.02	0.00	-0.05	0.03	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO <sub>2</sub>	9 028.53	3 181.94	5.00	2.80	5.73	0.42	-0.01	0.05	-0.03	0.35	0.12
1A2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO <sub>2</sub>	3 930.58	1 899.01	2.50	2.75	3.72	0.06	0.00	0.03	0.01	0.10	0.01
1A2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO <sub>2</sub>	200.34	269.06	5.00	5.00	7.07	0.00	0.00	0.00	0.01	0.03	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Peat	CO <sub>2</sub>	0.00	0.00	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
1A3a	Fuel combustion - Domestic Aviation	CO <sub>2</sub>	3.74	1.56	1.00	5.00	5.10	0.00	0.00	0.00	0.00	0.00	0.00
1A3b	Fuel combustion - Road Transportation	CO <sub>2</sub>	4 503.02	7 538.72	1.00	5.00	5.10	1.85	0.09	0.12	0.43	0.17	0.21
1A3c	Fuel combustion - Railways	CO <sub>2</sub>	372.29	81.29	1.00	5.00	5.10	0.00	0.00	0.00	-0.01	0.00	0.00
1A3d	Fuel combustion - Domestic Navigation - Liquid Fuels	CO <sub>2</sub>	0.02	5.32	1.00	5.00	5.10	0.00	0.00	0.00	0.00	0.00	0.00
1A3e	Fuel combustion - Other Transportation	CO <sub>2</sub>	1 813.95	20.93	1.00	5.00	5.10	0.00	-0.01	0.00	-0.06	0.00	0.00
1A4	Fuel combustion - Other Sectors - Liquid Fuels	CO <sub>2</sub>	580.74	213.78	5.00	3.60	6.16	0.00	0.00	0.00	0.00	0.02	0.00
1A4	Fuel combustion - Other Sectors - Solid Fuels	CO <sub>2</sub>	6 852.15	306.43	5.00	4.00	6.40	0.00	-0.04	0.00	-0.17	0.03	0.03
1A4	Fuel combustion - Other Sectors - Gaseous Fuels	CO <sub>2</sub>	3 634.43	3 677.32	2.50	2.75	3.72	0.23	0.03	0.06	0.09	0.20	0.05
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Liquid Fuels	CO <sub>2</sub>	34.99	1.43	5.00	3.60	6.16	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Solid Fuels	CO <sub>2</sub>	216.08	0.48	5.00	4.00	6.40	0.00	0.00	0.00	-0.01	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Gaseous Fuels	CO <sub>2</sub>	154.75	59.16	2.50	2.75	3.72	0.00	0.00	0.00	0.00	0.00	0.00



CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A5	Fuel combustion - Other (Not specified elsewhere) - Mobile- Liquid Fuels	CO <sub>2</sub>	70.04	4.24	5.00	3.60	6.16	0.00	0.00	0.00	0.00	0.00	0.00
1B1	Fugitive emissions from fuels - Solid Fuels	CO <sub>2</sub>	19.76	48.30	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
1B2a	Fugitive emissions from fuels - oil. NG and Other - Oil	CO <sub>2</sub>	39.69	30.65	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2b	Fugitive emissions from fuels - oil. NG and Other - Natural gas	CO <sub>2</sub>	17.18	2.57	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
1B2c	Fugitive emissions from fuels - oil. NG and Other - Venting and flaring	CO <sub>2</sub>	0.23	0.03	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
2A1	Mineral Industry - Cement Production	CO <sub>2</sub>	1 464.50	1 294.23	0.75	1.67	1.83	0.01	0.01	0.02	0.02	0.02	0.00
2A2	Mineral Industry - Lime Production	CO <sub>2</sub>	794.92	475.06	0.66	2.33	2.42	0.00	0.00	0.01	0.00	0.01	0.00
2A3	Mineral Industry - Glass Production	CO <sub>2</sub>	7.88	12.52	1.36	0.00	1.36	0.00	0.00	0.00	0.00	0.00	0.00
2A4	Mineral Industry - Other Process Uses of Carbonates	CO <sub>2</sub>	446.73	210.09	3.16	0.00	3.16	0.00	0.00	0.00	0.00	0.01	0.00
2B1	Chemical Industry - Ammonia Production	CO <sub>2</sub>	331.77	509.92	1.50	3.59	3.89	0.00	0.01	0.01	0.02	0.02	0.00
2B5	Chemical Industry - Carbide Production	CO <sub>2</sub>	0.00	31.82	1.45	7.14	7.29	0.00	0.00	0.00	0.00	0.00	0.00
2B8	Chemical Industry - Petrochemical and Carbon Black Production	CO <sub>2</sub>	428.80	376.47	1.12	13.24	13.29	0.03	0.00	0.01	0.04	0.01	0.00
2B10	Chemical Industry - Other	CO <sub>2</sub>	0.00	0.00	2.00	2.00	2.83	0.00	0.00	0.00	0.00	0.00	0.00
2C1	Metal Industry - Iron and Steel Production	CO <sub>2</sub>	4 167.97	3 733.94	2.23	5.77	6.19	0.67	0.03	0.06	0.17	0.18	0.06
2C2	Metal Industry - Ferroalloys Production	CO <sub>2</sub>	296.74	7.77	1.62	0.00	1.62	0.00	0.00	0.00	0.00	0.00	0.00
2C3	Metal Industry - Aluminium Production	CO <sub>2</sub>	121.32	6.88	1.79	3.36	3.81	0.00	0.00	0.00	0.00	0.00	0.00
2C5	Metal Industry - Lead Production	CO <sub>2</sub>	0.00	0.08	1.50	20.00	20.06	0.00	0.00	0.00	0.00	0.00	0.00
2D	Non-energy Products from Fuels and Solvent Use	CO <sub>2</sub>	50.49	60.21	6.89	22.39	23.43	0.00	0.00	0.00	0.01	0.01	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
3G	Liming	CO <sub>2</sub>	45.73	16.40	3.04	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
3H	Urea Application	CO <sub>2</sub>	15.29	56.07	3.93	50.00	50.15	0.01	0.00	0.00	0.04	0.00	0.00
4A1	Forest Land - Forest Land Remaining Forest Land	CO <sub>2</sub>	-5 999.27	-6 671.90	20.00	82.84	85.22	118.34	-0.06	0.10	-5.19	2.92	35.51
4A2	Forest Land - Land Converted to Forest Land	CO <sub>2</sub>	-2 263.04	-337.60	3.00	57.81	57.88	0.48	0.01	0.01	0.59	0.02	0.34
4B1	Cropland - Cropland Remaining Cropland	CO <sub>2</sub>	-950.94	-689.34	3.00	75.00	75.06	3.34	0.00	0.01	-0.32	0.05	0.10
4B2	Cropland - Land Converted to Cropland	CO <sub>2</sub>	466.51	35.05	3.00	83.58	83.63	0.01	0.00	0.00	-0.22	0.00	0.05
4C2	Grassland - Land Converted to Grassland	CO <sub>2</sub>	-195.77	-27.85	3.00	83.58	83.63	0.01	0.00	0.00	0.08	0.00	0.01
4E2	Settlements - Land Converted to Settlements	CO <sub>2</sub>	96.59	76.94	3.00	86.07	86.13	0.05	0.00	0.00	0.05	0.01	0.00
4F2	Other land - Land Converted to Other Land	CO <sub>2</sub>	293.10	88.78	3.00	86.07	86.13	0.07	0.00	0.00	-0.05	0.01	0.00
4G	Harvested Wood Products	CO <sub>2</sub>	-470.41	-291.90	10.00	50.00	50.99	0.28	0.00	0.00	-0.07	0.06	0.01
5C	Incineration and Open Burning of Waste	CO <sub>2</sub>	6.68	2.38	5.00	31.10	31.50	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Liquid Fuels	CH <sub>4</sub>	3.83	1.37	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Solid Fuels	CH <sub>4</sub>	3.57	0.62	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Gaseous Fuels	CH <sub>4</sub>	1.16	1.01	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Other Fossil Fuels	CH <sub>4</sub>	0.31	1.62	5.00	50.00	50.25	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Biomass	CH <sub>4</sub>	0.63	10.95	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CH <sub>4</sub>	3.12	0.23	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CH <sub>4</sub>	18.22	2.73	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CH <sub>4</sub>	1.97	0.95	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CH <sub>4</sub>	1.79	2.45	5.00	50.00	50.25	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Peat	CH <sub>4</sub>	0.00	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Biomass	CH <sub>4</sub>	3.05	9.07	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A3a	Fuel combustion - Domestic Aviation	CH <sub>4</sub>	0.00	0.00	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A3b	Fuel combustion - Road Transportation	CH <sub>4</sub>	32.63	5.20	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A3c	Fuel combustion - Railways	CH <sub>4</sub>	0.58	0.14	1.00	40.00	40.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3d	Fuel combustion - Domestic Navigation - Liquid Fuels	CH <sub>4</sub>	0.00	0.01	1.00	40.00	40.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3e	Fuel combustion - Other Transportation	CH <sub>4</sub>	0.89	0.01	1.00	40.00	40.01	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Liquid Fuels	CH <sub>4</sub>	1.40	0.83	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	384.16	12.83	3.00	50.00	50.09	0.00	0.00	0.00	-0.12	0.00	0.01
1A4	Fuel combustion - Other Sectors - Gaseous Fuels	CH <sub>4</sub>	9.11	9.16	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	40.46	159.27	3.00	50.00	50.09	0.08	0.00	0.00	0.11	0.01	0.01

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Liquid Fuels	CH <sub>4</sub>	0.04	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Solid Fuels	CH <sub>4</sub>	0.06	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Gaseous Fuels	CH <sub>4</sub>	0.38	0.15	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Biomass	CH <sub>4</sub>	0.00	0.08	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Mobile- Liquid Fuels	CH <sub>4</sub>	0.22	0.01	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1B1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	794.91	133.93	5.00	7.00	8.60	0.00	0.00	0.00	-0.02	0.01	0.00
1B2a	Fugitive emissions from fuels - oil. NG and Other - Oil	CH <sub>4</sub>	13.24	5.96	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2b	Fugitive emissions from fuels - oil. NG and Other - Natural gas	CH <sub>4</sub>	1 387.74	349.21	2.00	5.00	5.39	0.00	0.00	0.01	-0.02	0.02	0.00
1B2c	Fugitive emissions from fuels - oil. NG and Other - Venting and flaring	CH <sub>4</sub>	659.46	74.68	2.00	5.00	5.39	0.00	0.00	0.00	-0.02	0.00	0.00
2B1	Chemical Industry - Ammonia Production	CH <sub>4</sub>	0.30	0.35	1.50	10.00	10.11	0.00	0.00	0.00	0.00	0.00	0.00
2B10	Chemical Industry - Other	CH <sub>4</sub>	0.00	0.00	2.00	10.00	10.20	0.00	0.00	0.00	0.00	0.00	0.00
2C1	Metal Industry - Iron and Steel Production	CH <sub>4</sub>	13.83	15.59	2.23	400.00	400.01	0.05	0.00	0.00	0.06	0.00	0.00
2C2	Metal Industry - Ferroalloys Production	CH <sub>4</sub>	0.00	0.13	1.50	10.00	10.11	0.00	0.00	0.00	0.00	0.00	0.00
3A	Enteric Fermentation	CH <sub>4</sub>	3 120.02	1037.25	13.10	14.91	19.85	0.53	-0.01	0.02	-0.08	0.30	0.09
3B	Manure Management	CH <sub>4</sub>	716.20	101.32	6.50	9.41	11.43	0.00	0.00	0.00	-0.03	0.01	0.00
4A1	Forest Land - Forest Land Remaining Forest Land	CH <sub>4</sub>	12.37	15.14	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
4A2	Forest Land - Land Converted to Forest Land	CH <sub>4</sub>	0.08	0.00	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
5A	Solid Waste Disposal	CH <sub>4</sub>	781.78	1162.62	17.35	20.31	26.71	1.20	0.01	0.02	0.26	0.44	0.26
5B	Biological Treatment of Solid Waste	CH <sub>4</sub>	72.69	209.00	8.42	62.23	62.80	0.22	0.00	0.00	0.17	0.04	0.03
5C	Incineration and Open Burning of Waste	CH <sub>4</sub>	24.88	7.53	5.00	50.00	50.25	0.00	0.00	0.00	0.00	0.00	0.00
5D	Wastewater Treatment and Discharge	CH <sub>4</sub>	181.83	93.10	4.44	31.44	31.75	0.01	0.00	0.00	0.01	0.01	0.00
1A1	Fuel combustion - Energy Industries - Liquid Fuels	N <sub>2</sub> O	7.04	2.23	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Solid Fuels	N <sub>2</sub> O	48.12	6.50	3.00	50.00	50.09	0.00	0.00	0.00	-0.01	0.00	0.00
1A1	Fuel combustion - Energy Industries - Gaseous Fuels	N <sub>2</sub> O	1.09	0.96	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Other Fossil Fuels	N <sub>2</sub> O	0.40	2.04	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Biomass	N <sub>2</sub> O	0.80	13.82	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	N <sub>2</sub> O	6.05	0.43	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	N <sub>2</sub> O	25.54	3.67	3.00	50.00	50.09	0.00	0.00	0.00	-0.01	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	N <sub>2</sub> O	1.86	0.90	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	N <sub>2</sub> O	2.26	3.09	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Peat	N <sub>2</sub> O	0.00	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A2	Fuel combustion - Manufacturing Industries and Construction - Biomass	N <sub>2</sub> O	3.84	16.26	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A3a	Fuel combustion - Domestic Aviation	N <sub>2</sub> O	0.03	0.01	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3b	Fuel combustion - Road Transportation	N <sub>2</sub> O	50.23	74.08	1.00	50.00	50.01	0.02	0.00	0.00	0.04	0.00	0.00
1A3c	Fuel combustion - Railways	N <sub>2</sub> O	38.08	8.22	1.00	50.00	50.01	0.00	0.00	0.00	-0.01	0.00	0.00
1A3d	Fuel combustion - Domestic Navigation - Liquid Fuels	N <sub>2</sub> O	0.00	0.04	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3e	Fuel combustion - Other Transportation	N <sub>2</sub> O	0.84	0.01	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Liquid Fuels	N <sub>2</sub> O	8.88	15.74	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A4	Fuel combustion - Other Sectors - Solid Fuels	N <sub>2</sub> O	24.59	1.23	3.00	50.00	50.09	0.00	0.00	0.00	-0.01	0.00	0.00
1A4	Fuel combustion - Other Sectors - Gaseous Fuels	N <sub>2</sub> O	1.73	1.73	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Biomass	N <sub>2</sub> O	5.57	21.19	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Liquid Fuels	N <sub>2</sub> O	0.07	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Solid Fuels	N <sub>2</sub> O	0.80	0.00	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Gaseous Fuels	N <sub>2</sub> O	0.07	0.03	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Biomass	N <sub>2</sub> O	0.00	0.01	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Mobile- Liquid Fuels	N <sub>2</sub> O	1.47	0.09	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1B1	Fugitive emissions from fuels - Solid Fuels	N <sub>2</sub> O	0.00	0.08	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1B2a	Fugitive emissions from fuels - oil, NG and Other - Oil	N <sub>2</sub> O	0.14	0.12	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	N <sub>2</sub> O	0.03	0.00	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
2B1	Chemical Industry - Ammonia Production	N <sub>2</sub> O	0.29	0.33	1.50	10.00	10.11	0.00	0.00	0.00	0.00	0.00	0.00
2B2	Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	1 072.65	46.73	2.57	0.00	2.57	0.00	-0.01	0.00	0.00	0.00	0.00
2B10	Chemical Industry - Other	N <sub>2</sub> O	0.00	0.00	2.00	10.00	10.20	0.00	0.00	0.00	0.00	0.00	0.00
2C7	Metal Industry - Other	N <sub>2</sub> O	0.92	1.18	2.23	400.00	400.01	0.00	0.00	0.00	0.00	0.00	0.00
2G3	Other Product Manufacture and Use	N <sub>2</sub> O	14.58	56.71	9.13	0.00	9.13	0.00	0.00	0.00	0.00	0.01	0.00
3B	Manure Management	N <sub>2</sub> O	560.67	207.53	6.50	248.03	248.11	3.31	0.00	0.00	-0.15	0.03	0.02
3D1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	801.87	460.51	59.37	36.22	69.54	1.28	0.00	0.01	0.06	0.60	0.36
3D2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	611.26	302.75	77.10	103.30	128.90	1.90	0.00	0.00	0.06	0.51	0.26
4A1	Forest Land - Forest Land Remaining Forest Land	N <sub>2</sub> O	6.48	7.93	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
4A2	Forest Land - Land Converted to Forest Land	N <sub>2</sub> O	0.04	0.00	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
4B2	Cropland - Land Converted to Cropland	N <sub>2</sub> O	94.82	7.84	75.00	100.00	125.00	0.00	0.00	0.00	-0.05	0.01	0.00
4C2	Grassland - Land Converted to Grassland	N <sub>2</sub> O	1.10	0.27	75.00	100.00	125.00	0.00	0.00	0.00	0.00	0.00	0.00
4E2	Settlements - Land Converted to Settlements	N <sub>2</sub> O	4.88	5.44	75.00	100.00	125.00	0.00	0.00	0.00	0.01	0.01	0.00
4F2	Other land - Land Converted to Other Land	N <sub>2</sub> O	10.95	5.57	75.00	100.00	125.00	0.00	0.00	0.00	0.00	0.01	0.00
5B	Biological Treatment of Solid Waste	N <sub>2</sub> O	41.29	103.50	8.42	93.34	93.72	0.12	0.00	0.00	0.12	0.02	0.02

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
5C	Incineration and Open Burning of Waste	N <sub>2</sub> O	0.39	0.12	5.00	100.00	100.12	0.00	0.00	0.00	0.00	0.00	0.00
5D	Wastewater Treatment and Discharge	N <sub>2</sub> O	75.12	93.62	6.74	31.44	32.16	0.01	0.00	0.00	0.03	0.01	0.00
2C3	Metal Industry - Aluminium Production	PFCs	213.92	0.01	1.50	10.05	10.16	0.00	0.00	0.00	-0.01	0.00	0.00
2F1	Refrigeration and air conditioning	PFCs	0.00	0.00	2.10	11.00	11.20	0.00	0.00	0.00	0.00	0.00	0.00
2G1	Electrical equipment	SF <sub>6</sub>	0.06	14.70	2.00	0.00	2.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL			64 533.04	28 298.03		134.97							37.71
<b>Total Uncertainties</b>			<b>-56.15</b>	<b>Uncertainty in total inventory %:</b>		<b>11.62</b>					<b>Trend uncertainty %:</b>		<b>6.14</b>

**Table A3.2: Approach 1 uncertainty with LULUCF assessment in 1990 (emissions in Gg of CO<sub>2</sub> eq., uncertainty in %)**

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
2F1	Refrigeration and Air conditioning	F – gases	0.00	398.92	2.10	0.00	2.10	0.00	0.01	0.01	0.00	0.02	0.00
2F2	Foam Blowing Agents	F – gases	0.00	1.73	8.21	0.00	8.21	0.00	0.00	0.00	0.00	0.00	0.00
2F3	Fire Protection	F – gases	0.00	27.20	13.49	0.00	13.49	0.00	0.00	0.00	0.00	0.01	0.00
2F4	Aerosols	F – gases	0.00	10.04	10.08	0.00	10.08	0.00	0.00	0.00	0.00	0.00	0.00



CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A1	Fuel combustion - Energy Industries - Liquid Fuels	CO2	3819.21	1385.82	5.00	3.60	6.16	0.13	0.00	0.02	-0.02	0.15	0.02
1A1	Fuel combustion - Energy Industries - Solid Fuels	CO2	12861.05	2783.58	2.50	3.60	4.38	0.76	-0.04	0.04	-0.16	0.15	0.05
1A1	Fuel combustion - Energy Industries - Gaseous Fuels	CO2	2293.69	2026.18	2.50	2.75	3.72	0.02	0.02	0.03	0.04	0.11	0.01
1A1	Fuel combustion - Energy Industries - Other Fossil Fuels	CO2	35.61	157.57	5.00	5.00	7.07	0.00	0.00	0.00	0.01	0.02	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CO2	2867.64	277.03	5.00	3.60	6.16	0.07	-0.02	0.00	-0.05	0.03	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CO2	9028.53	3181.94	5.00	2.80	5.73	0.64	-0.01	0.05	-0.03	0.35	0.12
1A2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CO2	3930.58	1899.01	2.50	2.75	3.72	0.05	0.00	0.03	0.01	0.10	0.01
1A2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CO2	200.34	269.06	5.00	5.00	7.07	0.00	0.00	0.00	0.01	0.03	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Peat	CO2	0.00	0.00	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
1A3a	Fuel combustion - Domestic Aviation	CO2	3.74	1.56	1.00	5.00	5.10	0.00	0.00	0.00	0.00	0.00	0.00
1A3b	Fuel combustion - Road Transportation	CO2	4 503.02	7 538.72	1.00	5.00	5.10	0.13	0.09	0.12	0.43	0.17	0.21
1A3c	Fuel combustion - Railways	CO2	372.29	81.29	1.00	5.00	5.10	0.00	0.00	0.00	-0.01	0.00	0.00
1A3d	Fuel combustion - Domestic Navigation - Liquid Fuels	CO2	0.02	5.32	1.00	5.00	5.10	0.00	0.00	0.00	0.00	0.00	0.00
1A3e	Fuel combustion - Other Transportation	CO2	1 813.95	20.93	1.00	5.00	5.10	0.02	-0.01	0.00	-0.06	0.00	0.00
1A4	Fuel combustion - Other Sectors - Liquid Fuels	CO2	580.74	213.78	5.00	3.60	6.16	0.00	0.00	0.00	0.00	0.02	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A4	Fuel combustion - Other Sectors - Solid Fuels	CO2	6 852.15	306.43	5.00	4.00	6.40	0.46	-0.04	0.00	-0.17	0.03	0.03
1A4	Fuel combustion - Other Sectors - Gaseous Fuels	CO2	3 634.43	3 677.32	2.50	2.75	3.72	0.04	0.03	0.06	0.09	0.20	0.05
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Liquid Fuels	CO2	34.99	1.43	5.00	3.60	6.16	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Solid Fuels	CO2	216.08	0.48	5.00	4.00	6.40	0.00	0.00	0.00	-0.01	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Gaseous Fuels	CO2	154.75	59.16	2.50	2.75	3.72	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Mobile- Liquid Fuels	CO2	70.04	4.24	5.00	3.60	6.16	0.00	0.00	0.00	0.00	0.00	0.00
1B1	Fugitive emissions from fuels - Solid Fuels	CO2	19.76	48.30	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
1B2a	Fugitive emissions from fuels - oil. NG and Other - Oil	CO2	39.69	30.65	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2b	Fugitive emissions from fuels - oil. NG and Other - Natural gas	CO2	17.18	2.57	2.00	5.00	5.39	0.00	0.00	0.00	0.00	0.00	0.00
1B2c	Fugitive emissions from fuels - oil. NG and Other - Venting and flaring	CO2	0.23	0.03	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
2A1	Mineral Industry - Cement Production	CO2	1 464.50	1 294.23	0.75	1.67	1.83	0.00	0.01	0.02	0.02	0.02	0.00
2A2	Mineral Industry - Lime Production	CO2	794.92	475.06	0.66	2.33	2.42	0.00	0.00	0.01	0.00	0.01	0.00
2A3	Mineral Industry - Glass Production	CO2	7.88	12.52	1.36	0.00	1.36	0.00	0.00	0.00	0.00	0.00	0.00
2A4	Mineral Industry - Other Process Uses of Carbonates	CO2	446.73	210.09	3.16	0.00	3.16	0.00	0.00	0.00	0.00	0.01	0.00
2B1	Chemical Industry - Ammonia Production	CO2	331.77	509.92	1.50	3.59	3.89	0.00	0.01	0.01	0.02	0.02	0.00
2B5	Chemical Industry - Carbide Production	CO2	0.00	31.82	1.45	7.14	7.29	0.00	0.00	0.00	0.00	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
2B8	Chemical Industry - Petrochemical and Carbon Black Production	CO2	428.80	376.47	1.12	13.24	13.29	0.01	0.00	0.01	0.04	0.01	0.00
2B10	Chemical Industry - Other	CO2	0.00	0.00	2.00	2.00	2.83	0.00	0.00	0.00	0.00	0.00	0.00
2C1	Metal Industry - Iron and Steel Production	CO2	4 167.97	3 733.94	2.23	5.77	6.19	0.16	0.03	0.06	0.17	0.18	0.06
2C2	Metal Industry - Ferroalloys Production	CO2	296.74	7.77	1.62	0.00	1.62	0.00	0.00	0.00	0.00	0.00	0.00
2C3	Metal Industry - Aluminium Production	CO2	121.32	6.88	1.79	3.36	3.81	0.00	0.00	0.00	0.00	0.00	0.00
2C5	Metal Industry - Lead Production	CO2	0.00	0.08	1.50	20.00	20.06	0.00	0.00	0.00	0.00	0.00	0.00
2D	Non-energy Products from Fuels and Solvent Use	CO2	50.49	60.21	6.89	22.39	23.43	0.00	0.00	0.00	0.01	0.01	0.00
3G	Liming	CO2	45.73	16.40	3.04	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
3H	Urea Application	CO2	15.29	56.07	3.93	50.00	50.15	0.00	0.00	0.00	0.04	0.00	0.00
4A1	Forest Land - Forest Land Remaining Forest Land	CO2	-5 999.27	-6 671.90	20.00	82.84	85.22	62.76	-0.06	0.10	-5.19	2.92	35.51
4A2	Forest Land - Land Converted to Forest Land	CO2	-2 263.04	-337.60	3.00	57.81	57.88	4.12	0.01	0.01	0.59	0.02	0.34
4B1	Cropland - Cropland Remaining Cropland	CO2	-950.94	-689.34	3.00	75.00	75.06	1.22	0.00	0.01	-0.32	0.05	0.10
4B2	Cropland - Land Converted to Cropland	CO2	466.51	35.05	3.00	83.58	83.63	0.37	0.00	0.00	-0.22	0.00	0.05
4C2	Grassland - Land Converted to Grassland	CO2	-195.77	-27.85	3.00	83.58	83.63	0.06	0.00	0.00	0.08	0.00	0.01
4E2	Settlements - Land Converted to Settlements	CO2	96.59	76.94	3.00	86.07	86.13	0.02	0.00	0.00	0.05	0.01	0.00
4F2	Other land - Land Converted to Other Land	CO2	293.10	88.78	3.00	86.07	86.13	0.15	0.00	0.00	-0.05	0.01	0.00
4G	Harvested Wood Products	CO2	-470.41	-291.90	10.00	50.00	50.99	0.14	0.00	0.00	-0.07	0.06	0.01
5C	Incineration and Open Burning of Waste	CO2	6.68	2.38	5.00	31.10	31.50	0.00	0.00	0.00	0.00	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A1	Fuel combustion - Energy Industries - Liquid Fuels	CH <sub>4</sub>	3.83	1.37	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Solid Fuels	CH <sub>4</sub>	3.57	0.62	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Gaseous Fuels	CH <sub>4</sub>	1.16	1.01	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Other Fossil Fuels	CH <sub>4</sub>	0.31	1.62	5.00	50.00	50.25	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Biomass	CH <sub>4</sub>	0.63	10.95	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	CH <sub>4</sub>	3.12	0.23	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	CH <sub>4</sub>	18.22	2.73	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	CH <sub>4</sub>	1.97	0.95	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	CH <sub>4</sub>	1.79	2.45	5.00	50.00	50.25	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Peat	CH <sub>4</sub>	0.00	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Biomass	CH <sub>4</sub>	3.05	9.07	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A3a	Fuel combustion - Domestic Aviation	CH <sub>4</sub>	0.00	0.00	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A3b	Fuel combustion - Road Transportation	CH <sub>4</sub>	32.63	5.20	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A3c	Fuel combustion - Railways	CH <sub>4</sub>	0.58	0.14	1.00	40.00	40.01	0.00	0.00	0.00	0.00	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A3d	Fuel combustion - Domestic Navigation - Liquid Fuels	CH <sub>4</sub>	0.00	0.01	1.00	40.00	40.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3e	Fuel combustion - Other Transportation	CH <sub>4</sub>	0.89	0.01	1.00	40.00	40.01	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Liquid Fuels	CH <sub>4</sub>	1.40	0.83	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Solid Fuels	CH <sub>4</sub>	384.16	12.83	3.00	50.00	50.09	0.09	0.00	0.00	-0.12	0.00	0.01
1A4	Fuel combustion - Other Sectors - Gaseous Fuels	CH <sub>4</sub>	9.11	9.16	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Biomass	CH <sub>4</sub>	40.46	159.27	3.00	50.00	50.09	0.00	0.00	0.00	0.11	0.01	0.01
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Liquid Fuels	CH <sub>4</sub>	0.04	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Solid Fuels	CH <sub>4</sub>	0.06	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Gaseous Fuels	CH <sub>4</sub>	0.38	0.15	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Biomass	CH <sub>4</sub>	0.00	0.08	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Mobile- Liquid Fuels	CH <sub>4</sub>	0.22	0.01	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1B1	Fugitive emissions from fuels - Solid Fuels	CH <sub>4</sub>	794.91	133.93	5.00	7.00	8.60	0.01	0.00	0.00	-0.02	0.01	0.00
1B2a	Fugitive emissions from fuels - oil, NG and Other - Oil	CH <sub>4</sub>	13.24	5.96	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2b	Fugitive emissions from fuels - oil, NG and Other - Natural gas	CH <sub>4</sub>	1 387.74	349.21	2.00	5.00	5.39	0.01	0.00	0.01	-0.02	0.02	0.00
1B2c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	CH <sub>4</sub>	659.46	74.68	2.00	5.00	5.39	0.00	0.00	0.00	-0.02	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
2B1	Chemical Industry - Ammonia Production	CH <sub>4</sub>	0.30	0.35	1.50	10.00	10.11	0.00	0.00	0.00	0.00	0.00	0.00
2B10	Chemical Industry - Other	CH <sub>4</sub>	0.00	0.00	2.00	10.00	10.20	0.00	0.00	0.00	0.00	0.00	0.00
2C1	Metal Industry - Iron and Steel Production	CH <sub>4</sub>	13.83	15.59	2.23	400.00	400.01	0.01	0.00	0.00	0.06	0.00	0.00
2C2	Metal Industry - Ferroalloys Production	CH <sub>4</sub>	0.00	0.13	1.50	10.00	10.11	0.00	0.00	0.00	0.00	0.00	0.00
3A	Enteric Fermentation	CH <sub>4</sub>	3 120.02	1 037.25	13.10	14.91	19.85	0.92	-0.01	0.02	-0.08	0.30	0.09
3B	Manure Management	CH <sub>4</sub>	716.20	101.32	6.50	9.41	11.43	0.02	0.00	0.00	-0.03	0.01	0.00
4A1	Forest Land - Forest Land Remaining Forest Land	CH <sub>4</sub>	12.37	15.14	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
4A2	Forest Land - Land Converted to Forest Land	CH <sub>4</sub>	0.08	0.00	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
5A	Solid Waste Disposal	CH <sub>4</sub>	781.78	1162.62	17.35	20.31	26.71	0.10	0.01	0.02	0.26	0.44	0.26
5B	Biological Treatment of Solid Waste	CH <sub>4</sub>	72.69	209.00	8.42	62.23	62.80	0.01	0.00	0.00	0.17	0.04	0.03
5C	Incineration and Open Burning of Waste	CH <sub>4</sub>	24.88	7.53	5.00	50.00	50.25	0.00	0.00	0.00	0.00	0.00	0.00
5D	Wastewater Treatment and Discharge	CH <sub>4</sub>	181.83	93.10	4.44	31.44	31.75	0.01	0.00	0.00	0.01	0.01	0.00
1A1	Fuel combustion - Energy Industries - Liquid Fuels	N <sub>2</sub> O	7.04	2.23	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Solid Fuels	N <sub>2</sub> O	48.12	6.50	3.00	50.00	50.09	0.00	0.00	0.00	-0.01	0.00	0.00
1A1	Fuel combustion - Energy Industries - Gaseous Fuels	N <sub>2</sub> O	1.09	0.96	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Other Fossil Fuels	N <sub>2</sub> O	0.40	2.04	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A1	Fuel combustion - Energy Industries - Biomass	N <sub>2</sub> O	0.80	13.82	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A2	Fuel combustion - Manufacturing Industries and Construction - Liquid Fuels	N <sub>2</sub> O	6.05	0.43	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Solid Fuels	N <sub>2</sub> O	25.54	3.67	3.00	50.00	50.09	0.00	0.00	0.00	-0.01	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Gaseous Fuels	N <sub>2</sub> O	1.86	0.90	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Other Fossil Fuels	N <sub>2</sub> O	2.26	3.09	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Peat	N <sub>2</sub> O	0.00	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A2	Fuel combustion - Manufacturing Industries and Construction - Biomass	N <sub>2</sub> O	3.84	16.26	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A3a	Fuel combustion - Domestic Aviation	N <sub>2</sub> O	0.03	0.01	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3b	Fuel combustion - Road Transportation	N <sub>2</sub> O	50.23	74.08	1.00	50.00	50.01	0.00	0.00	0.00	0.04	0.00	0.00
1A3c	Fuel combustion - Railways	N <sub>2</sub> O	38.08	8.22	1.00	50.00	50.01	0.00	0.00	0.00	-0.01	0.00	0.00
1A3d	Fuel combustion - Domestic Navigation - Liquid Fuels	N <sub>2</sub> O	0.00	0.04	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A3e	Fuel combustion - Other Transportation	N <sub>2</sub> O	0.84	0.01	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Liquid Fuels	N <sub>2</sub> O	8.88	15.74	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00
1A4	Fuel combustion - Other Sectors - Solid Fuels	N <sub>2</sub> O	24.59	1.23	3.00	50.00	50.09	0.00	0.00	0.00	-0.01	0.00	0.00
1A4	Fuel combustion - Other Sectors - Gaseous Fuels	N <sub>2</sub> O	1.73	1.73	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A4	Fuel combustion - Other Sectors - Biomass	N <sub>2</sub> O	5.57	21.19	3.00	50.00	50.09	0.00	0.00	0.00	0.01	0.00	0.00

CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Liquid Fuels	N <sub>2</sub> O	0.07	0.00	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Solid Fuels	N <sub>2</sub> O	0.80	0.00	3.00	5.00	5.83	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Gaseous Fuels	N <sub>2</sub> O	0.07	0.03	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Stationary - Biomass	N <sub>2</sub> O	0.00	0.01	1.00	50.00	50.01	0.00	0.00	0.00	0.00	0.00	0.00
1A5	Fuel combustion - Other (Not specified elsewhere) - Mobile- Liquid Fuels	N <sub>2</sub> O	1.47	0.09	3.00	50.00	50.09	0.00	0.00	0.00	0.00	0.00	0.00
1B1	Fugitive emissions from fuels - Solid Fuels	N <sub>2</sub> O	0.00	0.08	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2a	Fugitive emissions from fuels - oil, NG and Other - Oil	N <sub>2</sub> O	0.14	0.12	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
1B2c	Fugitive emissions from fuels - oil, NG and Other - Venting and flaring	N <sub>2</sub> O	0.03	0.00	5.00	7.00	8.60	0.00	0.00	0.00	0.00	0.00	0.00
2B1	Chemical Industry - Ammonia Production	N <sub>2</sub> O	0.29	0.33	1.50	10.00	10.11	0.00	0.00	0.00	0.00	0.00	0.00
2B2	Chemical Industry - Nitric Acid Production	N <sub>2</sub> O	1 072.65	46.73	2.57	0.00	2.57	0.00	-0.01	0.00	0.00	0.00	0.00
2B10	Chemical Industry - Other	N <sub>2</sub> O	0.00	0.00	2.00	10.00	10.20	0.00	0.00	0.00	0.00	0.00	0.00
2C7	Metal Industry - Other	N <sub>2</sub> O	0.92	1.18	2.23	400.00	400.01	0.00	0.00	0.00	0.00	0.00	0.00
2G3	Other Product Manufacture and Use	N <sub>2</sub> O	14.58	56.71	9.13	0.00	9.13	0.00	0.00	0.00	0.00	0.01	0.00
3B	Manure Management	N <sub>2</sub> O	560.67	207.53	6.50	248.03	248.11	4.65	0.00	0.00	-0.15	0.03	0.02
3D1	Direct N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	801.87	460.51	59.37	36.22	69.54	0.75	0.00	0.01	0.06	0.60	0.36
3D2	Indirect N <sub>2</sub> O Emissions From Managed Soils	N <sub>2</sub> O	611.26	302.75	77.10	103.30	128.90	1.49	0.00	0.00	0.06	0.51	0.26



CRT	IPCC Category	GAS	BASE YEAR EMISSIONS OR REMOVALS	YEAR 2023 EMISSIONS OR REMOVALS	ACTIVITY DATA UNCERTAINTY	EMISSION FACTOR UNCERTAINTY	COMBINED UNCERTAINTY	CONTRIBUTION TO VARIANCE BY CATEGORY IN YEAR 2023	TYPE A SENSITIVITY	TYPE B SENSITIVITY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY EMISSION FACTOR / ESTIMATION PARAMETER UNCERTAINTY	UNCERTAINTY IN TREND IN NATIONAL EMISSIONS INTRODUCED BY ACTIVITY DATA UNCERTAINTY	UNCERTAINTY INTRODUCED INTO THE TREND IN TOTAL NATIONAL EMISSIONS
4A1	Forest Land - Forest Land Remaining Forest Land	N <sub>2</sub> O	6.48	7.93	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
4A2	Forest Land - Land Converted to Forest Land	N <sub>2</sub> O	0.04	0.00	5.00	5.00	7.07	0.00	0.00	0.00	0.00	0.00	0.00
4B2	Cropland - Land Converted to Cropland	N <sub>2</sub> O	94.82	7.84	75.00	100.00	125.00	0.03	0.00	0.00	-0.05	0.01	0.00
4C2	Grassland - Land Converted to Grassland	N <sub>2</sub> O	1.10	0.27	75.00	100.00	125.00	0.00	0.00	0.00	0.00	0.00	0.00
4E2	Settlements - Land Converted to Settlements	N <sub>2</sub> O	4.88	5.44	75.00	100.00	125.00	0.00	0.00	0.00	0.01	0.01	0.00
4F2	Other land - Land Converted to Other Land	N <sub>2</sub> O	10.95	5.57	75.00	100.00	125.00	0.00	0.00	0.00	0.00	0.01	0.00
5B	Biological Treatment of Solid Waste	N <sub>2</sub> O	41.29	103.50	8.42	93.34	93.72	0.00	0.00	0.00	0.12	0.02	0.02
5C	Incineration and Open Burning of Waste	N <sub>2</sub> O	0.39	0.12	5.00	100.00	100.12	0.00	0.00	0.00	0.00	0.00	0.00
5D	Wastewater Treatment and Discharge	N <sub>2</sub> O	75.12	93.62	6.74	31.44	32.16	0.00	0.00	0.00	0.03	0.01	0.00
2C3	Metal Industry - Aluminium Production	PFCs	213.92	0.01	1.50	10.05	10.16	0.00	0.00	0.00	-0.01	0.00	0.00
2F1	Refrigeration and air conditioning	PFCs	0.00	0.00	2.10	11.00	11.20	0.00	0.00	0.00	0.00	0.00	0.00
2G1	Electrical equipment	SF <sub>6</sub>	0.06	14.70	2.00	0.00	2.00	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL			64 533.04	28 298.03		79.47							37.71
<b>Total Uncertainties</b>			<b>-56.15</b>	<b>Uncertainty in total inventory %:</b>		<b>8,91</b>					<b>Trend uncertainty %:</b>		<b>6.14</b>

## ANNEX 4. QUALITY ASSURANCE/QUALITY CONTROL PLAN

**Table A4.1: The Quality Assurance/Quality Control Plan 2025 - Internal**

ACTIVITY		WHO	CHECK-IN	TIME SCHEDULE	RECORD
1.	Evaluation of Improvement plans for the year 2025	Sectoral experts NS coordinator Deputy of NS coordinator	Quality manager MŽP SR – NFP	15.01.2025	Improvement plan for the year 2025 for every sector
2.	Tasks and financial plan of NS – preparation for the year 2025.	NS coordinator Deputy of NS coordinator	MŽP SR – NFP Quality manager Head of the SHMÚ	12.02.2025	Information on budget, capacity (personal, external, internal), training plan, meetings and business trips plan, plan of QA/QC activities for the inventory year 2023.
3.	Update of capacity incorporating updates for each sector	Sectoral experts (SE) Deputy of SE	MŽP SR – NFP Quality manager Head of the SHMÚ	28.02.2025	Responsibilities matrix for 2025 Description of work activities
4.	Work assignment and contracts signing for each sector for the year 2025	NS coordinator Deputy of NS coordinator	MŽP SR - NFP Head of the SHMÚ	31.03.2025	Frame contracts with the sectoral experts Specification of tasks for a given year (improvement plan) Nomination letters for sectoral experts
5.	Plan of QA/QC activities for the emission inventory on overall and sectoral level	Sectoral experts (SE) Deputy of SE	NS coordinator Deputy of NS coordinator Quality manager	10.03.2025	Description QA/QC activities in each sectoral chapters for the year 2025
6.	Key sources and uncertainty management for each sector for the inventory year 2023	Sectoral expert for uncertainty Sectoral experts NS coordinator	Deputy of NS coordinator Quality manager	15.03.2025	Report on key sources and uncertainty evaluation for year 2023 Template for the key sources and uncertainty evaluation for year 2023
7.	Final evaluation of emission data on sectoral level based on the external audit of the European Commission	Sectoral experts NS coordinator	Deputy of NS coordinator Quality manager MŽP SR – NFP	31.05.2025	Verification protocols Description of changes Updated sectoral report
8.	Workshop – meeting of experts, ministries, SNE; Program: evaluation of results, finding from the reviews, proposals for improvement, proposal for the inventory plan for NID 2026	Sectoral experts NS coordinator Deputy of NS coordinator	MŽP SR – NFP Quality manager	April 2025 September 2025 December 2025	Report from the meeting
9.	Completeness check of emission inventory for the year 2025	Sectoral experts	NS coordinator Deputy of NS coordinator	30.09.2025	Report from completeness check

ACTIVITY	WHO	CHECK-IN	TIME SCHEDULE	RECORD	
		Quality manager MŽP SR – NFP			
10.	Methodical updates, recalculation list on sectoral level, according to IPCC 2006 GL	Sectoral experts	NS coordinator Deputy of NS coordinator Quality manager	31.10. 2025	Report of emission for each sector, for inventory year 2024
11.	Sectoral final documents delivery	Sectoral experts	NS coordinator Deputy of NS coordinator Quality manager	30.11. 2025	Delivery protocols Drafts of sectoral reports for the inventory year 2024
12.	Participation in individual evaluations and cooperation in preparing of view on the review assessment by the UNFCCC secretariat	Sectoral experts	NS coordinator Deputy of NS coordinator Quality manager	continuously	Sectoral assessment reports

**Table A4.2: The Quality Assurance/Quality Control Plan 2025 - External**

ACTIVITY	WHO	CHECK-IN	TIME SCHEDULE	RECORD	
1.	Annual Report 2024 submission according to the Regulation (EÚ) 2018/1999 Article 26 and Implementing regulation 2020/1208/EU, Article 7: - Preliminary Emission GHG inventory for years 2023; - Indicators for the year 2023; - Preliminary National Inventory 2025; - SEF tables for the year 2024.	NS coordinator Sectoral experts National administrator	Ministry of Environment of the Slovak Republic – NFP Deputy of NS coordinator	15. 01. 2025	Annual Report SVK 2023 - complete Components of the SVK NID 2025 - incomplete CRT 1990 - 2023 SEF tables 2024 Tables according directive (EU) 2020/1208
2.	Repeated Annual Report 2025 submission according to the Regulation 2018/1999/EU, Article 26 and Implementing regulation 2020/1208/EU, Article 8-24:	NS coordinator Sectoral experts National administrator	Ministry of Environment of the Slovak Republic – NFP Deputy of NS coordinator	15. 03. 2025	Indicators form for the year 2023 CRT tables 1990-2023 SVK NID 2025 - final Tables according directive (EU) 2020/1208

ACTIVITY	WHO	CHECK-IN	TIME SCHEDULE	RECORD
- Emission GHG inventory for year 2023; - Indicators for the year 2023; - National Inventory Document for year 2025; - SEF tables for the year 2024.				
3. ESR comprehensive review 2025	NS coordinator Deputy of NS coordinator Sectoral experts	Technical Expert Review Team	15. 02. 2025 20. 04. 2025	Report from the review until 30. 06. 2025 (depending on the findings and their solution)
4. Nomination letters for the sectoral experts – update for the year 2025.	Ministry of Environment of the Slovak Republic – NFP	Deputy of NS coordinator	15. 04. 2025	Nomination Letters List of nominated sectoral experts for the year 2025.
5. National Inventory SVK NID 2025 submission to the secretariat UNFCCC by ETF software: - Emission GHG inventory for the years 1990-2023; - National Inventory Document 2025; - Information from the National Registry for the year 2024.	NS coordinator Sectoral experts National Registry	Deputy of NS coordinator Ministry of Environment of the Slovak Republic – NFP	15. 04. 2025	CRT tables 1990-2023 SEF 2024 SVK NID 2025 published on the official web of the UNFCCC
6. Publicity of the SVK NID 2025 and emissions data on the official web of the SVK NS.	NS coordinator Deputy of NS coordinator	Ministry of Environment of the Slovak Republic – NFP	15. 04. 2025	Update of data on <a href="https://oeab.shmu.sk/">https://oeab.shmu.sk/</a>
7. Completion and updating of the SVK NID 2025 on the basis of Initial Assessment by the EU review.	NS coordinator Sectoral experts	Deputy of NS coordinator Ministry of Environment of the Slovak Republic – NFP	6 weeks after 15. 04. 2025	Repeated Emission GHG inventory and SVK NID 2025 submission (if relevant)
8. UNFCCC in-country review on the first BTR 2024	MŽP SR		8.	UNFCCC in-country review on the first BTR 2024
9. Audit of the status of the preparation of the emission GHG inventory for the year 2024 – check days.	NS coordinator Sectoral experts	Deputy of NS coordinator Ministry of Environment of the Slovak Republic – NFP	30. 06. 2025 30. 09. 2025	Report from the coordination meetings of the NS
10. Proxy Inventory SVK 2024 according Regulation	NS coordinator	Deputy of NS coordinator Ministry of	31. 07. 2025	Proxy inventory of GHG

ACTIVITY	WHO	CHECK-IN	TIME SCHEDULE	RECORD
2018/1999/EU, Article 26 and Implementing regulation 2020/1208/EU, Article 7	Sectoral experts	Environment of the Slovak Republic – NFP		
11. Data delivering to the Statistical Office of the Slovak Republic. Distribution of the SVK NID 2025 to the relevant institutions.	NS coordinator Sectoral experts	Deputy of NS coordinator Ministry of Environment of the Slovak Republic – NFP	31. 10. 2025	Statistical record Emission GHG inventory for the years 1990-2023
12. Measures and objectives for improvements in QA/QC procedure of GHG emission inventory for relevant sectors based on the preliminary results of the review NID SVK 2025.	Sectoral experts Deputy of NS coordinator	NS coordinator Ministry of Environment of the Slovak Republic – NFP	30. 11. 2025	Report and Improvement plan for the year 2026

**Table A4.3:** List of UNFCCC main findings and recommendations, and status of implementation

CRT CATEGORY / ISSUE	REVIEW RECOMMENDATION	REVIEW REPORT / PARAGRAPH	MS RESPONSE / STATUS OF IMPLEMENTATION	CHAPTER/SECTION IN THE NID
Energy - 1.A.4 Other sectors – solid fuels – CH <sub>4</sub> , (E.6, 2021) (E.17, 2019), (E.36, 2017) Accuracy	Estimate and report CH <sub>4</sub> emissions from solid fuels for category 1.A.4 using at least a tier 2 methodology (in accordance with the 2006 IPCC Guidelines) if the emissions are identified as key.	E.2 - ARR 2022 (sent on 4. 4. 2023)	Partly Implemented. Complete will be possible in the future submission due to capacity and budget constraints and difficulties connected with the EFs estimation in services and households.	Implemented in the SVK NIR 2023 (Chapters 3.2.5 & 3.2.9)
LULUCF - 4. General (LULUCF) – CO <sub>2</sub> (L.1, 2021) (L.1, 2019)(L.1, 2017) (L.1, 2016)(L.1, 2015) (66, 2014)(44, 2013), Accuracy	Continue the ongoing technical research to provide reliable data for estimating CSC in living biomass, dead organic matter, and soil organic matter.	L.1 - ARR 2022 (sent on 4. 4. 2023)	Partly implemented in the SVK NIR 2023	Partly implemented in the SVK NIR 2023, Chapter 6
LULUCF - Land representation (L.10, 2021) Transparency	Provide in the NIR an explanation for the cause of the abrupt increase in the areas of settlements and decrease in other land occurring around 1995 and report land representation data for 2016 onward.	L.5 - ARR 2022 (sent on 4. 4. 2023)	Implemented in the SVK NIR 2023	Partly implemented in the SVK NIR 2023, Chapter 6.1

## ANNEX 5. ENERGY BALANCE OF THE ŠÚ SR FOR THE YEAR 2023

ACTIVITY/FUELS	Anthracite	Coking Coal	Other Bituminous Coal	Brown Coal and Lignite	Hard Coal Coke	Brown Coal & Peat Briquettes	Patent Fuel	Coal Tar	Coke Oven Gas	Blast Furnace Gas	Oxygen Steel Furnace Gas
UNITS	<i>TJ</i>										
Primary Production	-	-	-	8 753	-	-	-	-	-	-	-
Import	464	70 548	9 889	4 756	4 561	819	56	-	-	-	-
Export	-	-	-	-	366	-	-	1 842	-	-	-
Stock Changes	283	901	-328	942	338	-	-	-	-	-	-
Gross Inland Consumption	747	71 449	9 561	14 451	4 533	819	56	-1 842	-	-	-
Transformation Input	386	71 449	3 578	13 303	46 260	-	-	-	793	1 288	306
Electricity Production - Thermal Equipment	386	-	3 578	13 303	-	-	-	-	793	1 282	291
of which: Public	386	-	2 513	13 280	-	-	-	-	-	-	-
Autoproducers	-	-	1 065	23	-	-	-	-	793	1 282	291
Nuclear Plants	-	-	-	-	-	-	-	-	-	-	-
Coke Ovens	-	58 683	-	-	-	-	-	-	-	-	-
Blast Furnaces	-	12 766	-	-	46 260	-	-	-	-	-	-
Refineries	-	-	-	-	-	-	-	-	-	-	-
Heat Production	-	-	-	-	-	-	-	-	-	6	15
Transformation Output	-	-	-	-	43 613	-	-	1 842	10 814	20 576	3 105
Electricity Production - Thermal Equipment	-	-	-	-	-	-	-	-	-	-	-
of which: Public	-	-	-	-	-	-	-	-	-	-	-
Autoproducers	-	-	-	-	-	-	-	-	-	-	-
Nuclear Plants	-	-	-	-	-	-	-	-	-	-	-
Coke Ovens	-	-	-	-	43 613	-	-	1 842	10 814	-	-
Blast Furnaces	-	-	-	-	-	-	-	-	-	20 576	3 105
Refineries	-	-	-	-	-	-	-	-	-	-	-
Heat Production	-	-	-	-	-	-	-	-	-	-	-
Exchanges and Transfers, Backflows	-	-	-	-	-	-	-	-	-	-	-
Product Transferred	-	-	-	-	-	-	-	-	-	-	-
Backflows from Petrochemical Sector	-	-	-	-	-	-	-	-	-	-	-

ACTIVITY/FUELS	Anthracite	Coking Coal	Other Bituminous Coal	Brown Coal and Lignite	Hard Coal Coke	Brown Coal & Peat Briquettes	Patent Fuel	Coal Tar	Coke Oven Gas	Blast Furnace Gas	Oxygen Steel Furnace Gas
UNITS	<i>TJ</i>										
Consumption of the Energy Sector	-	-	-	-	-	-	-	-	3 273	11 363	-
Distribution Losses	-	-	-	-	-	-	-	-	118	1 624	483

### 1<sup>st</sup> continuation

ACTIVITY/FUELS	Anthracite	Coking Coal	Other Bituminous Coal	Brown Coal and Lignite	Hard Coal Coke	Brown Coal & Peat Briquettes	Patent Fuel	Coal Tar	Coke Oven Gas	Blast Furnace Gas	Oxygen Steel Furnace Gas
UNITS	<i>TJ</i>										
Final Consumption	361	-	5 983	1 148	1 886	819	56	-	6 630	6 301	2 316
Final Non - Energy Consumption	258	-	-	-	760	-	-	-	-	-	-
of which: Chemical Industry	-	-	-	-	-	-	-	-	-	-	-
Final Energy Consumption	103	-	5 983	1 148	1 126	819	56	-	6 630	6 301	2 316
Industry	103	-	4 754	528	1 098	-	-	-	6 630	6 301	2 316
of which: Iron and steel	103	-	4 235	-	113	-	-	-	6 626	6 301	2 316
Non - ferrous metals	-	-	-	-	56	-	-	-	-	-	-
Chemical	-	-	-	-	28	-	-	-	-	-	-
Non - metallic minerals	-	-	519	482	845	-	-	-	4	-	-
Mining and quarrying	-	-	-	-	-	-	-	-	-	-	-
Food, beverages and tobacco	-	-	-	-	56	-	-	-	-	-	-
Textile and leather	-	-	-	-	-	-	-	-	-	-	-
Pulp, paper and print	-	-	-	-	-	-	-	-	-	-	-
Mach. and transport equipment	-	-	-	46	-	-	-	-	-	-	-
Not elsewhere specified	-	-	-	-	-	-	-	-	-	-	-
Transport	-	-	-	-	-	-	-	-	-	-	-
Other Sectors	-	-	1 229	620	28	819	56	-	-	-	-
of which: Households	-	-	300	402	28	567	-	-	-	-	-
Agriculture	-	-	-	-	-	-	-	-	-	-	-
Commercial and public services	-	-	929	218	-	252	56	-	-	-	-

2<sup>nd</sup> continuation

ACTIVITY/FUELS	Natural Gas	Crude Oil and NGL	Refinery Feedstock <sup>1</sup>	Refinery Gas	LPG	Naphtha	Gasoline	Kerosene
UNITS	<i>TJ</i>							
Primary Production	1 752	84	8 553	-	-	-	-	-
Import	159 699	227 561	-	-	2 392	880	9 794	87
Export	-	42	-	-	2 760	4 488	28 372	2 771
Stock Changes	-10 319	-8 022	-	-	-92	704	-615	-87
Gross Inland Consumption	151 132	219 581	8 553	-	-460	-2 904	-19 193	-2 771
Transformation Input	30 593	219 581	30 630	167	-	-	-	-
Electricity Production - Thermal Equipment	22 781	-	-	167	-	-	-	-
of which: Public	21 811	-	-	-	-	-	-	-
Autoproducers	970	-	-	167	-	-	-	-
Nuclear Plants	-	-	-	-	-	-	-	-
Coke Ovens	-	-	-	-	-	-	-	-
Blast Furnaces	-	-	-	-	-	-	-	-
Refineries	-	219 581	30 630	-	-	-	-	-
Heat Production	7 812	-	-	-	-	-	-	-
Transformation Output	-	-	-	14 815	7 820	25 300	46 247	3 984
Electricity Production - Thermal Equipment	-	-	-	-	-	-	-	-
of which: Public	-	-	-	-	-	-	-	-
Autoproducers	-	-	-	-	-	-	-	-
Nuclear Plants	-	-	-	-	-	-	-	-
Coke Ovens	-	-	-	-	-	-	-	-
Blast Furnaces	-	-	-	-	-	-	-	-
Refineries	-	-	-	14 815	7 820	25 300	46 247	3 984
Heat Production	-	-	-	-	-	-	-	-
Exchanges and Transfers, Backflows	-6 609	-	22 077	-	-2 668	-5 632	-	-
Product Transferred	-6 609	-	13 777	-	-	-	-	-
Backflows from Petrochemical Sector	-	-	8 300	-	-2 668	-5 632	-	-



ACTIVITY/FUELS	Natural Gas	Crude Oil and NGL	Refinery Feedstock <sup>1</sup>	Refinery Gas	LPG	Naphtha	Gasoline	Kerosene
UNITS	<i>TJ</i>							
Consumption of the Energy Sector	3 200	-	-	10 993	-	-	-	-
Distribution Losses	2 849	-	-	-	-	-	-	-

### 3<sup>rd</sup> continuation

ACTIVITY/FUELS	Natural Gas	Crude Oil and NGL	Refinery Feedstock <sup>1</sup>	Refinery Gas	LPG	Naphtha	Gasoline	Kerosene
UNITS	<i>TJ</i>							
Final Consumption	106 460	-	-	3 655	4 692	16 764	27 054	1 213
Final Non - Energy Consumption	12 553	-	-	-	2 208	16 764	-	-
of which: Chemical Industry	12 553	-	-	-	2 208	16 764	-	-
Final Energy Consumption	93 907	-	-	3 655	2 484	-	27 054	1 213
Industry	30 410	-	-	3 655	368	-	44	-
of which: Iron and steel	5 879	-	-	-	-	-	44	-
Non - ferrous metals	1 262	-	-	-	-	-	-	-
Chemical	3 594	-	-	3 655	-	-	-	-
Non - metallic minerals	3 246	-	-	-	92	-	-	-
Mining and quarrying	1 040	-	-	-	-	-	-	-
Food, beverages and tobacco	4 811	-	-	-	92	-	-	-
Textile and leather	319	-	-	-	-	-	-	-
Pulp, paper and print	1 444	-	-	-	-	-	-	-
Mach. and transport equipment	5 666	-	-	-	46	-	-	-
Not elsewhere specified	3 149	-	-	-	138	-	-	-
Transport	313	-	-	-	1 610	-	27 010	1213
Other Sectors	63 184	-	-	-	506	-	-	-
of which: Households	43 949	-	-	-	276	-	-	-
Agriculture	1 223	-	-	-	138	-	-	-
Commercial and public services	18 012	-	-	-	92	-	-	-

<sup>1</sup> include Additives, Oxygenates and Other Hydrocarbons

4<sup>th</sup> continuation

ACTIVITY/FUELS	Natural Gas	Crude Oil and NGL	Refinery Feedstock <sup>1</sup>	Refinery Gas	LPG	Naphtha	Gasoline	Kerosene
UNITS	<i>mil. m<sup>3</sup></i>	<i>1 000 t</i>	<i>1 000 t</i>	<i>1 000 t</i>	<i>1 000 t</i>	<i>1 000 t</i>	<i>1 000 t</i>	<i>1 000 t</i>
Final Consumption	3 011	-	-	131	102	381	616	28
Final Non - Energy Consumption	355	-	-	-	48	381	-	-
of which: Chemical Industry	355	-	-	-	48	381	-	-
Final Energy Consumption	2 656	-	-	131	54	-	616	28
Industry	860	-	-	131	8	-	1	-
of which: Iron and steel	166	-	-	-	-	-	1	-
Non - ferrous metals	36	-	-	-	-	-	-	-
Chemical	102	-	-	131	-	-	-	-
Non - metallic minerals	92	-	-	-	2	-	-	-
Mining and quarrying	29	-	-	-	-	-	-	-
Food, beverages and tobacco	136	-	-	-	2	-	-	-
Textile and leather	9	-	-	-	-	-	-	-
Pulp, paper and print	41	-	-	-	-	-	-	-
Mach. and transport equipment	160	-	-	-	1	-	-	-
Not elsewhere specified	89	-	-	-	3	-	-	-
Transport	9	-	-	-	35	-	615	28
Other Sectors	1 787	-	-	-	11	-	-	-
of which: Households	1 244	-	-	-	6	-	-	-
Agriculture	34	-	-	-	3	-	-	-
Commercial and public services	509	-	-	-	2	-	-	-

<sup>1</sup> include Additives, Oxygenates and Other Hydrocarbons

5<sup>th</sup> continuation

ACTIVITY/FUELS	Diesel Oil	Light Fuel Oil	HFO - low Sulphur (<1%)	HFO - high Sulphur (>=1%)	White Spirit SBP	Lubricants	Bitumen	Paraffin Waxes	Petroleum Coke	Other Products
UNITS	<i>TJ</i>									
Primary Production	-	-	-	-	-	-	-	-	-	-
Import	55 619	1 261	162	2 666	387	2 459	8 310	173	2 583	5 750
Export	86 584	1 058	4 242	8 685	215	375	3 544	-	-	13 784
Stock Changes	1 346	-81	-81	-566	-	-	-	-	-276	-381
Gross Inland Consumption	-29 619	122	-4 161	-6 585	172	2 084	4 766	173	2 307	-8 415
Transformation Input	-	-	81	2 909	-	-	-	-	-	-
Electricity Production - Thermal Equipment	-	-	81	2 909	-	-	-	-	-	-
of which: Public	-	-	81	-	-	-	-	-	-	-
Autoproducers	-	-	-	2 909	-	-	-	-	-	-
Nuclear Plants	-	-	-	-	-	-	-	-	-	-
Coke Ovens	-	-	-	-	-	-	-	-	-	-
Blast Furnaces	-	-	-	-	-	-	-	-	-	-
Refineries	-	-	-	-	-	-	-	-	-	-
Heat Production	-	-	-	-	-	-	-	-	-	-
Transformation Output	111 785	1 058	4 242	16 685	-	-	-	-	1 619	11 712
Electricity Production - Thermal Equipment	-	-	-	-	-	-	-	-	-	-
of which: Public	-	-	-	-	-	-	-	-	-	-
Autoproducers	-	-	-	-	-	-	-	-	-	-
Nuclear Plants	-	-	-	-	-	-	-	-	-	-
Coke Ovens	-	-	-	-	-	-	-	-	-	-
Blast Furnaces	-	-	-	-	-	-	-	-	-	-
Refineries	111 785	1 058	4 242	16 685	-	-	-	-	1 619	11 712
Heat Production	-	-	-	-	-	-	-	-	-	-
Exchanges and Transfers, Backflows	-	-	-	-	-	-	-	-	-	-
Product Transferred	-	-	-	-	-	-	-	-	-	-
Backflows from Petrochemical Sector	-	-	-	-	-	-	-	-	-	-
Consumption of the Energy Sector	-	-	-	-	-	-	-	-	1 618	-
Distribution Losses	-	-	-	-	-	-	-	-	-	-

### 6<sup>th</sup> continuation

ACTIVITY/FUELS	Diesel Oil	Light Fuel Oil	HFP - low sulphur (<1%)	HFO - high Sulphur (>=1%)	White Spirit SBP	Lubricants	Bitumen	Paraffin Waxes	Petroleum Coke	Other Products
UNITS	<i>TJ</i>									
Final Consumption	82 166	1 180	-	7 191	172	2 084	4 766	173	2 308	3 297
Final Non - Energy Consumption	-	936	-	-	172	2 084	4 766	173	620	3 297
of which: Chemical Industry	-	936	-	-	-	-	-	-	-	3 297
Final Energy Consumption	82 166	244	-	7 191	-	-	-	-	1 688	-
Industry	420	163	-	7 191	-	-	-	-	1 688	-
of which: Iron and steel	-	-	-	-	-	-	-	-	-	-
Non - ferrous metals	-	-	-	-	-	-	-	-	-	-
Chemical	-	-	-	7 191	-	-	-	-	-	-
Non - metallic minerals	42	-	-	-	-	-	-	-	1 688	-
Mining and quarrying	126	-	-	-	-	-	-	-	-	-
Food, beverages and tobacco	42	122	-	-	-	-	-	-	-	-
Textile and leather	-	-	-	-	-	-	-	-	-	-
Pulp, paper and print	-	-	-	-	-	-	-	-	-	-
Mach. and transport equipment	-	-	-	-	-	-	-	-	-	-
Not elsewhere specified	210	41	-	-	-	-	-	-	-	-
Transport	79 769	-	-	-	-	-	-	-	-	-
Other Sectors	1 977	81	-	-	-	-	-	-	-	-
of which: Households	-	-	-	-	-	-	-	-	-	-
Agriculture	1 977	-	-	-	-	-	-	-	-	-
Commercial and public services	-	81	-	-	-	-	-	-	-	-

### 7<sup>th</sup> continuation

ACTIVITY/FUELS	Nuclear Heat	Solar Heat	Geoth. Heat	Ambient heat	Heat	Wood and Charcoal	MSW	Bio-gas	ISW	Wind energy	Hydro Energy	Solar Electricity	EE	Liquid Bio-fuels	Total
UNITS	<i>TJ</i>														
Primary Production	177 266	405	366	4 721	-	48 857	2 971	4 461	6 683	14	16 870	2 178	-	7 745	291 679
Import	-	-	-	-	61	259	-	-	490	-	-	-	38 336	3 637	613 659
Export	-	-	-	-	-	373	-	-	-	-	-	-	50 656	4 251	214 408

ACTIVITY/FUELS	Nuclear Heat	Solar Heat	Geoth. Heat	Ambient heat	Heat	Wood and Charcoal	MSW	Bio-gas	ISW	Wind energy	Hydro Energy	Solar Electricity	EE	Liquid Bio-fuels	Total
UNITS	<i>TJ</i>														
Stock Changes	-	-	-	-	-	-270	-	-	80	-	-	-	-	37	-16 487
Gross Inland Consumption	177 266	405	366	4 721	61	48 473	2 971	4 461	7 253	14	16 870	2 178	-12 320	7 168	674 443
Transformation Input	175 483	-	334	-	-	16 627	1 765	3 554	536	-	-	-	-	-	619 623
Electricity Production - Thermal Equipment	-	-	-	-	-	13 674	1 765	3 479	517	-	-	-	-	-	65 006
of which: Public	-	-	-	-	-	7 772	-	968	318	-	-	-	-	-	47 129
Autoproducers	-	-	-	-	-	5 902	1 765	2 511	199	-	-	-	-	-	17 877
Nuclear Plants	175 483	-	-	-	-	-	-	-	-	-	-	-	-	-	175 483
Coke Ovens	-	-	-	-	-	-	-	-	-	-	-	-	-	-	58 683
Blast Furnaces	-	-	-	-	-	-	-	-	-	-	-	-	-	-	59 026
Refineries	-	-	-	-	-	-	-	-	-	-	-	-	-	-	250 211
Heat Production	-	-	334	-	-	2 953	-	75	19	-	-	-	-	-	11 214
Transformation Output	-	-	-	-	24 781	-	-	-	-	-	-	-	87 358	-	437 356
Electricity Production - Thermal Equipment	-	-	-	-	15 064	-	-	-	-	-	-	-	21 359	-	36 423
of which: Public	-	-	-	-	12 952	-	-	-	-	-	-	-	13 129	-	26 081
Autoproducers	-	-	-	-	2 112	-	-	-	-	-	-	-	8 230	-	10 342
Nuclear Plants	-	-	-	-	-	-	-	-	-	-	-	-	65 999	-	65 999
Coke Ovens	-	-	-	-	-	-	-	-	-	-	-	-	-	-	56 269
Blast Furnaces	-	-	-	-	-	-	-	-	-	-	-	-	-	-	23 681
Refineries	-	-	-	-	-	-	-	-	-	-	-	-	-	-	245 267
Heat Production	-	-	-	-	9 717	-	-	-	-	-	-	-	-	-	9 717
Exchanges and Transfers, Backflows	-1 783	-2	-	-4 721	6 506	-	-	-	-	-14	-16 870	-2 178	19 062	-7 168	0
Product Transferred	-1 783	-2	-	-4 721	6 506	-	-	-	-	-14	-16 870	-2 178	19 062	-7 168	0
Backflows from Petrochemical Sector	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0
Consumption of the Energy Sector	-	2	-	-	2 835	-	-	-	-	-	-	-	13 496	-	46 780
Distribution Losses	-	-	-	-	3 923	11	-	-	-	-	-	-	4 021	-	13 029

8<sup>th</sup> continuation

ACTIVITY/FUELS	Nuclear Heat	Solar Heat	Geoth. Heat	Ambient heat	Heat	Wood and Charcoal	MSW	Bio-gas	ISW	Wind energy	Hydro Energy	Solar Electricity	EE	Liquid Bio-fuels	Total
UNITS	<i>TJ</i>														
Final Consumption	-	401	32	-	24 590	31 835	1 206	907	6 717	-	-	-	76 583	-	430 946
Final Non - Energy Consumption	-	-	-	-	-	-	-	-	-	-	-	-	-	-	44 591
of which: Chemical Industry	-	-	-	-	-	-	-	-	-	-	-	-	-	-	35 758
Final Energy Consumption	-	401	32	-	24 590	31 835	1 206	907	6 717	-	-	-	76 583	-	386 355
Industry	-	-	-	-	2 244	12 812	-	38	6 704	-	-	-	31 871	-	119 338
of which: Iron and steel	-	-	-	-	25	39	-	-	-	-	-	-	5 980	-	31 661
Non - ferrous metals	-	-	-	-	74	-	-	-	-	-	-	-	749	-	2 141
Chemical	-	-	-	-	320	3	-	-	950	-	-	-	3 373	-	19 114
Non - metallic minerals	-	-	-	-	201	95	-	-	5 724	-	-	-	2 376	-	15 314
Mining and quarrying	-	-	-	-	2	-	-	-	-	-	-	-	212	-	1 380
Food, beverages and tobacco	-	-	-	-	72	208	-	11	-	-	-	-	2 203	-	7 617
Textile and leather	-	-	-	-	26	1	-	-	-	-	-	-	497	-	843
Pulp, paper and print	-	-	-	-	974	9 876	-	3	-	-	-	-	3 064	-	15 361
Mach. and transport equipment	-	-	-	-	361	213	-	24	30	-	-	-	9 320	-	15 706
Not elsewhere specified	-	-	-	-	189	2 377	-	-	-	-	-	-	4 097	-	10 201
Transport	-	-	-	-	-	-	-	-	-	-	-	-	2 542	-	112 457
Other Sectors	-	401	32	-	22 346	19 023	1206	869	13	-	-	-	42 170	-	154 560
of which: Households	-	321	-	-	15 987	18 666	-	-	-	-	-	-	21 053	-	101 549
Agriculture	-	-	32	-	68	278	-	554	-	-	-	-	727	-	4 997
Commercial and public services	-	80	-	-	6 291	79	1206	315	13	-	-	-	20 390	-	48 014

---

Title: **NATIONAL INVENTORY DOCUMENT OF THE SLOVAK REPUBLIC 2025**

Issued: SLOVAK HYDROMETEOROLOGICAL INSTITUTE

Authors: JURAJ LABOVSKÝ (CHAPTER 3)  
JÁN HORVÁTH (CHAPTER 3)  
VLADIMÍR DANIELIK (CHAPTERS 3, 4)  
KRISTÍNA TONHAUZER (CHAPTERS 5, 7)  
ONDREJ PASTIERIK (CHAPTER 5)  
ZUZANA PALKOVIČOVÁ (CHAPTER 5)  
TIBOR PRIWITZER (CHAPTERS 6, 11)  
IVAN BARKA (CHAPTERS 6, 11)  
PAVEL PAVLENDÁ (CHAPTERS 6, 11)  
MICHAL SVIČEK (CHAPTER 6)  
PAVOL BEZÁK (CHAPTER 6)  
KRISTÍNA BUCHOVÁ (CHAPTER 6)  
ŠTEFAN POLLÁK (CHAPTER 6)  
IGOR BODÍK (CHAPTER 7)  
MAREK HRABČÁK (CHAPTER 7)  
ZUZANA JONÁČEK (CHAPTER 7)

Editors JANKA SZEMESOVÁ  
ALEXANDRA NADŽADYOVÁ  
PETRA KRŠÁKOVÁ  
LENKA ZETOCHOVÁ  
PATRÍCIA NAVRATILOVÁ

Date MARCH 2025

Version 2

Pages 503

ISBN 978-80-99929-74-7

EAN 9788099929747

